

Ichthyotic Keratoderma, Spasticity, Hypomyelination and Dysmorphic Facial Features (IKSHD): A Comprehensive Disease Report

Disease: Ichthyotic Keratoderma, Spasticity, Hypomyelination And Dysmorphic Facial Features (IKSHD) **Causal gene:** *ELOVL1* (ELOVL fatty acid elongase 1) **OMIM phenotype:** #618527 | **OMIM gene:** 611813 | *HGNC:* 14417 | *Locus:* 1p34.2 *Suggested MONDO mapping:** MONDO term for "ichthyotic keratoderma, spasticity, hypomyelination, and dysmorphic facial features"

Summary

Ichthyotic Keratoderma, Spasticity, Hypomyelination and Dysmorphic Facial Features (IKSHD) is an ultra-rare (<1/1,000,000) congenital, multisystem disorder that sits at the interface of dermatology and neurology. It is caused by pathogenic variants in *ELOVL1*, the gene encoding the endoplasmic-reticulum (ER) fatty-acid elongase that performs the **first and rate-limiting condensation step** of very-long-chain fatty acid (VLCFA) biosynthesis. The classic autosomal-dominant form of IKSHD arises from a **recurrent de novo heterozygous missense variant, c.494C>T (p.Ser165Phe)**, identified independently in multiple unrelated pediatric probands without a family history of disease ([PMID: 29496980](#); [PMID: 30487246](#); [PMID: 42101073](#)).

Mechanistically, IKSHD is a **VLCFA-biosynthesis deficiency**. ELOVL1 elongates saturated and monounsaturated C22–C26 VLCFAs, which are esterified into ceramides (essential for the epidermal permeability barrier) and myelin sphingolipids (essential for central white-matter integrity). Loss of ELOVL1 function depletes these VLCFA-containing complex lipids, producing the two-compartment clinical signature: **ichthyotic keratoderma** in skin and **hypomyelination with progressive spasticity** in the central nervous system, accompanied by

dysmorphic facial features and nystagmus. Notably, IKSHD is the biochemical *mirror image* of VLCFA-catabolism leukodystrophies such as X-linked adrenoleukodystrophy (X-ALD), where VLCFAs *accumulate*; both nonetheless converge on white-matter pathology and spasticity.

A defining feature of the *ELOVL1* disease spectrum is a **graded, allele-dose-dependent severity**. Monoallelic de novo p.Ser165Phe produces classic IKSHD with *mild* cerebral hypomyelination, whereas **biallelic (recessive) variants** cause a more severe and recognizable hypomyelinating leukodystrophy with a complex movement disorder (head tremor, myoclonus, dysarthria), 100%-penetrant developmental delay and spasticity, and corpus callosum hypoplasia ([PMID: 40590574](#)). No disease-specific or *ELOVL1*-targeted therapy exists; management is symptomatic and multidisciplinary — topical emollients, keratolytics and retinoids for the skin, and antispasticity agents plus physical/occupational/speech therapy for the neurological features.

1. Disease Information

Overview. IKSHD is a congenital multisystem genodermatosis-plus-leukodystrophy characterized by the tetrad encoded in its name: (1) **ichthyotic keratoderma** (generalized scaling with palmoplantar involvement), (2) **spasticity** (upper-motor-neuron signs, progressive), (3) **hypomyelination** (deficient central white-matter myelin on MRI), and (4) **dysmorphic facial features**. Additional recurrent features include **nystagmus** and developmental delay. The disorder was delineated in 2018 when whole-exome sequencing of two unrelated pediatric probands with this shared phenotype identified a novel heterozygous *ELOVL1* variant ([PMID: 29496980](#)).

Key identifiers.

Resource	Identifier
OMIM (phenotype)	#618527
OMIM (gene <i>ELOVL1</i>)	*611813
HGNC	14417
Gene locus	1p34.2
Orphanet	Ultra-rare; listed under ELOVL1-related disorders (specific ORPHAcode as assigned)
ICD-11	Best mapped under hereditary ichthyoses / hereditary spastic conditions (no dedicated code)
MeSH	No dedicated MeSH descriptor; indexed via <i>ELOVL1</i> , "Ichthyosis," "Hypomyelination"
MONDO	Map to the ELOVL1-related IKSHD phenotype entry

Synonyms / alternative names. IKSHD; ELOVL1-related ichthyosis–spasticity–hypomyelination disorder; ELOVL1-related neuroichthyosis. The biallelic form is described as "hypomyelinating leukodystrophy, movement disorder, and ichthyosis" ([PMID: 40590574](#)).

Nature of information. Evidence is derived from **individual patient reports and small case series** (aggregated at the disease level via OMIM/Orphanet), not from large EHR cohorts — consistent with an ultra-rare Mendelian disorder.

2. Etiology

Primary cause (genetic). IKSHD is a **monogenic disorder** caused by pathogenic variants in *ELOVL1*. The classic dominant form results from a **recurrent de novo heterozygous missense variant p.Ser165Phe (c.494C>T)**; a more severe form results from **biallelic** variants. There is no environmental or infectious etiology.

Genetic risk factors. The causal variant is the risk factor. The recurrent p.Ser165Phe change is the principal reported dominant allele; biallelic loss-of-function/hypomorphic variants define the recessive end. Because ELOVL1 sits within a network of six mammalian ELOVL elongases

(ELOVL1–7), **other ELOVL genes are candidate modifiers**: patient fibroblasts show dysregulation of other ELOVL enzymes, which may buffer or aggravate the VLCFA deficit (PMID: 40590574).

Environmental risk factors. None established. As a de novo–dominant disorder, advanced parental age is a theoretical (unproven) contributor to de novo variant occurrence. Sex distribution is approximately 1:1.

Protective factors. None established genetically or environmentally. Mechanistically, residual/compensatory activity of paralogous elongases (ELOVL3, ELOVL6, ELOVL7) that also handle VLCFA substrates could partially mitigate severity, but this is inferred, not demonstrated.

Gene–environment interactions. No specific GxE interactions have been documented. Skin-barrier compromise means that **environmental factors affecting congenital ichthyosis generally** (low humidity, temperature extremes, infection risk through a defective barrier) can modulate cutaneous morbidity, but these are downstream and non-specific.

3. Phenotypes

IKSHD is a two-compartment disorder (skin + CNS) with facial dysmorphism. Frequencies below are best-characterized for the **biallelic** cohort (n=7; PMID: 40590574); the monoallelic form has similar features but *milder* hypomyelination.

Phenotype	Type	HPO suggestion	Onset	Frequency (biallelic cohort)	Severity/course
Ichthyotic keratoderma / ichthyosis	Physical/skin sign	HP:0008064 (Ichthyosis); HP:0007479 (palmoplantar keratoderma)	Congenital/neonatal	5/7 (71%)	Chronic, lifelong
Developmental delay	Behavioral/cognitive	HP:0001263	Infancy	7/7 (100%)	Stable–progressive
Progressive spasticity	Neurological sign	HP:0001257 (Spasticity); HP:0002061 (lower-limb spasticity)	Childhood	7/7 (100%)	Progressive
Head tremor	Neurological sign	HP:0002346	Childhood	7/7 (100%)	Persistent
Dysarthria	Neurological sign	HP:0001260	Childhood	6/6 (100%)	Progressive
Myoclonus	Neurological sign	HP:0001336	Childhood	6/7 (86%)	Episodic/persistent
Nystagmus	Neuro-ophthalmic sign	HP:0000639	Infancy/childhood	5/6 (83%)	Persistent
Hypomyelination (MRI)	Imaging/lab abnormality	HP:0002500; HP:0006808 (leukoencephalopathy)	Congenital	6/6 (100%)	Non-progressive on MRI
Corpus callosum hypoplasia	Imaging abnormality	HP:0002079	Congenital	5/6 (83%)	Static
Dysmorphic facial features	Physical sign	HP:0001999	Congenital	Recurrent in IKSHD	Static

Quality-of-life impact. The combination of progressive spasticity, a complex movement disorder, dysarthria, and developmental delay imposes substantial motor and communication disability, likely requiring mobility aids, assistive communication, and long-term caregiver support. The ichthyotic skin adds chronic pruritus/xerosis, cosmetic burden, thermoregulatory and barrier-infection concerns. Disease-specific QoL instruments have not been reported; generic pediatric-disability and dermatologic-QoL tools would apply.

4. Genetic / Molecular Information

Causal gene. *ELOVL1* (ELOVL fatty acid elongase 1; HGNC:14417; OMIM 611813; locus 1p34.2). *ELOVL1* catalyzes elongation of saturated and monounsaturated C22–C26 VLCFAs*, the rate-limiting condensation step of the ER elongation cycle (PMID: 30487246; PMID: 16564093).

Pathogenic variants.

Variant	Nomenclature	Zygosity/ inheritance	Classification	Type	Disease
p.Ser165Phe	c.494C>T	Heterozygous, de novo (AD)	Pathogenic	Missense	Classic IKSHD (#618527)
Biallelic <i>ELOVL1</i> variants	(various)	Homozygous/ compound- het (AR)	Pathogenic/ likely pathogenic	Missense + LoF/ hypomorphic	Severe hypomyelinating leukodystrophy + ichthyosis

- **Allele frequency.** The recurrent p.Ser165Phe is de novo and absent/ultra-rare in population databases (gnomAD), consistent with a highly penetrant pathogenic change.
- **Somatic vs germline.** Germline (de novo in dominant cases; inherited in recessive cases).
- **Functional consequence.** The biochemical readout is **loss of elongase output** — reduced plasma C24:0 and C26:0 in a biallelic patient, and reduced VLCFA elongation in patient fibroblasts (PMID: 40590574). The dominant p.Ser165Phe likely acts by impairing enzyme activity (loss-of-function/dominant effect on the elongation complex); a dominant-negative contribution on the shared ER four-enzyme complex is plausible but not definitively established.

Modifier genes. Paralogous elongases (**ELOVL3, ELOVL6, ELOVL7**) and downstream ceramide-synthesis machinery are candidate modifiers; fibroblast studies show compensatory dysregulation of other ELOVLs ([PMID: 40590574](#)).

Epigenetic information. No disease-specific DNA-methylation or histone-modification signature has been reported for IKSHD.

Chromosomal abnormalities. None; IKSHD is a single-nucleotide/small-variant disorder, not a copy-number or structural disorder.

5. Environmental Information

IKSHD is a **purely genetic** disorder. No **environmental toxins, radiation, pollution, occupational exposures, lifestyle factors, or infectious agents** cause or trigger it. Environmental exposures are relevant only as non-specific modifiers of the skin-barrier phenotype (e.g., dry/cold climates worsening scaling; barrier breaches raising cutaneous infection risk), analogous to other congenital ichthyoses.

6. Mechanism / Pathophysiology

Ordered causal chain

1. A pathogenic *ELOVL1* variant (heterozygous de novo **p.Ser165Phe**, or **biallelic** variants) **leads to** reduced ELOVL1 elongase activity in the endoplasmic reticulum.
2. Reduced ELOVL1 activity **results in** impaired condensation — the first, rate-limiting step of the four-enzyme ER VLCFA-elongation cycle — **causing** decreased synthesis of saturated/monounsaturated **C22–C26+ very-long-chain fatty acids** (measurable as reduced plasma C24:0 and C26:0). ([PMID: 39946831](#); [PMID: 40590574](#))
3. VLCFA deficiency **branches** into two tissue-specific consequences:
4. **Skin branch:** Reduced VLCFAs **lead to** depletion of ultra-long-chain **ceramides** and diminished stratum-corneum lipid lamellae, **causing** a defective epidermal permeability barrier that manifests as **ichthyotic keratoderma** (inferred from *Elovl1*-knockout mouse: barrier failure and reduced $\geq$ C26 ceramides; [PMID: 23689133](#)).


5. **CNS branch:** Reduced VLCFAs **lead to** shortened-chain **myelin sphingolipids, causing** hypomyelination and, consequently, progressive **spasticity** and a complex movement disorder (supported by Elov11-deficient mouse: reduced myelin-sphingolipid chain length and impaired motor coordination; [PMID: 32123819](#)).
6. Deficient central myelination and cortico-spinal tract dysfunction **result in** the observed upper-motor-neuron signs (spasticity, hyperreflexia), dysarthria, nystagmus, head tremor, and myoclonus.
7. Allele dose **modulates** severity: monoallelic p.Ser165Phe → *mild* hypomyelination (classic IKSHD); biallelic variants → *more severe* hypomyelinating leukodystrophy with corpus callosum hypoplasia and 100%-penetrant motor/developmental involvement (inferred from genotype–phenotype comparison; [PMID: 42101073](#); [PMID: 40590574](#)).

Detail by category

- **Molecular pathways.** VLCFA/sphingolipid biosynthesis (ER fatty-acid elongation cycle: condensation → reduction → dehydration → reduction). ELOVL1 catalyzes the **rate-limiting condensation** step ([PMID: 39946831](#)). Downstream: ceramide synthesis (skin) and myelin sphingolipid assembly (CNS).
- **Cellular processes.** Epidermal terminal differentiation / cornification (keratinocytes); oligodendrocyte myelination (CNS). Membrane biogenesis broadly — VLCFAs are esterified into sphingolipids, glycerophospholipids and ether lipids ([PMID: 39946831](#)).
- **Protein dysfunction.** Loss of elongase catalytic output; possible ER misfolding/dominant effects on the shared elongation complex (inferred). ELOVL defects are noted to contribute to disorders at the metabolic–neurodegenerative interface partly via misfolded enzymes in the ER/Golgi ([PMID: 39946831](#)).
- **Metabolic changes.** Reduced VLCFA/sphingolipid pool (C24:0, C26:0 down). This is the **opposite** of VLCFA-catabolism disorders (X-ALD), where ABCD1 loss impairs peroxisomal VLCFA degradation and VLCFAs *accumulate* ([PMID: 42320861](#); [PMID: 42469918](#)).
- **Biochemical abnormalities.** Enzyme (elongase) deficiency; reduced ceramide chain length in skin (Elov11-KO mouse) and reduced myelin sphingolipid chain length in brain.
- **Immune involvement.** Not a primary feature; no autoimmunity/immunodeficiency reported (barrier failure may secondarily raise infection risk).
- **Tissue-damage mechanisms.** Primarily a *developmental/biosynthetic deficiency* (dysmyelination/hypomyelination and defective cornification) rather than degenerative injury such as oxidative stress or fibrosis.

Ontology suggestions. GO biological processes: **fatty acid elongation** (GO:0030497), **very long-chain fatty acid metabolic process** (GO:0000038), **sphingolipid biosynthetic process** (GO:0030148), **myelination** (GO:0042552), **keratinocyte differentiation** (GO:0030216), **establishment of skin barrier** (GO:0061436). GO cellular component: **endoplasmic reticulum membrane** (GO:0005789). CL cell types: **keratinocyte** (CL:0000312), **oligodendrocyte** (CL:0000128). CHEBI: **very long-chain fatty acid** (CHEBI:27283), **ceramide** (CHEBI:17761).

Mechanistic diagram



7. Anatomical Structures Affected

- **Organ level.** Primary: **skin** (integumentary system) and **brain/central white matter** (nervous system). Secondary: eyes/oculomotor pathways (nystagmus). Body systems: **integumentary** and **central nervous** systems predominate.
- **Tissue/cell level.** **Epidermis** — keratinocytes/stratum corneum (CL:0000312). **CNS white matter** — oligodendrocytes and myelin sheaths (CL:0000128). Corpus callosum hypoplasia indicates commissural white-matter involvement.
- **Subcellular level.** **Endoplasmic reticulum membrane** (GO:0005789) — site of the ELOVL1 elongation complex; downstream sphingolipid-rich plasma/myelin membranes.

- **Localization (UBERON).** Skin/epidermis (UBERON:0001003 / UBERON:0001834), palms and soles (palmoplantar keratoderma), brain white matter (UBERON:0002316), corpus callosum (UBERON:0002336). Involvement is **bilateral/generalized** (skin) and **symmetric** (central white matter).

8. Temporal Development

- **Onset. Congenital / neonatal** for ichthyosis and hypomyelination; neurological signs (spasticity, movement disorder, developmental delay) emerge and evolve through **infancy and childhood**. Onset pattern is **chronic/insidious**.
- **Progression.** Cutaneous disease is **chronic and lifelong**. **Spasticity is progressive**; developmental delay is 100% penetrant in the biallelic cohort. Notably, brain MRI **hypomyelination is described as non-progressive** (PMID: 40590574) — i.e., a static hypomyelination rather than an actively demyelinating course, even as clinical motor signs progress.
- **Disease course pattern.** Progressive neuromotor disability on a background of static white-matter hypomyelination; chronic skin disease.
- **Duration.** Lifelong.
- **Remission / critical periods.** No spontaneous remission. Given that myelination is a developmental process, an **early developmental window** would be the theoretical opportunity for any future disease-modifying (e.g., substrate/lipid-supplementation) intervention — inferred, not demonstrated.

9. Inheritance and Population

- **Epidemiology.** Ultra-rare, prevalence <1/1,000,000. Only a small number of patients are reported worldwide (two founding probands plus subsequent case reports for the dominant form; a 7-patient series for the biallelic form). Precise incidence/prevalence figures are not established.
- **Inheritance patterns.**
- **Autosomal dominant, de novo:** classic IKSHD via recurrent p.Ser165Phe (no family history in probands) (PMID: 29496980; PMID: 42101073).

- **Autosomal recessive:** biallelic variants → severe hypomyelinating leukodystrophy ([PMID: 40590574](#)).
- **Penetrance / expressivity.** High/complete penetrance for the reported pathogenic genotypes; in the biallelic cohort, developmental delay, spasticity, and head tremor were **100% penetrant**. Expressivity is graded by allele dose (mono → milder; biallelic → more severe).
- **Genetic anticipation / mosaicism / founder effects.** Not reported; not a repeat-expansion disorder. Germline mosaicism theoretically possible for de novo cases but undocumented. Consanguinity is relevant for the recessive form (as for AR disorders generally).
- **Carrier frequency.** Not established; expected very low given ultra-rarity.
- **Demographics.** No ethnic predilection established. **Sex ratio ~1:1.** Age distribution: pediatric-onset; affected individuals present from birth/infancy.

10. Diagnostics

- **Clinical recognition.** The **combination of congenital ichthyosis + spasticity + MRI hypomyelination + dysmorphic facial features (± nystagmus)** should prompt consideration of IKSHD/ELOVL1-related disease.
- **Laboratory / biomarkers.** **Plasma VLCFA profiling** is the key biochemical test: IKSHD shows **reduced** C24:0 and C26:0 (a *deficiency* pattern), which distinguishes it from X-ALD's VLCFA *accumulation* ([PMID: 40590574](#); [PMID: 42320861](#)). Fibroblast elongation assays and stable-isotope substrate assays confirm reduced ELOVL1 activity ([PMID: 30487246](#)).
- **Imaging.** **Brain MRI** demonstrating **hypomyelination** (± corpus callosum hypoplasia) is central; hypomyelination is non-progressive on serial imaging.
- **Genetic testing (definitive).** **Whole-exome (WES) or whole-genome sequencing (WGS)** is the diagnostic method of choice, given IKSHD was defined by WES ([PMID: 29496980](#)). Targeted single-gene/panel testing for *ELOVL1* (in ichthyosis, leukodystrophy, or hereditary-spastic panels) can confirm the recurrent **c.494C>T (p.Ser165Phe)** or biallelic variants. Chromosomal microarray/karyotype/FISH/mtDNA/repeat-expansion testing are **not applicable** (single-nucleotide disorder).
- **Skin biopsy.** May show ichthyosis-consistent histology (hyperkeratosis); not specific.
- **Differential diagnosis.** X-linked adrenoleukodystrophy and other VLCFA-related leukodystrophies (distinguished by VLCFA *accumulation* vs *deficiency*); other congenital

ichthyoses with neurological involvement (e.g., Sjögren–Larsson syndrome — ALDH3A2; MEDNIK; trichothiodystrophy; Chanarin–Dorfman); other hypomyelinating leukodystrophies (e.g., PLP1-related). Genetic testing resolves these.

- **Screening.** No newborn-screening program exists for IKSHD. Cascade genetic testing is relevant in recessive families. (X-ALD newborn screening, based on C26:0-lysoPC elevation, will *not* detect ELOVL1 deficiency, which lowers VLCFAs.)

11. Outcome / Prognosis

- **Survival/mortality.** No formal survival statistics are published. Unlike neonatal-lethal complete Elov11 loss in mice (which die from barrier failure; [PMID: 23689133](#)), human patients survive into childhood and beyond, indicating residual elongase function. Life expectancy is not well defined but the disorder is chronic rather than rapidly fatal.
- **Morbidity/function.** Substantial: progressive spasticity, complex movement disorder, dysarthria, developmental delay, and chronic skin disease produce combined motor, communication, and cognitive disability. Many patients likely require mobility and communication support.
- **Disease course.** Progressive neuromotor decline on static hypomyelination; lifelong ichthyosis. Complications include those of severe spasticity (contractures, mobility loss), swallowing/communication impairment, and skin-barrier-related issues (infection, dehydration).
- **Prognostic factors.** **Allele dose** is the dominant prognostic determinant — biallelic disease is more severe with 100%-penetrant developmental delay/spasticity and corpus callosum hypoplasia, versus milder monoallelic IKSHD. Plasma VLCFA depletion severity is a candidate biochemical prognostic marker (inferred).

12. Treatment

No disease-specific or ELOVL1-targeted therapy exists. Management is **symptomatic and multidisciplinary** ([PMID: 40590574](#); dermatologic evidence base [PMID: 23870202](#), [PMID: 36928932](#)).

Skin (congenital ichthyosis care).

Intervention	Class / mechanism	Evidence	NCIT suggestion
Topical emollients	Barrier repair/hydration	Therapeutic benefit, good safety in systematic review (PMID: 23870202)	Emollient
Keratolytics (urea, propylene glycol, lactic acid)	Desquamation	Standard of care (PMID: 36928932)	Keratolytic Agent
Calcipotriol ointment	Vitamin D analog	Benefit + good safety (PMID: 23870202)	Calcipotriene
Topical retinoids (tazarotene, isotretinoin/TMB-001)	Retinoid; normalizes keratinization	Liartzole benefit (PMID: 23870202); Phase IIb TMB-001 reduced scaling (PMID: 36928932)	Retinoid
Systemic retinoids (acitretin)	Oral retinoid	Effective but limited by teratogenicity/adverse effects (PMID: 36928932)	Acitretin

Neurological features (symptomatic). Antispasticity agents (e.g., baclofen, botulinum toxin — NCIT: Baclofen, OnabotulinumtoxinA), physiotherapy, occupational therapy, speech/language therapy, orthopedic management of contractures, and supportive/movement-disorder management. **No controlled trials exist specifically for IKSHD**; neurological care follows generic spasticity/leukodystrophy pathways.

Advanced/experimental therapeutics. No approved gene, cell, RNA-based, or targeted therapy. Conceptually, because IKSHD is a *biosynthetic deficiency*, **substrate/lipid-supplementation or elongase-augmentation** strategies are rational future directions (untested). Note that this is opposite to X-ALD, where *substrate-reduction* (e.g., targeting ELOVL1) is a therapeutic goal ([PMID: 42469918](#)) — highlighting ELOVL1's dual relevance across VLCFA disorders.

Pharmacogenomics. None specific to IKSHD.

13. Prevention

- **Primary prevention.** Not applicable to the genetic cause. **Genetic counseling** is central: recurrence risk is low for de novo dominant cases but ~25% for future pregnancies in recessive families. **Prenatal/preimplantation genetic testing** is feasible once the familial variant is known.
 - **Secondary prevention.** Early genetic diagnosis (WES/WGS) enables early institution of skin-barrier care, spasticity management, and developmental/rehabilitation support to limit complications.
 - **Tertiary prevention.** Prevent complications of spasticity (contracture prophylaxis via physiotherapy/orthotics), maintain skin-barrier integrity to reduce infection/dehydration, nutritional and communication support.
 - **Immunization / public-health / prophylaxis.** Standard pediatric care; no disease-specific vaccine or prophylaxis. Routine infection-prevention measures are prudent given barrier compromise.
 - **Counseling.** Genetic counseling for families is recommended, distinguishing de novo dominant (low sibling recurrence, but germline-mosaicism caveat) from recessive (25% recurrence) inheritance.
-

14. Other Species / Natural Disease

- **Taxonomy / orthologs.** *Elovl1* is conserved across mammals. Mouse *Elovl1* (NCBI Gene) is the principal experimental ortholog; the enzyme's elongation function is evolutionarily conserved.
- **Natural disease in other species.** No well-characterized naturally occurring *ELOVL1* disorder in companion animals or wildlife is documented (OMIA). The disease knowledge derives from human patients and engineered mouse models rather than spontaneous animal disease.
- **Comparative biology.** The mouse recapitulates key mechanisms: **Elovl1 knockout** causes lethal epidermal-barrier failure with reduced $\geq$ C26 ceramides (PMID: 23689133); **Elovl1-deficient mice** show reduced myelin-sphingolipid chain length and impaired motor coordination (PMID: 32123819) — demonstrating strong cross-species conservation of both the skin and CNS mechanisms.

- **Transmission.** Not applicable (non-infectious, non-zoonotic).

15. Model Organisms

- **Mouse (*Mus musculus*)** is the primary model.
 - **Elovl1 knockout:** neonatal-lethal from epidermal permeability-barrier defects; diminished stratum-corneum lipid lamellae and reduced $\geq$ C26 ceramides — establishing ELOVL1 as essential for barrier formation and ceramide chain length (PMID: 23689133).
 - **Elovl1-deficient (hypomorphic/partial) mice:** reduced myelin-sphingolipid chain length and poorer motor coordination — modeling the CNS hypomyelination/motor phenotype of human IKSHD (PMID: 32123819).
 - **In vitro models.** Patient **fibroblasts** and **transfected HEK293 cells** with VLCFA GC-MS profiling and stable-isotope substrate assays quantify ELOVL1 activity and demonstrate the elongation defect and compensatory ELOVL dysregulation (PMID: 30487246; PMID: 40590574).
 - **Genetic-model types available.** Knockout and hypomorphic mouse lines; humanized knock-in models carrying p.Ser165Phe would be a valuable (not-yet-reported) resource.
 - **Phenotype recapitulation.** Excellent for both compartments: skin-barrier/ceramide defect and myelin-sphingolipid/motor-coordination defect are both reproduced.
 - **Limitations.** Complete knockout is neonatally lethal (barrier failure), limiting study of the CNS phenotype in nulls; dysmorphic facial features and the human dominant-variant-specific effects are not well captured by simple knockouts.
-

Key Findings (with statistical evidence)

F001 — IKSHD is caused by a recurrent de novo heterozygous *ELOVL1* p.Ser165Phe. WES in two unrelated pediatric probands without family history identified a shared novel heterozygous *ELOVL1* variant; the same **c.494C>T (p.Ser165Phe)** de novo variant was independently confirmed in additional patients. *"To identify the cause of a similar phenotype of ichthyotic keratoderma, spasticity, mild hypomyelination (on MRI) and dysmorphic features (IKSHD)..."* (PMID: 29496980); *"Genetic analysis identified the de novo pathogenic variant p.Ser165Phe in ELOVL1."* (PMID: 42101073).

F002 — ELOVL1 elongates C22–C26 VLCFAs required for skin barrier and myelin. *"Elovl1 knockout mice died shortly after birth due to epidermal barrier defects" and "Elovl1 is a key determinant of epidermal Cer chain length and is essential for permeability barrier formation" (PMID: 23689133); "ELOVL fatty acid elongase 1 catalyses elongation of saturated and monounsaturated C22-C26-VLCFAs" (PMID: 30487246).*

F003 — Elovl1-deficient mice recapitulate CNS features. *"Reduced chain length in myelin sphingolipids and poorer motor coordination in mice deficient in the fatty acid elongase" — linking Elovl1 loss to the hypomyelination/motor phenotype (PMID: 32123819).*

F004 — Biallelic ELOVL1 variants cause a more severe hypomyelinating leukodystrophy. *In 7 patients: "Common clinical features included ichthyosis (5/7), developmental delay (7/7), progressive spasticity (7/7), nystagmus (5/6), and a complex movement disorder characterized by pronounced head tremor (7/7), myoclonus (6/7), and dysarthria (6/6)"; "Brain MRI revealed non-progressive hypomyelination (6/6) and hypoplasia of the corpus callosum (5/6)"; "Plasma VLCFA analysis in one patient showed reduced concentrations of C24:0 and C26:0" (PMID: 40590574).*

F005 — IKSHD is a VLCFA-biosynthesis defect, opposite to VLCFA-catabolism leukodystrophies. *"Variants in genes encoding enzymes responsible for catalyzing the first and rate limiting step in the production of VLCFAs, elongation of VLCFAs (ELOVLs), underlie a novel group of metabolic disorders" (PMID: 40590574); contrast with "X-linked adrenoleukodystrophy (ALD)... resulting in the systemic accumulation of very-long-chain fatty acids (VLCFAs)" (PMID: 42320861).*

F006 — ELOVL1 performs the rate-limiting condensation step of the ER four-enzyme elongation cycle. *"Elongation occurs in the endoplasmic reticulum (ER) through the actions of a complex of four ER-embedded enzymes, which includes the ELOVL proteins" and "which catalyze the first and rate-limiting step of the FA elongation cycle" (PMID: 39946831); "the fatty acid elongases can be divided into two major groups: (a) enzymes... involved in the elongation of saturated and monounsaturated VLCFA (ELOVL1, 3 and 6)" (PMID: 16564093).*

F007 — No disease-specific therapy; management is symptomatic. *"Topical treatments including emollients, calcipotriol ointment, and liarozole cream seem to have therapeutic benefit and a good safety profile" (PMID: 23870202); "Emollients and keratolytics are frequently used to manage symptoms of congenital ichthyosis... Systemic retinoid treatment is complicated by teratogenicity and dose-limiting adverse effects" (PMID: 36928932).*

F008 — Genotype–phenotype architecture is a graded severity spectrum. *"Monoallelic pathogenic variants in ELOVL1 have been described in association to a condition characterized by ichthyosis, spasticity, nystagmus and cerebral hypomyelination, although a similar but more*

severe presentation has been reported in patients with biallelic variants" (PMID: 42101073); "biallelic variants in ELOVL1 are associated with a unique and recognizable phenotype of hypomyelinating leukodystrophy, ichthyosis, and a complex movement disorder including progressive spastic..." (PMID: 40590574).

Mechanistic Model / Interpretation

IKSHD is best understood as a **single upstream biochemical lesion — deficient VLCFA elongation — producing two downstream tissue-specific lipid deficiencies**. Because ELOVL1 catalyzes the rate-limiting condensation step of ER fatty-acid elongation, its dysfunction throttles the supply of C22–C26+ VLCFAs. Those VLCFAs are the acyl backbones of two critical lipid classes: **epidermal ceramides** (whose ultra-long chains build the stratum-corneum lipid lamellae) and **myelin sphingolipids** (whose chain length underpins white-matter integrity). Depleting the first yields **ichthyotic keratoderma**; depleting the second yields **hypomyelination**, from which upper-motor-neuron dysfunction (spasticity) and a complex movement disorder emerge. The mouse genetics cleanly separate and confirm both arms: complete knockout kills neonates through the skin arm, while partial deficiency reveals the myelin/motor arm.

The **allele-dose gradient** — mild monoallelic IKSHD versus severe biallelic leukodystrophy — is the disorder's most instructive genetic feature and implies a **quantitative threshold** relationship between residual elongase output and phenotype severity. This threshold logic also frames the therapeutic hypothesis: because the defect is a *deficiency*, restoring VLCFA supply (substrate/lipid supplementation, or enhancing residual/paralogous elongase activity) is mechanistically rational — the inverse of X-ALD, where the therapeutic aim is to *reduce* VLCFAs (even using ELOVL1 as a drug target). Positioning IKSHD opposite X-ALD on a single VLCFA axis is both diagnostically useful (plasma VLCFAs low vs high) and conceptually unifying.

Evidence Base

PMID	Contribution	Type
29496980	Defines IKSHD; dominant <i>ELOVL1</i> variant in 2 unrelated probands	Human clinical
30487246	De novo <i>ELOVL1</i> mutation; enzyme substrate specificity (C22–C26 VLCFA)	Human clinical + in vitro
42101073	Confirms recurrent de novo p.Ser165Phe; states mono vs biallelic gradient	Human clinical
40590574	Biallelic <i>ELOVL1</i> → severe leukodystrophy; phenotype frequencies; reduced plasma C24:0/C26:0	Human clinical + in vitro
23689133	Elovl1-KO mouse: barrier failure, reduced ≥C26 ceramides	Model organism
32123819	Elovl1-deficient mouse: shortened myelin sphingolipids, motor deficits	Model organism
39946831	ELOVL biology; ER four-enzyme complex; rate-limiting step	Review
16564093	ELOVL1/3/6 elongate saturated/monounsaturated VLCFA	Review
42320861	X-ALD VLCFA accumulation (contrast)	Human clinical
42469918	ELOVL1 as substrate-reduction target in ALD (contrast/therapeutics)	Review
23870202	Systematic review of congenital-ichthyosis treatments	Systematic review
36928932	Topical isotretinoin (TMB-001) Phase IIb in congenital ichthyosis	Clinical trial
41165046	Commentary on the biallelic <i>ELOVL1</i> series	Correspondence

Note on citation: PMID: [32123819](#) was flagged as a snippet-abstract mismatch during investigation; the finding it supports (Elovl1 deficiency → shortened myelin sphingolipids and motor deficits) is corroborated by the broader mechanistic literature and the biallelic-patient VLCFA data, but the exact quote should be re-verified against the source.

Limitations and Knowledge Gaps

1. **Very small evidence base.** The disorder rests on a handful of case reports (dominant form) and a single 7-patient series (biallelic form); frequencies, penetrance, and natural history are imprecise.
 2. **Prognosis/survival data are absent.** No formal life-expectancy or survival statistics exist.
 3. **Molecular mechanism of the dominant p.Ser165Phe** (simple loss-of-function vs dominant-negative on the ER complex) is not definitively resolved.
 4. **Human brain lipidomics** directly demonstrating shortened myelin sphingolipids in patients are lacking; the CNS mechanism is largely inferred from mouse.
 5. **No epigenetic, transcriptomic, or proteomic disease signatures** have been reported.
 6. **No therapeutic trials** target the underlying VLCFA deficiency; all skin-treatment evidence is extrapolated from congenital ichthyosis generally.
 7. **One key mouse citation had a snippet mismatch** and warrants source re-verification.
 8. **Ontology mapping** (MONDO, ICD-11, Orphanet ORPHAcode) is incompletely standardized for this newly delineated entity.
-

Proposed Follow-up Experiments / Actions

1. **Establish a patient registry** capturing dominant vs biallelic genotypes with longitudinal MRI, plasma VLCFA, motor, and dermatologic outcomes to define natural history and prognosis.
2. **Generate a humanized p.Ser165Phe knock-in mouse** to test whether the dominant variant acts by loss-of-function or dominant-negative mechanism and to model the CNS phenotype without neonatal lethality.
3. **Perform patient plasma/CSF and, where available, brain lipidomics** to directly confirm VLCFA-ceramide and myelin-sphingolipid depletion and to identify candidate biomarkers of severity.
4. **Test VLCFA/lipid-supplementation and elongase-augmentation strategies** in patient fibroblasts and mouse models as a mechanism-based therapeutic proof-of-concept (the inverse of X-ALD substrate reduction).

5. **Screen paralogous elongases (ELOVL3/6/7) and ceramide-synthesis genes** as modifiers to explain the mono-to-biallelic severity gradient.
 6. **Standardize diagnostics:** promote plasma VLCFA profiling (deficiency pattern) plus *ELOVL1* sequencing as a combined diagnostic panel, and formalize MONDO/ICD-11/Orphanet coding.
 7. **Re-verify the mouse myelin-sphingolipid citation** ([PMID: 32123819](#)) against the primary source.
-

Report compiled from a 5-iteration autonomous investigation: 8 confirmed findings, 24 papers reviewed. Evidence types span human clinical case reports/series, model-organism (mouse), and in vitro/biochemical studies.

Generated by OpenScientist — Scientific Hypothesis Agent for Novel Discovery