Trichorhinophalangeal Syndrome Type I (TRPS I): A Comprehensive Disease-Characterization Report

Disease: Trichorhinophalangeal Syndrome Type I **MONDO ID:** MONDO:0008596 · **OMIM:** 190350 · **Orphanet:** ORPHA:77258 · **Category:** Mendelian (autosomal dominant) **Causal gene:** *TRPS1* (8q23.3) · **HGNC:** 12340 · **OMIM:** 604386

Summary

Trichorhinophalangeal syndrome type I (TRPS I) is a rare (~1 in 100,000; minimum point prevalence ~0.5/100,000) autosomal-dominant craniofacial–skeletal dysplasia caused by **heterozygous loss-of-function/haploinsufficient variants in *TRPS1***, a gene on chromosome 8q23.3 encoding a GATA-type zinc-finger transcriptional repressor. The disorder is defined by a clinical triad reflected in its name: **tricho** (fine, sparse, slow-growing scalp hair and thin lateral eyebrows), **rhino** (a bulbous, pear-shaped nose with underdeveloped/hypoplastic alae nasi), and **phalangeal** (brachydactyly with pathognomonic cone-shaped epiphyses). Short stature, hip dysplasia, and progressive premature osteoarthrosis round out the phenotype. Intelligence and life expectancy are normal.

Mechanistically, TRPS1 is a master transcriptional regulator of **endochondral ossification, hair-follicle morphogenesis, and odontogenesis**. Reduced TRPS1 activity **de-represses RUNX2, Indian hedgehog (Ihh), SOX9, STAT3, and PTHrP in growth-plate cartilage**, uncoupling chondrocyte differentiation from perichondrial mineralization; it **disrupts a Wnt-inhibitor/transcription-factor network required for hair-follicle development**; and it **dysregulates Dspp-dependent dentin mineralization**. Most pathogenic variants are **truncating (nonsense/frameshift) LOF alleles causing haploinsufficiency**, whereas **missense variants in the exon-6 GATA DNA-binding domain act in a dominant-negative fashion** and produce the more severe, brachydactyly-predominant TRPS III end of the allelic spectrum. TRPS II (Langer–Giedion syndrome) is a contiguous-gene deletion removing *TRPS1* plus *EXT1* (± *RAD21*), adding multiple exostoses and sometimes intellectual disability.

There is no disease-modifying or curative therapy. Management is **symptomatic and multidisciplinary**: recombinant human growth hormone (rhGH) in the subset with documented GH deficiency, bisphosphonates for severe osteoporosis, orthopedic surgery for hip/joint disease, dental care for supernumerary/impacted teeth, and genetic counseling for the 50% transmission risk. TRPS1 biology extends well beyond the clinical triad—to kidney development/fibrosis (BMP7/TGF-β/Smad) and to oncology, where TRPS1 is now a highly sensitive, largely breast-specific immunohistochemistry marker.

Section 1 — Disease Information

Overview. TRPS I is a hereditary skeletal dysplasia with a characteristic clinical presentation, classified into types I, II, and III based on phenotype and genotype (PMID: 21844942). TRPS I—the most common subtype—is caused by heterozygous mutations in the single gene *TRPS1*; it exhibits distinctive craniofacial, ectodermal, and skeletal abnormalities with **extremely variable expressivity**, such that many mild cases remain undiagnosed (PMID: 30458885).

Key identifiers.

Resource	Identifier
MONDO	MONDO:0008596
OMIM	190350 (TRPS I)
Orphanet	ORPHA:77258
MeSH	Langer-Giedion Syndrome / Tricho-rhino-phalangeal syndrome
ICD-10	Q87.1 (congenital malformation syndromes predominantly associated with short stature)
Gene (HGNC)	*TRPS1*, HGNC:12340, OMIM 604386, locus 8q23.3 (older literature: 8q24.1/8q24.12)

Synonyms / alternative names. Tricho-rhino-phalangeal syndrome type I; TRPS1; TRP syndrome type I; trichorhinophalangeal dysplasia syndrome (PMID: 3958163); "Giedion syndrome" is used for the broader TRPS spectrum (PMID: 37147064).

Information source. The knowledge base here is derived primarily from **aggregated disease-level resources** (OMIM, Orphanet) supplemented by **individual-patient clinical reports and cohort studies** (103-patient international cohort, PMID: 25792522; 20-child Korean cohort, PMID: 42361898; 15-patient single-center series, PMID: 38574886) and from **mouse/in-vitro mechanistic studies**. No local datasets or EHR data were used; this is a literature synthesis.

Section 2 — Etiology

Primary cause (genetic). TRPS I is **monogenic and autosomal dominant**, caused by heterozygous pathogenic variants in *TRPS1*. In an international cohort of 103 molecularly/cytogenetically confirmed patients, TRPS I was present in 85 individuals (22 missense, 62 "other" mutations) versus 14 with TRPS II (PMID: 25792522). The review by Saeki et al. states the disorder "occurs in 1 in 100,000 people globally and is caused by several types of mutations of the TRPS1 gene" (PMID: 40995872).

Genetic risk factors. The causal variant in *TRPS1* is the sole established risk factor. **No susceptibility loci or modifier genes outside *TRPS1* have been established in humans**, though *RUNX2* is a genetic modifier of the growth-plate phenotype in mice (double-heterozygote rescue; PMID: 18424451).

Environmental risk factors. None known. TRPS I is a Mendelian disorder with no demonstrated environmental or infectious contribution. A substantial fraction of cases are de novo ("fresh") mutations (PMID: 10405449).

Protective factors. No genetic or environmental protective factors have been described. *Not applicable* for this Mendelian disorder.

Gene–environment interactions. None documented. *Not applicable.*

Section 3 — Phenotypes

The phenotype is variable and presents a wide clinical spectrum (PMID: 31343132). The defining features, per the 103-patient cohort (PMID: 25792522), are **fine/sparse hair, thick medial eyebrows, broad/bulbous nasal ridge and tip, underdeveloped nasal alae, short hands/feet, cone-shaped epiphyses, and reduced linear growth**.

Phenotype	Type	HPO term (suggested)	Onset	Frequency / severity
Sparse, fine, slow-growing scalp hair	Ectodermal/ physical	HP:0002213 (sparse hair)	Congenital/ childhood	Near-universal; variable; non-terminal hairs, low density (PMID: 40790426)
Thin lateral eyebrows / thick medial eyebrows	Physical	HP:0045025 (lateral eyebrow thinning)	Childhood	Common; diagnostic clue
Bulbous/pear-shaped nose, hypoplastic alae nasi	Physical	HP:0000414 (bulbous nose), HP:0000430 (underdeveloped nasal alae)	Congenital	Near-universal; characteristic
Brachydactyly, short metacarpals/ phalanges	Skeletal	HP:0001156 (brachydactyly)	Childhood (radiographic)	Near-universal; **severe with exon-6 missense** (PMID: 42361898)
Cone-shaped epiphyses	Skeletal/ radiographic	HP:0010579 (cone-shaped epiphysis)	Childhood	Pathognomonic radiographic sign
Short stature	Growth	HP:0004322 (short stature)	Prenatal + postnatal	Common; variable; GH deficiency in a subset
Hip dysplasia / Perthes-like changes	Skeletal	HP:0001385 (hip dysplasia)	Childhood–adult	Frequent; dominant morbidity (PMID: 3958163)
Progressive premature osteoarthrosis	Skeletal	HP:0002758 (osteoarthritis)	Childhood → adult	**100% in one cohort** (PMID: 38574886); progressive
	Skeletal/lab			

Phenotype	Type	HPO term (suggested)	Onset	Frequency / severity
Osteopenia / low bone mass		HP:0000938 (osteopenia)	Childhood–adult	Common (PMID: 25792522)
Dental anomalies (supernumerary/ impacted teeth, microdontia)	Physical	HP:0011069 (supernumerary tooth), HP:0000699 (microdontia)	Childhood	Variable; up to 45 teeth reported (PMID: 37147064)
Protruding ears	Physical	HP:0000411 (protruding ear)	Congenital	Common

Age of onset. Most features are **congenital or manifest in childhood**; radiographic cone-shaped epiphyses appear in childhood; joint disease often precedes diagnosis and can present as late as adulthood (>40 years; PMID: 3958163).

Progression. Craniofacial and digital features are **stable/structural**; joint disease and osteoarthrosis are **progressive**.

Quality-of-life impact. The dominant QoL burden is **lifelong joint disease**—reported by all patients in one cohort, often before diagnosis (PMID: 38574886)—plus cosmetic concerns (hair, nose, stature) and dental complications. Formal EQ-5D/SF-36 data are not available for TRPS I (*knowledge gap*).

Section 4 — Genetic / Molecular Information

Causal gene. *TRPS1* (8q23.3; OMIM 604386), encoding a **GATA-type transcription factor with nine zinc-finger motifs** that functions primarily as a **transcriptional repressor** (PMID: 21630221).

Pathogenic variants — spectrum and classification. Most variants are **truncating loss-of-function**. In the landmark mutation survey, "seven of the nine mutations result in a premature stop codon, leading to a truncated, nonfunctional TRPS1 protein. The final two mutations are missense mutations in the GATA DNA binding zinc finger" (PMID: 11950061). The Korean cohort reported **truncating 65%, exon-6 missense 15%, exonic deletions 20%** (PMID: 42361898).

Representative variants (all ACMG pathogenic/likely-pathogenic, germline, heterozygous):

Variant (cDNA / protein)	Type	Phenotype note	PMID
c.2086C>T (p.Arg696Ter)	Nonsense	Classic TRPS I, familial	30458885
c.2854_2858del (p.Asn952ArgfsTer2)	Frameshift	TRPS I, 4-generation family	36093893
c.880_882delAAG	In-frame del	Disturbed RUNX2/STAT3 repression	36990068
c.2783A>G (p.Tyr928Cys)	Missense (GATA domain)	**Severe** brachydactyly + short stature	30914275
c.2762G>A (p.Arg921Gln)	Missense	Dental anomalies, impaired dentin mineralization	32442662
p.Ala932Ser (exon 6, GATA)	Missense	TRPS III	28468609
c.3809_3811delACTinsCATGTTGTG	Indel (Ikaros-like ZnF)	TRPS I + supernumerary teeth	27706911

Functional consequences. Truncating variants → **haploinsufficiency (LOF)**. **GATA-domain missense variants act as dominant-negatives**, disturbing the protein's transcriptional repression of targets such as RUNX2 and STAT3 despite normal cellular localization and expression (PMID: 36990068).

Allele frequency. Pathogenic *TRPS1* variants are private/rare and essentially absent from population databases (gnomAD); the disorder is fully penetrant and dominant.

Modifier genes. *RUNX2* dosage modifies the skeletal phenotype in mouse models (double-heterozygote rescue; PMID: 18424451). No validated human modifiers.

Epigenetic information. No disease-specific DNA-methylation/histone signature has been defined for TRPS I (*knowledge gap*). Note that TRPS1 itself acts on chromatin as a transcriptional regulator.

Chromosomal abnormalities. Large deletions at **8q23–q24** cause the contiguous-gene TRPS II (Langer–Giedion), removing *TRPS1*, *RAD21*, and *EXT1*; a ~3.2-Mb critical region defines typical LGS (PMID: 35290978; PMID: 41683676). Intragenic deletions/microdeletions of *TRPS1* alone cause TRPS I.

Section 5 — Environmental Information

Environmental factors. None established. TRPS I is purely genetic. **Lifestyle factors.** None. *Not applicable.* **Infectious agents.** None. *Not applicable.*

(Any environmental contribution would be limited to general modifiers of growth/bone health; no TRPS-specific data exist.)

Section 6 — Mechanism / Pathophysiology

Ordered causal chain (initiating lesion → clinical manifestation)


Skeletal/chondrogenic branch: 1. A heterozygous *TRPS1* variant (truncating LOF → haploinsufficiency, **or** exon-6 GATA-domain missense → dominant-negative) **results in** reduced functional TRPS1 transcriptional-repressor activity in cartilage. 2. Reduced TRPS1 **leads to** de-repression of its direct targets **RUNX2, SOX9, STAT3, PTHrP** and increased **Indian hedgehog (Ihh)** expression/signaling in the growth plate (PMID: 18424451; PMID: 24709795). 3. Loss of TRPS1–RUNX2 repression **results in** delayed chondrocyte differentiation but accelerated perichondrial mineralization—i.e., **uncoupling of chondrocyte differentiation from perichondrial mineralization** (PMID: 18424451). 4. Dysregulated chondrocyte condensation, proliferation, maturation, apoptosis, and phalangeal-joint formation (downstream of Gdf5 signaling; TRPS1 also directly interacts with activated Gli3) **leads to** abnormal endochondral ossification (PMID: 24709795). 5. Abnormal endochondral ossification **results in** the **cone-shaped epiphyses, brachydactyly, short stature, hip dysplasia, and premature osteoarthrosis** observed clinically.

Hair-follicle branch: 1. Reduced TRPS1 in dermal papilla / peri-hair-peg mesenchyme **leads to** downregulation of a network of **secreted Wnt inhibitors and transcription factors** and upregulation of extracellular-matrix proteins (PMID: 22115758). 2. (TRPS1 here acts as both repressor *and* **direct transcriptional activator**, binding target-gene promoters to activate them — PMID: 22115758.) 3. Disrupted Wnt-dependent follicle morphogenesis **results in** reduced/absent follicles → **sparse, fine, slow-growing, non-terminal hair** (PMID: 40790426).

Dental branch: 1. Dysregulated TRPS1 in odontoblasts **alters** direct repression of the **Dspp** promoter and expression of mineralization genes (Alpl, Phospho1) (PMID: 22508542; PMID: 25128529). 2. This **leads to** impaired dentin mineralization and, via APC/RUNX2/Wnt dysregulation, **supernumerary/impacted teeth and dental anomalies** (PMID: 32442662; PMID: 27706911).

(Steps 2–4 of the skeletal branch are directly demonstrated in mouse models and in-vitro assays; their quantitative applicability to human tissue is inferred.)

ASCII schematic



Molecular pathways and processes

- **Pathways:** Indian hedgehog (Ihh) signaling; Wnt signaling (via secreted Wnt inhibitors); Gdf5/BMP signaling; TGF-β/Smad (chiefly in kidney). KEGG/Reactome: Hedgehog signaling, Wnt signaling pathway.
- **Cellular processes (GO):** chondrocyte differentiation (GO:0002062), endochondral ossification (GO:0001958), regulation of transcription (GO:0006357), hair follicle morphogenesis (GO:0031069), apoptotic process (GO:0006915), biomineral tissue development (GO:0031214).
- **Protein dysfunction:** truncation → nonfunctional protein (LOF); GATA-domain missense → impaired DNA binding and loss of repression (dominant-negative) (PMID: 11950061; PMID: 36990068).

- **Cell types (CL):** chondrocyte (CL:0000138), odontoblast (CL:0000060), dermal papilla / perichondrial mesenchymal cells, osteoblast (CL:0000062).
- **Pleiotropy beyond the triad:** In kidney, Trps1 acts downstream of Bmp7 in mesenchymal–epithelial transition; its loss activates **TGF-β/Smad3**, suppressing ureteric-bud branching (PMID: 23537899), and haploinsufficiency promotes renal fibrosis via increased Arkadia and reduced Smad7 (PMID: 20507941). Trps1 also promotes tubular repair after ischemia-reperfusion via Pde4d/AKT (PMID: 27466160).
- **Molecular profiling:** Microarray of mutant mouse whisker pads identified downregulated Wnt inhibitors/transcription factors and upregulated ECM proteins (PMID: 22115758); ChIP-seq in kidney/cancer contexts confirms direct promoter binding.

Section 7 — Anatomical Structures Affected

Organ/system level. - **Primary:** skeleton (hands, feet, phalanges, hips, long bones), skin appendages (hair follicles), nose/craniofacial structures, teeth. - **Body systems:** musculoskeletal (dominant), integumentary (ectodermal), craniofacial. - **Secondary/pleiotropic (chiefly from model organisms):** kidney (development, fibrosis); oncologic relevance (breast and other GATA-factor-expressing tumors).

UBERON suggestions: phalanx (UBERON:0003221), epiphysis (UBERON:0006716), hip joint (UBERON:0001464), growth plate (UBERON:0006233), hair follicle (UBERON:0002073), nose (UBERON:0000004), tooth (UBERON:0001091), dentine (UBERON:0001751), kidney (UBERON:0002113).

Tissue/cell level. Cartilage (growth-plate chondrocytes, CL:0000138), perichondrium/mesenchyme, dermal papilla and peri-hair-peg mesenchymal cells, odontoblasts (CL:0000060), osteoblasts (CL:0000062).

Subcellular level (GO Cellular Component). Nucleus (GO:0005634) — TRPS1 is a nuclear transcription factor with sequence-specific DNA binding at target-gene promoters.

Localization / lateralization. Skeletal involvement is **bilateral and generally symmetric** (hands, feet, hips); hair and facial features are midline/bilateral.

Section 8 — Temporal Development

Onset. Largely **congenital** facial/ectodermal features; **childhood** radiographic skeletal signs; onset pattern is **chronic/insidious**. Diagnosis is often delayed—median age at diagnosis 15 years in one series (PMID: 38574886), with some patients first diagnosed after age 40 (PMID: 3958163).

Progression. Craniofacial/digital morphology is structurally **stable**. **Joint disease is progressive** — Perthes-like hip dysplasia and "severe and progressive degenerative arthritis" (PMID: 3958163); premature osteoarthrosis (PMID: 3621840). Growth retardation is prenatal and postnatal.

Disease course/duration. Chronic, lifelong; non-remitting. No spontaneous remission; interventions (rhGH, bisphosphonates, orthopedic surgery) modify specific complications.

Critical periods. The childhood/prepubertal window is key for growth intervention—GH therapy is most effective when initiated earlier and continued longer in prepubertal/early-pubertal stages (PMID: 36990068).

Section 9 — Inheritance and Population

Epidemiology. Prevalence **~1 in 100,000** globally (PMID: 40995872); **minimum point prevalence 0.5/100,000 (95% CI 0.3–0.8)** in a single-center series (PMID: 38574886). Incidence not formally established; many mild cases are undiagnosed.

Inheritance. Autosomal dominant. Both inherited and de novo ("fresh") mutations occur (PMID: 10405449). Transmission risk to offspring is **50%**.

Penetrance / expressivity. High penetrance but **markedly variable expressivity**, even within families (PMID: 25792522; PMID: 30458885).

Genotype–phenotype correlation. Exon-6 GATA-domain missense variants → severe brachydactyly (consistent across cohorts; PMID: 42361898; PMID: 30914275), and define TRPS III. Stature shows no clear genotype correlation (PMID: 42361898). Facial features do not differ between TRPS I and II (PMID: 25792522).

Anticipation / mosaicism / founder effects / consanguinity / carrier frequency. No genetic anticipation (not a repeat-expansion disorder). Germline mosaicism not specifically characterized. No founder effects or consanguinity role (dominant, often de novo). Carrier frequency *not applicable* (dominant).

Demographics. Reported worldwide (European, Turkish, Korean, Chinese, Thai, Moroccan, Cypriot cohorts). Female predominance noted in some series (e.g., 73% female; PMID: 38574886), likely ascertainment-related rather than a true sex effect. No established sex-ratio bias expected in a dominant disorder.

Section 10 — Diagnostics

Clinical/radiographic. Diagnosis is based on **characteristic facial features** + **sparse hair** + **brachydactyly with cone-shaped epiphyses on hand/foot X-rays** (PMID: 21844942). Cone-shaped epiphyses are the hallmark radiographic sign (PMID: 42705688). **Trichoscopy** reveals non-terminal, thin, sparse hairs with low density and increased single-hair follicular units; a **high occipital hairline** is proposed as a diagnostic pearl (PMID: 40790426). DXA documents low bone mineral density/osteopenia. Endocrine work-up (GH stimulation tests) identifies the GH-deficient subset (PMID: 22964620). Full dental/radiographic examination is recommended to detect supernumerary/impacted teeth (PMID: 37147064).

Genetic testing. Molecular confirmation by **TRPS1 sequencing** (Sanger / next-generation / whole-exome) is the gold standard (PMID: 36990068; PMID: 42705688). **Chromosomal microarray / molecular karyotyping** detects 8q23–q24 contiguous-gene deletions distinguishing TRPS II (PMID: 36866988; PMID: 35290978). Skeletal-dysplasia NGS panels, CMA, and MLPA are all employed (PMID: 40626694). WES is valuable when the clinical picture is atypical or overlaps mimics.

Clinical criteria / differential diagnosis. Differentials include **cleidocranial dysplasia** (RUNX2; supernumerary teeth; PMID: 30095610), **FBXO11-related IDDFBA syndrome** (TRPS-like facies but with developmental delay; PMID: 38679370), juvenile idiopathic arthritis and other skeletal dysplasias presenting with non-inflammatory musculoskeletal complaints (PMID: 40626694), and TRPS II/III within the spectrum.

Screening. No population newborn screening (rare dominant disorder). **Cascade testing** of at-risk relatives and prenatal/preimplantation testing are feasible once the familial variant is known.

Section 11 — Outcome / Prognosis

Survival/mortality. Life expectancy is normal. TRPS I is not life-limiting; intelligence is normal in TRPS I (PMID: 10405449).

Morbidity. The **dominant morbidity is joint disease**—reported by **all patients** in one cohort, often before diagnosis (PMID: 38574886); "morbidity mostly concerns joint problems" in the 103-patient cohort (PMID: 25792522). Progressive premature osteoarthrosis and Perthes-like hip dysplasia may require hip replacement (PMID: 3958163; PMID: 3621840). Osteopenia/osteoporosis can cause fractures (PMID: 24357341).

Complications. Degenerative arthritis, hip deformity, fractures from low BMD, dental impaction/malocclusion, short stature. Occasional rare findings: renal cyst/stone (PMID: 3621840), cerebral hamartoma/long-bone cysts (rare; PMID: 36866988).

Prognostic factors. Exon-6 GATA-domain missense predicts more severe brachydactyly (PMID: 42361898; PMID: 30914275). GH-deficiency status predicts GH-therapy response (PMID: 22964620).

Section 12 — Treatment

There is no causal/curative treatment; management is symptomatic and supportive, requiring coordination among pediatrics, endocrinology, orthopedics, dermatology, rehabilitation, and dentistry (PMID: 21844942; PMID: 31343132).

Intervention	Indication	Evidence	NCIT (suggested)
Recombinant human GH (rhGH)	Short stature with documented GH deficiency	Improved growth velocity & radiographic femoral epiphyses (PMID: 31884116); 5.7±0.9 cm/yr over 10 yr in GH-deficient patient, poor response if non-deficient (PMID: 22964620); improved linear growth over 2 yr in two patients (PMID: 36990068)	Recombinant Human Growth Hormone (NCIT:C1898)
Bisphosphonates (IV neridronate)	Severe osteoporosis / recurrent fractures	Off-label 2 mg/kg q3mo stopped new fractures and raised lumbar BMD by **+45%** (PMID: 24357341)	Bisphosphonate (NCIT:C1476)
Minoxidil (topical)	Hair thinning	Variable response (PMID: 42361898)	Minoxidil (NCIT:C47641)
Orthopedic surgery	Hip dysplasia, degenerative arthritis	Valuable for many (PMID: 21844942)	Orthopedic Surgery (NCIT:C15300)
Dental surgery / extraction	Impacted & supernumerary teeth	4 third molars + 11 supernumerary teeth removed under GA (PMID: 37147064)	Tooth Extraction (NCIT:C51818)
Cosmetic aids, rehabilitation (PT/OT)	Hair, stature, joint function	Supportive (PMID: 21844942)	—
Genetic counseling	50% transmission risk	Recommended for patient and family (PMID: 21844942)	Genetic Counseling (NCIT:C15648)

Pharmacogenomics / advanced therapeutics. No gene therapy, cell therapy, RNA-based, targeted, or immunotherapy exists for TRPS I. Animal-model work proposes **potential therapeutic targets** in the dysregulated pathways (RUNX2/Ihh, Wnt, TGF-β/Smad) but none are clinically translated (PMID: 40995872). No TRPS-specific clinical trials (NCT) are established.

Treatment outcomes/strategy. rhGH benefit is **conditional on GH deficiency** (PMID: 22964620)—a genotype/phenotype-guided (personalized) approach. Earlier, longer GH therapy in prepubertal/early-pubertal stages associates with better height outcomes (PMID: 36990068).

Section 13 — Prevention

Primary prevention. None possible for a dominant genetic disorder other than **reproductive options**: genetic counseling, prenatal diagnosis, and **preimplantation genetic testing** once the familial *TRPS1* variant is known.

Secondary prevention. Cascade genetic testing of at-risk relatives enables early diagnosis; early orthopedic, endocrine, and dental surveillance allows timely intervention (PMID: 30458885; PMID: 37147064).

Tertiary prevention. Prevent complications: DXA monitoring and bisphosphonates to prevent fractures (PMID: 24357341); orthopedic management to delay/manage degenerative arthritis; dental monitoring to prevent impaction complications.

Immunization / public-health / environmental interventions. *Not applicable* (non-infectious, non-environmental).

Counseling. Genetic counseling is central, emphasizing the 50% transmission risk and variable expressivity (PMID: 21844942).

Section 14 — Other Species / Natural Disease

Taxonomy / orthologs. *TRPS1* is conserved; the principal experimental ortholog is mouse **Trps1** (*Mus musculus*, NCBI Taxon:10090), with rat (*Rattus norvegicus*, Taxon:10116) used for kidney studies (PMID: 27466160).

Natural disease in other species / breeds. No naturally occurring TRPS-equivalent disease in companion animals or wildlife is documented in the reviewed literature (OMIA entry *not identified*; *knowledge gap*).

Comparative biology / evolutionary conservation. The TRPS1 regulatory functions in cartilage, hair follicle, kidney, and tooth are **evolutionarily conserved**, as mouse models recapitulate human skeletal, hair, and dental phenotypes (PMID: 40995872; PMID: 21630221).

Transmission / zoonosis. *Not applicable* (non-communicable genetic disorder).

Section 15 — Model Organisms

Rodent models are central to TRPS research—"several animal models have been established and studied intensively to address this rare disease" (PMID: 40995872).

Model	Type	Phenotype recapitulated	PMID
Trps1^Δgt/Δgt (GATA-domain deletion)	Knock-in/ deletion	Delayed chondrocyte differentiation, accelerated perichondrial mineralization, ↑Runx2/Ihh; reduced pelage follicles, absent vibrissae, ↓Wnt inhibitors	18424451; 22115758
Trps1-null	Knockout	Early postnatal lethality; expanded renal interstitium, impaired ureteric-bud branching (TGF-β/ Smad3)	23537899
Trps1^+/–	Haploinsufficient	Renal tubulointerstitial fibrosis via ↑Arkadia / ↓Smad7	20507941
Trps1^Δint/Δint (regulatory-element deletion, hypomorph)	Novel knock-in	Postnatal growth retardation, delayed secondary ossification centers, acetabular/hip dysplasia, patellar dislocation	38899779
Col1a1-Trps1 transgenic (overexpression)	Transgenic	Dentinogenesis-imperfecta-like dentin via Dspp-promoter repression	22508542
Preodontoblast cell lines (Trps1-deficient / -overexpressing)	In vitro	Dual role in dentin mineralization (Alpl/Phospho1)	25128529; 30691926

Model value and limitations. The **Trps1^Δint hypomorph** is notable for recapitulating *postnatal* hip-joint pathology relevant to the dominant human morbidity (PMID: 38899779). A key limitation is that **homozygous null mice are postnatally lethal**, whereas human TRPS I is

heterozygous and viable—so dosage-sensitive models (Δgt heterozygotes, hypomorphs) better mirror the human condition. Models have been used to dissect RUNX2/Ihh, Wnt, TGF-β/Smad, and Dspp mechanisms and to nominate therapeutic targets (PMID: 40995872). **Resource:** Mouse Genome Informatics (MGI) for *Trps1* alleles.

Mechanistic Model / Interpretation

TRPS1 is best understood as a **dosage-sensitive master transcriptional regulator** of mesenchymal differentiation. The unifying theme across all three cardinal tissues (cartilage, hair, tooth)—and the pleiotropic kidney phenotype—is that **TRPS1 sets the correct timing and level of differentiation programs**, principally by *repressing* key drivers (RUNX2, SOX9, STAT3, PTHrP, Dspp) but also, context-dependently, *activating* others (Wnt inhibitors, FOXA1). When TRPS1 activity falls below a threshold—whether by haploinsufficiency (truncating variants) or dominant-negative poisoning of the GATA domain (exon-6 missense)—these programs become **temporally and quantitatively uncoupled**, producing structural malformation rather than cell death.

The genotype–phenotype gradient is mechanistically coherent: truncating LOF alleles remove one functional copy (milder, classic TRPS I), whereas GATA-domain missense alleles produce a mutant protein that **actively interferes** with DNA binding, yielding more severe brachydactyly and the TRPS III phenotype (PMID: 11950061; PMID: 28468609). TRPS II layers an *EXT1*-deletion exostoses phenotype onto the *TRPS1* deficiency. This single axis—**TRPS1 dosage/function**—thus explains the entire TRPS I/III/II allelic and contiguous-gene spectrum.

Evidence Base

PMID	Role	Key contribution
25792522	Supports Disease Info, Genotype, Prognosis	103-patient cohort; mutation spectrum; morbidity = joints
40995872	Supports Epidemiology, Models	Prevalence ~1/100,000; animal-model synthesis; therapeutic targets
18424451	Supports Mechanism	Core skeletal mechanism: TRPS1–RUNX2 repression, ↑Ihh, uncoupling
42361898	Supports Genotype-Phenotype	Exon-6 missense → severe brachydactyly; genotype classes
30914275	Supports Genotype-Phenotype	GATA-domain missense → severe phenotype
31884116, 22964620	Support Treatment	rhGH efficacy conditional on GH deficiency
22115758	Supports Mechanism (hair)	TRPS1 activates Wnt-inhibitor network; dual repressor/activator
32442662, 22508542	Support Mechanism (dental)	Dental phenotype; Dspp-promoter repression mechanism
38574886	Supports Epidemiology, Prognosis	Prevalence 0.5/100,000; joint disease in 100%
20507941, 38647255	Support Pleiotropy	Renal fibrosis (TGF-β/Smad); TRPS1 as breast-cancer IHC marker
24357341	Supports Treatment	Bisphosphonate +45% BMD, no new fractures
11950061, 28468609	Support Variant Spectrum	Truncating LOF predominance; TRPS I/II/III distinctions
24709795	Supports Mechanism	Multi-target chondrogenic network (Sox9, Runx2, Stat3, Pthrp, Gli3)
38899779	Supports Models	Hypomorphic mouse recapitulates postnatal hip pathology

Challenging/nuancing evidence: The cancer literature (PMID: 26969828, 40197498) shows TRPS1 can be **context-dependently tumor-suppressive or oncogenic**, underscoring that "loss of function" is tissue-specific and that TRPS1 acts as both repressor and activator—caution against over-simplified LOF narratives.

Limitations and Knowledge Gaps

1. **No systematic natural-history/QoL data** — formal EQ-5D/SF-36/PROMIS outcomes for TRPS I are unavailable; cohorts are small and retrospective.
2. **Epidemiology uncertainty** — prevalence estimates (~1/100,000; minimum 0.5/100,000) derive from small/single-center series; incidence unknown; underdiagnosis of mild cases is acknowledged.
3. **Mechanism mostly from mouse/in-vitro** — the RUNX2/Ihh/Wnt/Dspp causal chain is well supported in models but less directly validated in human tissue; several steps are **inferred**.
4. **No disease-specific epigenetic signature** has been defined.
5. **No OMIA/veterinary natural-disease counterpart** identified; comparative biology rests on engineered rodent models.
6. **Human modifier genetics** largely unexplored; *RUNX2* is a mouse modifier only.
7. **Therapy evidence is anecdotal** — rhGH and bisphosphonate data come from case reports/small series, not controlled trials; no targeted/molecular therapy exists.

Proposed Follow-up Experiments / Actions

1. **Prospective multicenter TRPS registry** with standardized QoL instruments and longitudinal joint/skeletal outcomes to quantify natural history and GH/bisphosphonate efficacy.
2. **Patient-derived iPSC chondrocyte/organoid models** carrying truncating vs. GATA-domain-missense variants to directly test the haploinsufficiency vs. dominant-negative dichotomy on RUNX2/Ihh/SOX9 targets in human cells.
3. **Genome-wide TRPS1 ChIP-seq/CUT&RUN + RNA-seq** in human growth-plate chondrocytes and hair-follicle dermal papilla to map the direct target network and validate the dual repressor/activator model.

4. **Controlled or registry-based evaluation of bisphosphonates** for TRPS-associated osteoporosis, given the striking single-case +45% BMD response.
5. **Screen the RUNX2/Ihh and Wnt-inhibitor axes for druggable nodes** (e.g., Ihh-pathway modulation) using the Trps1^Δint hypomorphic mouse as a preclinical platform for hip-dysplasia/osteoarthrosis prevention.
6. **Define exon-6 missense dominant-negative mechanism** biochemically (DNA-binding, co-repressor recruitment) to support variant classification and prognostic counseling.

Report compiled from 12 confirmed findings and 53 reviewed papers across 5 investigation iterations. Evidence types: human clinical cohorts/case reports, mouse/rat model-organism studies, and in-vitro functional assays, as annotated throughout.
