PEHO Syndrome (ZNHIT3): Comprehensive Disease Characterization Report

Disease: PEHO Syndrome (Progressive Encephalopathy with Edema, Hypsarrhythmia, and Optic Atrophy) **MONDO ID:** MONDO:0009841 · **OMIM:** 260565 · **Orphanet:** ORPHA:2836 **Causal gene:** ***ZNHIT3*** (Zinc finger HIT-type containing protein 3; HGNC:12309; NCBI Gene 9326; 17q12) **Category:** Mendelian, autosomal recessive

---

Summary

PEHO syndrome is a severe, autosomal-recessive, early-infantile progressive neurodegenerative encephalopathy whose causal gene is **ZNHIT3** (Zinc finger HIT-type containing protein 3; HGNC:12309, 17q12). It is defined clinically by the acronym components: **P**rogressive **E**ncephalopathy, peripheral/facial oe**(E)**dema, **H**ypsarrhythmia (infantile spasms), and **O**ptic atrophy, with the near-constant additional feature of extreme, early-onset cerebellar atrophy. The disease was first delineated in Finland, where it occurs at elevated frequency because of a founder missense allele, **c.92C>T p.(Ser31Leu)** in *ZNHIT3* (NM_004773.3). Biallelic loss-of-function of ZNHIT3 is the primary cause; the first non-Finnish patient was a compound heterozygote carrying p.Ser31Leu together with a second rare allele p.Cys14Phe.

Mechanistically, PEHO syndrome is a **ribosomopathy**. ZNHIT3 is an evolutionarily conserved assembly chaperone required for incorporation of **box C/D small nucleolar RNAs (snoRNAs)** into functional snoRNP complexes that direct **2′-O-methylation of ribosomal RNA** and ribosome biogenesis. PEHO-modeling mutations destabilize the protein, lower box C/D snoRNA levels, impair rRNA processing, alter site-specific 2′-O-methylation, and perturb translation. In the developing cerebellum this produces **nucleolar stress** and activation of the **p53/p21 pathway**, driving apoptosis of cerebellar granule-cell progenitors. The resulting selective loss of inner granular layer neurons, with relative preservation of Purkinje cell number, is the pathological hallmark, matched clinically by profound hypotonia, absent developmental progress, optic atrophy and progressive infratentorial atrophy on neuroimaging.

There is **no curative therapy**; management is entirely supportive (seizure control, feeding support, physiotherapy, and family genetic counseling). However, the discovery of the nucleolar-stress → p53/p21 → granule-cell apoptosis axis is therapeutically significant: in conditional *Znhit3* knockout mice, **genetic or pharmacologic inhibition of p53/p21 signaling rescued cerebellar granule-cell development**, nominating this axis as a candidate target for future disease-modifying intervention. Prognosis is poor, with profound disability and death typically in infancy or childhood.

Key Findings

Finding 1 — PEHO is caused by biallelic ZNHIT3 loss-of-function, with a Finnish founder allele p.Ser31Leu

Homozygosity mapping combined with exome sequencing in Finnish families identified a homozygous missense substitution, **c.92C>T p.(Ser31Leu)** in *ZNHIT3* (NM_004773.3), as the primary cause of PEHO syndrome (PMID: 28335020). The affected residue lies in a conserved region of the zinc-finger HIT domain. The landmark report stated that *"a homozygous missense substitution of leucine for serine at codon 31 in ZNHIT3 was identified as the primary cause of PEHO syndrome,"* and functional work showed that *"transfection of cell lines with ZNHIT3 expression vectors showed that the PEHO syndrome mutant protein is unstable"* — establishing a **loss-of-function mechanism driven by protein instability**.

The disease is not exclusively Finnish. The first described non-Finnish case was a **compound heterozygote** carrying the founder p.Ser31Leu allele together with a novel second variant, **c.41G>T p.(Cys14Phe)** (PMID: 31048081): *"We identified biallelic missense variants in ZNHIT3 gene: the c.92C > T p.(Ser31Leu) variant (NM_004773.3), which is described previously as causing PEHO syndrome and the second novel variant c.41G > T p.(Cys14Phe)."* The c.41G>T allele was extremely rare (only 8 heterozygous carriers in gnomAD), consistent with a pathogenic, deleterious variant.

Variant summary table

Variant (cDNA)	Protein	Zygosity context	Population notes	Classification
c.92C>T	p.Ser31Leu	Homozygous (Finnish founder); also compound het	Enriched in Finland	Pathogenic
c.41G>T	p.Cys14Phe	Compound het with p.Ser31Leu	8 het carriers in gnomAD	Likely pathogenic
Novel biallelic variants (prenatal hydrops family)	—	Biallelic	Single family	Pathogenic (per P 39252897)

Variant types reported to date are predominantly **missense** affecting conserved zinc-finger residues, acting as partial loss-of-function/hypomorphic alleles (see Finding 7). Germline origin; no somatic contribution is relevant for this Mendelian disorder.

Finding 2 — PEHO is a ribosomopathy: ZNHIT3 loss disrupts box C/D snoRNP assembly, rRNA 2′-O-methylation, and triggers nucleolar stress → p53/p21

ZNHIT3 is *"an evolutionarily conserved protein required for ribosome biogenesis by mediating the assembly of small nucleolar RNAs (snoRNAs) of class C/D into ribonucleoprotein complexes (snoRNPs)"* (PMID: 39252897). Box C/D snoRNPs guide site-specific **2′-O-methylation** of rRNA, a modification essential for correct ribosome assembly and translational fidelity.

Yeast modeling using the ZNHIT3 ortholog **Hit1** demonstrated the molecular consequence directly: *"missense mutations modeling those found in PEHO syndrome patients cause a decrease in steady-state Hit1 protein levels, a significant reduction of box C/D snoRNA levels, and subsequent defects in rRNA processing and altered cellular translation"* (PMID: 35843310). RiboMethSeq analyses in that work showed altered site-specific 2′-O-methylation, linking ZNHIT3 dysfunction to a measurable defect in the rRNA modification landscape and in translation.

The in vivo consequence in the developing cerebellum was defined in conditional knockout mice: *"Transcriptomic analyses revealed activation of the p53/p21 pathway, rRNA processing defects, and nucleolar stress. Genetic or pharmacologic inhibition of p53/p21 signaling rescued granule cell progenitor development"* (PMID: 41857137). This establishes the complete molecular-to-cellular causal chain and identifies **p53/p21 as a druggable node**.

Finding 3 — Neuropathology: selective inner granular layer neuron loss with cerebellar and optic nerve atrophy

Autopsy study of 8 PEHO cases (Haltia & Somer, 1993) documented macroscopic cerebral and pronounced **cerebellar atrophy**, with histopathology concentrated in the cerebellar cortex and optic nerve (PMID: 8460530). The hallmark lesion is severe granule neuron depletion: *"There was a severe neuronal loss in the inner granular layer of the cerebellum. The Purkinje cells were relatively preserved in number although reduced in size, deformed and slightly disaligned."* Purkinje cells showed horizontally oriented dendrites, axonal torpedoes, and aberrant 200-kDa neurofilament immunoreactivity in perikarya, indicating cytoskeletal disorganization; the molecular layer was narrow and optic nerves atrophic.

The pathology is phenocopied in animal models: *"Both knockdown and genome editing of znhit3 in zebrafish embryos recapitulate the patients' cerebellar defects, microcephaly and oedema"* (PMID: 28335020), and rescue was achieved with wild-type but not mutant human ZNHIT3 mRNA, confirming variant pathogenicity.

Finding 4 — PEHO vs PEHO-like: genetic heterogeneity and differential diagnosis

A scoping review of 38 articles found that **cerebellar atrophy is the principal feature distinguishing true PEHO from PEHO-like syndromes**, while the classically "essential" criteria of optic atrophy and infantile spasms/hypsarrhythmia were less consistent than previously assumed (PMID: 33152950): *"Cerebellar atrophy was the main clinical difference between the two groups but data on optic atrophy and infantile spasms/hypsarrhythmia were not consistent with the previously essential diagnostic criteria."*

Several genetically distinct conditions mimic PEHO and must be excluded:

Condition	Gene	Inheritance	Distinguishing clue
True PEHO	*ZNHIT3*	AR	Extreme cerebellar granule-cell atrophy; ZNHIT3 biallelic variants
PEHO-like	*CCDC88A* (Girdin)	AR (biallelic LoF)	Cortical malformations (e.g., subcortical band heterotopia); PMIDs 30392057, 37798908, 39334473
"PEHO" phenocopy	*KIF1A* (motor domain)	De novo dominant	Dominant variant; may show complex IV deficiency; P 30385166
PCH6	*RARS2*	AR	Elevated lactate in blood/CSF; later-onset hypotonia; P 31536827

Of the PCH6 differential: *"Features distinguishing the proband's phenotype from PEHO syndrome were later appearance of hypotonia and elevated lactate levels in blood and cerebrospinal fluid"* (PMID: 31536827). This heterogeneity makes **molecular confirmation of ZNHIT3 essential** for a definitive PEHO diagnosis.

Finding 5 — Clinical diagnostic criteria and phenotype frequencies (Somer 1993)

The foundational clinical delineation (Somer, 1993) applied strict criteria: of 21 candidates, only 10 fulfilled true PEHO (PMID: 8301648). Key quantitative features: *"All were abnormal at birth showing hypotonia, drowsiness, or poor feeding. Head circumference was normal, but usually dropped to 2 SD below average during the first year of life. Visual fixation was either absent from birth or lost during the first months of life. Nine patients had peripheral oedema in early childhood. The mean age of onset of infantile spasms was 4.9 months."* All were extremely hypotonic, reached no motor milestones, and had brisk patellar reflexes; evoked potential studies showed abnormal brainstem/somatosensory responses with absent cortical SSEP. Nineteen PEHO patients were described in 14 Finnish families, consistent with autosomal recessive inheritance.

The phenotype was confirmed in non-Finnish (Australian/international) series (PMID: 12949965): *"Affected infants have facial dysmorphism and suffer from severe hypotonia, profound mental retardation, convulsions (often with a hypsarrhythmic EEG pattern), transient or persistent peripheral oedema, and optic atrophy. Cerebellar and brainstem atrophy are usually present on neuroimaging."*

Phenotype → HPO term suggestions with frequency

Phenotype	HPO term (suggested)	Onset	Frequency
Severe hypotonia	HP:0001290 (generalized hypotonia)	Neonatal	~100%
Profound intellectual disability / no development	HP:0002187	Infantile	~100%
Infantile spasms / hypsarrhythmia	HP:0012469 / HP:0002521	Mean 4.9 mo	High
Optic atrophy	HP:0000648	Birth–months	High
Acquired microcephaly	HP:0005484	First year	Typical (drops to −2 SD)
Peripheral/facial oedema	HP:0100820 / HP:0000316	Early childhood	~9/10
Cerebellar atrophy	HP:0001272	Early, progressive	Near-constant (defining)
Brainstem atrophy	HP:0002508	Early	Usual
Facial dysmorphism	HP:0001999	Congenital	Common
Abnormal evoked potentials / EEG	HP:0011185 (abnormal EEG)	Infantile	Reported

Finding 6 — Neuroradiology: early progressive infratentorial atrophy with abnormal myelination

Neuroimaging (CT/MR) of 21 clinical PEHO patients identified a core radiologic subgroup (PMID: 8352158): *"Twelve patients (group A) showed uniform changes with early progressive brain atrophy accentuated infratentorially, and abnormal myelination. The gyral pattern was*

normal." Group A (early progressive **infratentorial** — cerebellum/brainstem — atrophy, abnormal myelination, normal gyral pattern) corresponded on postmortem correlation to true PEHO, whereas group B (less progressive, supratentorial atrophy, abnormal gyri) was considered incompatible. A **normal gyral pattern with disproportionate cerebellar/brainstem atrophy** is therefore the characteristic imaging signature; in the genomic era, ZNHIT3 testing has superseded the historical requirement for neuropathology for conclusive diagnosis.

Finding 7 — Allelic severity spectrum: null lethality vs hypomorphic PEHO alleles; prenatal hydrops extends the phenotype

Complete *Znhit3* ablation in mouse **arrests embryos at the morula stage** with reduced translation and decreased snoRNA/rRNA abundance plus ribosome and splicing defects (PMID: 40178020): *"its ablation reduces protein translation and prevents mouse embryos development beyond the morula stage. The absence of ZNHIT3 leads to decreased snoRNA and rRNA abundance which causes defects of ribosomes and mRNA splicing."* This demonstrates that **null alleles are embryonic-lethal**, implying human PEHO missense alleles (e.g., p.Ser31Leu) are **partial loss-of-function/hypomorphic** — retaining enough residual function to permit birth but insufficient for normal cerebellar development.

The phenotypic spectrum extends prenatally: two fetuses with biallelic novel *ZNHIT3* variants presented with *"isolated hydrops during the early second trimester of pregnancy, resulting in intrauterine demise"* (PMID: 39252897), establishing a **severe prenatal/lethal hydrops form** at the severe end of the allelic series.

Finding 8 — Verified cross-reference identifiers

Confirmed via EBI OLS4 (MONDO:0009841, label "PEHO syndrome") and MyGene.info/Alliance of Genome Resources:

Resource	Identifier
MONDO	MONDO:0009841
OMIM (disease)	260565
Orphanet	ORPHA:2836
MeSH	C536317
MedGen / UMLS	342404 / C1850055
GARD	0004264
DOID	DOID:0080539
ICD-11 Foundation	976613527
Gene — HGNC	HGNC:12309
Gene — NCBI	9326
Gene — Ensembl	ENSG00000273611
Gene — UniProt	Q15649
Gene — OMIM	604500
Cytoband	17q12 (GRCh38 chr17:36,486,629–36,499,310)

Section-by-Section Report

1. Disease Information

PEHO syndrome is a rare, severe, autosomal-recessive infantile neurodegenerative encephalopathy caused by biallelic variants in **ZNHIT3**. The name is an acronym: **P**rogressive **E**ncephalopathy with (o)**E**dema, **H**ypsarrhythmia, and **O**ptic atrophy. Synonyms include *progressive encephalopathy–optic atrophy syndrome* and *progressive encephalopathy with edema, hypsarrhythmia and optic atrophy*. Identifiers are listed in Finding 8 (MONDO:0009841; OMIM:260565; ORPHA:2836; MeSH C536317; DOID:0080539; ICD-11 Foundation 976613527).

Information is derived from **aggregated disease-level resources** (OMIM, Orphanet) and **individual-patient case series/cohorts** (Finnish families, Australian series, individual case reports), together with model-organism and in vitro mechanistic studies.

2. Etiology

Causal factor: purely genetic — biallelic loss-of-function variants in *ZNHIT3*. **Genetic risk factors:** the **Finnish founder allele p.Ser31Leu (c.92C>T)** is the dominant risk allele in the Finnish population; rarer alleles (e.g., p.Cys14Phe) occur elsewhere. **Consanguinity / founder effect** is the principal population-genetic driver (homozygosity in Finnish families; founder enrichment). No environmental, lifestyle, infectious, toxic, or occupational risk factors are implicated — it is a monogenic Mendelian disorder. No **protective alleles or environmental protective factors** are known. **Gene–environment interactions** are not applicable. Carrier parents are unaffected heterozygotes.

3. Phenotypes

See Finding 5 table for the full phenotype list, types, onset, frequencies, and HPO suggestions. In brief: neonatal-onset severe generalized hypotonia (HP:0001290), profound developmental arrest (HP:0002187), infantile spasms with hypsarrhythmia beginning on average ~4.9 months (HP:0012469/HP:0002521), optic atrophy with absent visual fixation (HP:0000648), acquired microcephaly dropping to ~−2 SD in the first year (HP:0005484), transient/persistent peripheral and facial oedema (HP:0100820/HP:0000316, ~9/10 patients), facial dysmorphism (HP:0001999), and near-constant progressive cerebellar (HP:0001272) and brainstem (HP:0002508) atrophy. Severity is **severe-to-profound and progressive**; the disease is uniformly life-limiting. **Quality-of-life impact** is maximal: affected children reach no motor milestones, are non-verbal, cortically blind, dependent for all care, and require feeding and seizure management.

4. Genetic / Molecular Information

Causal gene: *ZNHIT3* (HGNC:12309; NCBI 9326; OMIM gene 604500; UniProt Q15649; 17q12). **Pathogenic variants:** predominantly **missense** affecting conserved zinc-finger residues — p.Ser31Leu (c.92C>T, founder) and p.Cys14Phe (c.41G>T), plus additional novel biallelic variants (prenatal hydrops family). Classification: p.Ser31Leu **pathogenic**; p.Cys14Phe **likely pathogenic** (gnomAD ~8 het carriers). **Allele frequency:** all pathogenic alleles are rare; p.Ser31Leu is enriched by founder effect in Finland. **Origin:** germline. **Functional consequence:** loss-of-function via protein instability (hypomorphic — null is embryonic-lethal;

Finding 7). **Modifier genes:** none established. **Epigenetic information:** none specific to PEHO; the mechanism involves rRNA 2′-O-methylation (an RNA modification) rather than DNA methylation. **Chromosomal abnormalities:** none — PEHO is a single-gene disorder, not a copy-number/structural syndrome.

5. Environmental Information

Not applicable. No environmental, lifestyle, toxicological, or infectious contributors are known or expected for this Mendelian ribosomopathy.

6. Mechanism / Pathophysiology

Ordered causal chain (initiating lesion → clinical manifestation):

1. **Biallelic hypomorphic *ZNHIT3* missense variants** (e.g., p.Ser31Leu) **lead to** reduced steady-state ZNHIT3 protein (protein instability; demonstrated in cell transfection and yeast Hit1 models).
2. Reduced ZNHIT3 **results in** impaired assembly of **box C/D snoRNAs into snoRNPs** (ZNHIT3 is the required assembly chaperone).
3. Defective snoRNP assembly **leads to** reduced box C/D snoRNA levels and **impaired site-specific 2′-O-methylation of rRNA** plus **rRNA processing defects** (demonstrated by RiboMethSeq in yeast; inferred to operate identically in human neural progenitors).
4. Defective ribosome biogenesis **results in altered/compromised translation** and **nucleolar stress** (demonstrated in yeast, mouse embryo, and conditional-KO mouse cerebellum).
5. Nucleolar stress **activates the p53/p21 pathway** (demonstrated transcriptomically in conditional *Znhit3*-KO cerebellum).
6. p53/p21 activation **drives apoptosis of cerebellar granule-cell progenitors** (the most proliferative, ribosome-demanding neuronal population) — **branch point**: the high biosynthetic demand of granule-cell progenitors is inferred to explain their selective vulnerability.
7. Granule-cell progenitor death **results in** severe **inner granular layer neuron loss** with secondary Purkinje-cell deformation and **cerebellar + brainstem + optic nerve atrophy** (demonstrated at autopsy and by neuroimaging).
8. Loss of cerebellar/visual/brainstem circuitry **manifests as** the clinical PEHO phenotype: hypotonia, developmental arrest, infantile spasms/hypsarrhythmia, optic atrophy, and progressive microcephaly.

Rescue branch: Genetic or pharmacologic **inhibition of p53/p21** restores granule-cell progenitor development in mice (PMID: 41857137), confirming steps 5–7 are causal and druggable.

```
ZNHIT3 LoF (unstable protein)
        |
        ▼
Box C/D snoRNP assembly failure ──► ↓snoRNAs ──► ↓rRNA 2'-O-methylation + rRNA processing defect
        |
        ▼
Impaired ribosome biogenesis / translation ──► NUCLEOLAR STRESS
        |
        ▼
   p53 / p21 activation ──(pharmacologic/genetic block = RESCUE)──► ✗
        |
        ▼
Granule-cell progenitor APOPTOSIS (selective vulnerability of high-biosynthesis cells)
        |
        ▼
Inner granular layer loss → cerebellar/brainstem/optic atrophy → PEHO clinical phenotype
```

Molecular pathways: ribosome biogenesis (box C/D snoRNP assembly), p53/p21 (TP53–CDKN1A) stress signaling. **Cellular processes:** nucleolar stress response, apoptosis, cell-cycle arrest, neurodevelopment. **Protein dysfunction:** loss-of-function via misfolding/instability of the ZNHIT3 zinc-finger. **GO term suggestions:** GO:0000492 (box C/D snoRNP assembly), GO:0042254 (ribosome biogenesis), GO:0031167 (rRNA methylation), GO:0006364 (rRNA processing), GO:0072331 (signal transduction by p53 class mediator), GO:0006915 (apoptotic process), GO:0021953 (central nervous system neuron differentiation). **CL term suggestions:** CL:0000120 (cerebellar granule cell), CL:0000121 (Purkinje cell), CL:0000047 (neuronal stem/progenitor cell). **CHEBI:** methylated-nucleotide / 2′-O-methyl modification entities.

7. Anatomical Structures Affected

Organ/system: central nervous system (nervous system). **Primary sites:** cerebellum — specifically the **inner granular layer** (UBERON:0002956 cerebellar cortex; granular layer), **brainstem** (UBERON:0002298), and **optic nerve** (UBERON:0000941). Secondary: cerebral cortex atrophy, corpus callosum hypoplasia. **Tissue/cell level:** nervous tissue; **cerebellar granule neurons** (CL:0000120) are the selectively depleted population, with secondary Purkinje cell (CL:0000121) deformation. **Subcellular level:** the **nucleolus** (GO:0005730) is the

locus of pathology (nucleolar stress; ribosome assembly); nucleus and ribosome (GO:0005840) are involved. **Localization/lateralization:** bilateral, symmetric atrophy; infratentorial predominance with a normal supratentorial gyral pattern.

8. Temporal Development

Onset: congenital/neonatal — infants are abnormal at birth (hypotonia, drowsiness, poor feeding). **Pattern:** chronic, relentlessly **progressive**. Head circumference is normal at birth then falls to ~−2 SD during the first year (acquired microcephaly); infantile spasms emerge at mean ~4.9 months; visual fixation is absent from birth or lost in the first months. **Progression rate:** early and progressive, with uniform infratentorial atrophy on serial imaging. **Course:** progressive neurodegeneration without remission; lifelong (and shortened). A **prenatal critical window** exists at the severe end (second-trimester hydrops with intrauterine demise). The developmental window of granule-cell proliferation is the critical period of vulnerability — and, per mouse data, the window in which p53/p21 modulation could theoretically intervene.

9. Inheritance and Population

Inheritance: autosomal recessive; carrier parents unaffected. **Penetrance:** complete in biallelic individuals (with variable severity/expressivity across the allelic series — from classic PEHO to prenatal lethal hydrops). **Epidemiology:** rare; best characterized in **Finland**, where 19 patients in 14 families were described, reflecting a **founder effect** (p.Ser31Leu). Precise global prevalence/incidence figures are not well established (ultra-rare; Orphanet ORPHA:2836). **Founder effect/consanguinity:** central to occurrence (Finnish founder allele; consanguinity in some non-Finnish families, particularly for PEHO-like *CCDC88A* cases). **Carrier frequency:** not precisely quantified outside Finland. **Sex ratio:** no strong sex predilection reported; autosomal. **Geographic distribution:** founder enrichment in Finland; sporadic cases worldwide (e.g., Australia; Middle East for PEHO-like). **Genetic anticipation / mosaicism:** not applicable/not reported.

10. Diagnostics

Definitive diagnosis is by **molecular genetic testing** of *ZNHIT3* (single-gene testing, epilepsy/neurodevelopmental gene panels, or whole-exome/whole-genome sequencing), which has superseded the historical requirement for neuropathology. **Clinical criteria** (Somer 1993): neonatal abnormality (hypotonia/drowsiness/poor feeding), acquired microcephaly, infantile spasms with hypsarrhythmic EEG, optic atrophy/absent visual fixation, peripheral oedema,

and progressive cerebellar/brainstem atrophy. **Imaging:** MRI showing early progressive infratentorial atrophy with abnormal myelination and normal gyral pattern (Finding 6). **Electrophysiology:** EEG hypsarrhythmia; abnormal brainstem auditory and somatosensory evoked potentials with absent cortical SSEP. **Laboratory:** no specific biomarker; **normal lactate** helps distinguish from PCH6/RARS2 (elevated lactate). **Differential diagnosis:** PEHO-like (*CCDC88A* — look for cortical malformations), *KIF1A*-related (de novo dominant; consider mitochondrial complex IV deficiency), PCH6 (*RARS2*; elevated lactate), and other early infantile epileptic encephalopathies (Findings 4, 5). **Screening:** carrier/cascade testing within affected families; prenatal/preimplantation testing feasible where the familial variants are known.

11. Outcome / Prognosis

Prognosis is **poor**. The disease causes profound, permanent disability — no developmental milestones, cortical blindness, intractable epilepsy, and complete care dependence. Progressive neurodegeneration leads to **death in infancy or childhood** (and intrauterine demise in the severe prenatal hydrops form). There is no recovery potential and no disease-modifying therapy currently. **Complications:** feeding difficulty/aspiration, respiratory infections, intractable seizures. **Prognostic factors:** genotype/allele severity (hypomorphic PEHO alleles vs more severe biallelic variants associated with prenatal hydrops). No validated prognostic biomarkers exist beyond the genotype and imaging burden.

12. Treatment

There is **no curative or disease-modifying therapy**. Management is **supportive and symptomatic**: - **Anticonvulsants** for infantile spasms/epilepsy (e.g., ACTH/vigabatrin and other antiseizure medications per standard infantile-spasm protocols) — symptom control, not disease-modifying. - **Nutritional support** (gastrostomy feeding for dysphagia). - **Physiotherapy, occupational and visual support**, and management of oedema. - **Genetic counseling** for families.

Experimental/translational direction: The mouse finding that **genetic or pharmacologic inhibition of p53/p21 signaling rescues granule-cell progenitor development** (PMID: 41857137) nominates **p53/p21 pathway inhibition** as a candidate therapeutic strategy — currently preclinical only, with no clinical trials (no NCT identifiers) for ZNHIT3-PEHO. No pharmacogenomic, gene-therapy, cell-therapy, RNA-based, or immunotherapy approaches are established. **NCIT suggestions:** supportive care, anticonvulsant therapy, nutritional support, physical therapy, genetic counseling.

13. Prevention

No primary prevention exists for this genetic disease. **Prevention centers on reproductive genetics: genetic counseling**, **carrier/cascade screening** of relatives of affected individuals, and **prenatal diagnosis or preimplantation genetic testing** where the familial *ZNHIT3* variants are known. In founder populations (Finland), awareness of p.Ser31Leu supports targeted carrier testing. **Tertiary prevention** = anticipatory management of complications (seizures, aspiration, infections). No immunization, behavioral, or public-health/environmental interventions are applicable.

14. Other Species / Natural Disease

No naturally occurring companion-animal or wildlife disease equivalent to PEHO is documented (no OMIA entry noted). **Orthologous genes** (relevant for modeling): mouse *Znhit3*, zebrafish *znhit3*, and the budding-yeast ortholog **Hit1** (*Saccharomyces cerevisiae*). The gene and its ribosome-biogenesis function are **evolutionarily conserved** from yeast to human, which is why yeast, zebrafish, and mouse faithfully model the molecular defect. No zoonotic or cross-species transmission is relevant (non-infectious genetic disease).

15. Model Organisms

Model	Type	Construct	Phenotype recapitulation	Key limitation	PMID
Yeast (Hit1)	In vitro/ cellular	PEHO-modeling missense mutations	↓ protein, ↓ box C/D snoRNA, rRNA processing + 2′-O-methylation/ translation defects	No neural phenotype	35843310
Zebrafish (*znhit3*)	Vertebrate	Knockdown + CRISPR/genome editing	Cerebellar defects, microcephaly, oedema; rescued by WT (not mutant) human ZNHIT3	Developmental not progressive adult model	28335020
Mouse — complete KO	Mammalian	Null ablation	Embryonic arrest at morula stage; ↓ translation, ↓ snoRNA/rRNA, ribosome/ splicing defects	Lethal — cannot model postnatal disease	40178020
Mouse — conditional KO	Mammalian	Spatiotemporal (cerebellar) KO	Granule-cell apoptosis, nucleolar stress, p53/p21 activation; rescued by p53/ p21 inhibition	Conditional context vs human hypomorphic allele	41857137

The **conditional mouse KO** is the most disease-relevant model (recapitulates granule-cell death and reveals the therapeutic p53/p21 node), while the **complete KO** defines the null-lethal severe boundary that frames human alleles as hypomorphic. **Resources:** MGI (*Znhit3*), ZFIN (*znhit3*), SGD (*HIT1*).

Mechanistic Model / Interpretation

PEHO syndrome is best understood as a **tissue-selective ribosomopathy**. The unifying model: a conserved ribosome-assembly chaperone (ZNHIT3) is partially lost; the most proliferative, biosynthesis-hungry neural population (cerebellar granule-cell progenitors) cannot tolerate the resulting ribosome deficit; nucleolar stress converges on p53/p21 and kills those progenitors; and the loss of the cerebellar granular layer, together with brainstem and optic nerve degeneration, produces the clinical syndrome. The strength of this model is that every link has direct experimental support across four orthogonal systems (yeast biochemistry, zebrafish development, mouse embryo, conditional mouse cerebellum), and the final link is **reversible in mice** by blocking p53/p21.

Two insights are particularly consequential for a knowledge base: 1. **Genotype–severity gradient.** Null = embryonic lethal; hypomorphic missense = classic PEHO; certain biallelic combinations = prenatal lethal hydrops. Severity tracks residual ZNHIT3/ribosome function. 2. **Therapeutic node.** The p53/p21 axis is downstream of the untreatable primary lesion yet upstream of the irreversible cell death — making it the rational point of intervention.

Evidence Base

PMID	Role	Contribution
28335020	Landmark (gene discovery)	Identifies ZNHIT3 p.Ser31Leu as cause; protein instability; zebrafish rescue
31048081	Supporting	First non-Finnish compound-het case (p.Ser31Leu/p.Cys14Phe)
35843310	Mechanism	Yeast Hit1 — snoRNA/rRNA/2′-O-methylation/ translation defects
41857137	Mechanism + therapy	Conditional mouse KO — p53/p21 activation; rescue by inhibition
39252897	Mechanism + phenotype	ZNHIT3 snoRNP function; prenatal hydrops extension
40178020	Allelic severity	Complete KO embryonic-lethal → human alleles hypomorphic
8460530	Neuropathology	Inner granular layer loss; Purkinje preservation
8301648	Clinical criteria	Diagnostic criteria, onset ages, frequencies (10/21 true PEHO)
8352158	Neuroradiology	Early progressive infratentorial atrophy signature
12949965	Clinical	Non-Finnish phenotype confirmation
33152950	Review	PEHO vs PEHO-like distinction (cerebellar atrophy)
31536827	Differential	PCH6/RARS2 — elevated lactate
30392057, 37798908, 39334473	Differential	CCDC88A PEHO-like syndrome
30385166, 34916088	Differential	KIF1A-related PEHO phenocopy

Limitations and Knowledge Gaps

- **Epidemiology is imprecise.** Exact global prevalence, incidence, and carrier frequency (outside the Finnish founder population) are not well quantified.
- **Mechanistic inference across species.** The snoRNA/rRNA 2′-O-methylation defect is directly shown in yeast; its precise quantitative operation in human cerebellar progenitors is inferred, not directly measured in patient tissue.
- **Selective vulnerability** of granule cells is biologically plausible (high biosynthetic demand) but not formally proven to be the sole determinant.
- **Therapeutic translation is preclinical.** p53/p21 inhibition rescue is demonstrated only in conditional mouse KO; no human trials, no safety/efficacy data, and timing (critical developmental window) is a major obstacle.
- **Genotype–phenotype map is sparse** — few pathogenic alleles are described; correlations between specific variants and severity (classic vs prenatal-lethal) rest on small numbers.
- **No validated biomarkers** for diagnosis monitoring or prognosis beyond genotype and imaging.

Proposed Follow-up Experiments / Actions

1. **Patient-derived models:** Generate iPSC-derived cerebellar organoids from PEHO (p.Ser31Leu) patients to directly measure rRNA 2′-O-methylation (RiboMethSeq), nucleolar stress, and p53/p21 activation in human neural tissue.
2. **Therapeutic validation:** Test pharmacologic p53/p21 inhibitors (and nucleolar-stress modulators) in conditional *Znhit3* mice for dose, timing window, and off-target/tumorigenic risk before considering translation.
3. **Allelic series functional assays:** Systematically quantify residual ZNHIT3 function/stability across reported and novel variants to build a predictive genotype–severity map (classic PEHO vs prenatal hydrops).
4. **Registry and natural history:** Establish an international *ZNHIT3*-PEHO registry to define prevalence, survival, and genotype–phenotype correlations with sufficient power.
5. **Biomarker discovery:** Explore whether ribosome/snoRNA signatures or nucleolar-stress markers are detectable in accessible patient samples (CSF, blood) for diagnosis/monitoring.

6. **Diagnostic optimization:** Ensure *ZNHIT3* is included on infantile epileptic encephalopathy and cerebellar atrophy gene panels, with reflex lactate testing to triage the RARS2/PCH6 differential.

Consensus Answer

PEHO syndrome (MONDO:0009841; OMIM:260565) is a severe autosomal-recessive early-infantile progressive neurodegenerative encephalopathy caused by biallelic loss-of-function variants in **ZNHIT3** (HGNC:12309; 17q12), most commonly the Finnish founder allele p.Ser31Leu. It is a ribosomopathy in which reduced/unstable ZNHIT3 impairs box C/D snoRNP assembly and rRNA 2′-O-methylation, causing nucleolar stress and p53/p21-dependent apoptosis of cerebellar granule-cell progenitors — producing the hallmark cerebellar atrophy together with profound hypotonia, infantile spasms/hypsarrhythmia, optic atrophy, oedema, and acquired microcephaly. There is no curative therapy (management is supportive), though preclinical mouse data show p53/p21 inhibition rescues the cerebellar phenotype, nominating a candidate therapeutic target.
