

Hypothesis-Search Report: Ibogaine GDNF/VTA Induced-Neuroplasticity Model of Opioid Use Disorder

Hypothesis ID: `ibogaine_gdnf_vta_neuroplasticity_model` **Disease:** Opioid Use Disorder (OUD), Psychiatric **KB status:** EMERGING **Search date:** 2026-08-28

Executive Judgment

Verdict: PARTIALLY SUPPORTED / UNRESOLVED for opioid use disorder.

The mechanistic core of the model — that ibogaine induces GDNF in the ventral tegmental area (VTA), activates Ret/ERK-MAPK signalling there, and thereby produces an anti-addiction effect that outlasts drug exposure — is **directly and rigorously established, but only for the ethanol reinforcer** (He et al. 2005,

[P 15659598](#)

sufficiency by intra-VTA GDNF microinjection *and* necessity by intra-VTA anti-GDNF antibody blockade). Its extension to opioids is currently an **inference across reinforcers**, not a demonstrated causal chain.

Three lines of evidence make the extension *plausible*: 1. VTA GDNF/Ret signalling is a general negative regulator of drug adaptations, including **morphine** (Messer et al. 2000,

[P 10798408](#)

Niwa et al. 2007,

[P 17538232](#)

2. Single-dose ibogaine durably reduces **morphine self-administration** in rats, with an aftereffect outlasting drug elimination (Glick et al. 1991,

[P 1868880](#)

3. Single-dose ibogaine durably reduces opioid craving/use in **humans** (Brown & Alper 2018,

P 28541119

~24-week

effect,

P 38518270

However, the **defining experiment of the model — intra-VTA anti-GDNF blockade of ibogaine's effect on opioid self-administration/relapse — has never been performed.**

Meanwhile, two competing mechanisms are *better evidenced in the opioid domain specifically*: - **Habenular $\alpha 3\beta 4$ nicotinic route**: the iboga congener 18-MC reduces morphine self-administration when injected into the medial habenula/interpeduncular nucleus but has **no effect from the VTA** (Glick et al. 2006,

P 16626688

and

does

not

raise

GDNF

(

P 21040239

- **G-protein-biased κ -opioid agonism**: noribogaine is a biased κ -agonist (Maillet et al. 2015,

P 26302653

and κ -agonist oxa-iboga analogs give durable single-dose suppression of heroin/fentanyl intake (Havel et al. 2024,

P 39304653

Finally, the human neurotrophic signal that tracks opioid craving is **BDNF, not GDNF** (Heberlein et al. 2011,

P 21890593

and rodent ibogaine induces BDNF broadly across NAcc/SN/PFC

(

P 30890941

qualifying the model's attribution of durability *specifically* to VTA GDNF.

Bottom line: The GDNF/VTA neuroplasticity model is a mechanistically coherent, experimentally validated account **for alcohol** and a plausible but **unproven** account **for opioids**. For OUD it should remain **EMERGING**, explicitly flagged as reinforcer-transferred inference, with the $\alpha 3\beta 4$ and κ models as co-equal (arguably better-evidenced) opioid-domain competitors.

Evidence Matrix

#	Citation (PMID)	Evidence type	Stance	Mechanistic claim tested	Key finding	Context / subtype
1	15659598 (He 2005)	Model organism + in vitro	Support (partial for OUD)	Intra-VTA GDNF is necessary & sufficient for ibogaine's anti-SA effect; ibogaine→Ret/ERK1	Intra-VTA GDNF mimics, anti-GDNF antibody blocks ibogaine's effect; phospho-Ret/ERK1 ↑ in SH-SY5Y	Ethanol SA, rat / cell line
2	18541917 (Carnicella 2008)	Model organism	Support (partial for OUD)	VTA GDNF→MAPK reduces consumption & relapse	Intra-VTA GDNF rapidly, dose-dependently ↓ ethanol SA and blocks reacquisition; MAPK-dependent	Ethanol, rat
3	21040239 (Carnicella 2010)	In vitro + model organism	Support / discriminator	Noribogaine (not 18-MC) drives GDNF; VTA site	Noribogaine ↑ GDNF mRNA & ↓ intra-VTA ethanol SA; 18-MC does neither	Ethanol; SH-SY5Y
4	10798408 (Messer 2000)	Model organism	Support (opioid bridge)	VTA GDNF modulates opioid adaptations	Intra-VTA GDNF blocks morphine biochemical adaptations; anti-GDNF Ab & GDNF+/- enhance drug responses; morphine ↓ VTA phospho-Ret	Morphine + cocaine, rat/mouse

#	Citation (PMID)	Evidence type	Stance	Mechanistic claim tested	Key finding	Context / subtype
5	17538232 (Niwa 2007)	Model organism	Support (opioid bridge)	GDNF induction blocks morphine reward	Leu-Ile ↓ morphine CPP/sensitization; absent in GDNF+/- mice	Morphine, mouse
6	1868880 (Glick 1991)	Model organism	Support (phenotype, opioid)	Single ibogaine durably ↓ opioid SA	Ibogaine ↓ morphine SA with aftereffect outlasting elimination; persists days-weeks; reinforcer-specific	Morphine SA , rat
7	33299186 (Cameron 2021)	Model organism	Support (qualifies locus)	Plasticity separable from hallucinogenesis	TBG promotes structural plasticity, ↓ heroin-seeking, non-hallucinogenic	Heroin-seeking, mouse
8	30890941 (Marton 2019)	Model organism	Qualifies	Neurotrophic response is GDNF-specific to VTA	Ibogaine ↑ BDNF broadly (NAcc, SN, PFC) and GDNF/BDNF across regions	Rat, no drug challenge
9	39304653 (Havel 2024)	Model organism + in vitro	Competing (κ)	κ-agonism drives durable opioid suppression	κ-agonist oxa-iboga analogs give long-lasting single-dose ↓ of morphine/heroin/fentanyl	Opioid SA/relapse, male rat

#	Citation (PMID)	Evidence type	Stance	Mechanistic claim tested	Key finding	Context / subtype
					intake & seeking	
10	26302653 (Maillet 2015)	In vitro / computational	Competing (κ)	Noribogaine is a κ substrate	Noribogaine = G-protein-biased κ agonist ($EC_{50} \approx 9 \mu M$), weak μ antagonist	Molecular pharmacology
11	9668680 (Glick & Maisonneuve 1998)	Review / model organism	Competing (κ /NMDA)	Opioid effect via κ + NMDA	Authors attribute opioid-SA effects to κ -agonism + NMDA antagonism; serotonergic route for alcohol	Rat SA
12	16626688 (Glick 2006)	Model organism	Competing ($\alpha 3\beta 4$), site challenge	Anti-morphine effect maps to habenula, not VTA	18-MC in MHb/IPN $\downarrow$ morphine SA; VTA administration had no effect; sucrose unaffected	Morphine SA rat
13	17447255 (Taraschenko 2007)	Model organism	Competing ($\alpha 3\beta 4$)	Habenular $\alpha 3\beta 4$ gates opioid DA	18-MC in MHb/IPN blocks sensitized accumbens DA to morphine	Morphine, rat
14		Model organism	Competing ($\alpha 3\beta 4$)	Withdrawal via habenula/LC	Intracerebral 18-MC	

#	Citation (PMID)	Evidence type	Stance	Mechanistic claim tested	Key finding	Context / subtype
	16289028 (Panchal 2005)				attenuates precipitated morphine withdrawal	Morphine withdrawal, rat
15	28541119 (Brown & Alper 2018)	Human clinical (observational)	Support (phenotype)	Single-dose durable human effect	50% opioid-free at 1 mo; SOWS 31→14 (p<0.001); improvements to 12 mo	OUD, n=30
16	38518270 (Cherian, primer)	Review	Support (phenotype)	Durability of craving reduction	RCT/OL: craving ↓ ~50% up to 24 weeks	OUD
17	33620733 (Knuijver 2022)	Human clinical (observational)	Qualifies (safety)	Cardiac/cerebellar risk	QTc +95 ms mean, 50% >500 ms; severe ataxia	OUD, n=14
18	21890593 (Heberlein 2011)	Human clinical	Qualifies / refutes biomarker	GDNF as human opioid biomarker	Serum GDNF unchanged & craving-unassociated; BDNF elevated & craving-correlated	Opiate dependence, n=28
19	37937505 (Ona 2023)	Review	Qualifies	MoA is polypharmacology	Ibogaine acts via complex multi-target modulation, no single MoA	SUD

Mechanistic Causal Chain

Proposed chain (seed hypothesis): single ibogaine dose → (ibogaine + long-lived noribogaine) → ↑ GDNF transcription in VTA → GDNF binds GFR α 1/Ret → ↑ phospho-Ret → ↑ ERK1/2-MAPK → structural/functional remodeling of mesolimbic (VTA dopaminergic) circuitry → durable ↓ opioid self-administration & relapse (outlasting drug clearance), dissociable from acute withdrawal and from the psychedelic experience → sustained abstinence/reduced craving in OUD patients

Link-by-link strength:

Edge	Strength	Evidence
ibogaine/noribogaine → ↑ GDNF (VTA/midbrain)	Strong (in vitro + rodent)	P 15659598 P 21040239 (ethanol context; SH-SY5Y + midbrain)
GDNF → Ret/ERK-MAPK activation	Strong	P 15659598 (phospho-Ret, ERK1); P 18541917 (MAPK necessity for ethanol)
VTA GDNF/MAPK → reduced ethanol SA & relapse	Strong (sufficiency + necessity)	P 15659598 P 18541917
VTA GDNF → reduced opioid adaptation/ reward	Moderate (bridge; not ibogaine-driven)	P 10798408 P 17538232 (morphine CPP/biochemical, not SA-with-blockade)
ibogaine → durable ↓ opioid SA (phenotype)	Moderate-strong (mechanism-agnostic)	P 1868880 (rat), 28541119 (human)
VTA GDNF necessity for ibogaine's OPIOID effect	MISSING — never tested	<i>No study performs intra-VTA anti-GDNF blockade in opioid SA</i>
durability sustained by self-amplifying VTA GDNF loop	Inferred / unverified in this search	Proposed in alcohol literature (Barak et al.); not retrievable/confirmed here — treat as known-unknown
durable effect dissociable from psychedelic experience	Supported indirectly	Non-hallucinogenic 18-MC & TBG retain anti-opioid/anti-seeking effects (P 16626688) P 33299186

Edge	Strength	Evidence
durable effect dissociable from acute withdrawal	Plausible	Withdrawal maps to habenula/LC (P 16289028); durability is a separate axis

Where the chain is strong: the intracellular module (GDNF→Ret→ERK) and the *ethanol* behavioural endpoint. **Where it is inferred:** every opioid-specific edge. **Where it is missing:** the necessity test in opioids, and the molecular basis of *durability* (self-amplifying loop) in any reinforcer as confirmed in this search.

Knowledge Gaps

- Opioid necessity edge is untested (highest priority).**
- Scope:** Does intra-VTA anti-GDNF neutralizing antibody (or GFRα1/Ret knockdown) abolish ibogaine's effect on **opioid** self-administration/reinstatement?
- Why it matters:** This is the model's own defining/discriminating experiment; without it, extension to OUD is inference.
- Checked:** Searched ibogaine/morphine SA + GDNF/VTA; found the sufficiency/necessity loop only for ethanol
([P 15659598](#))
and mechanism-agnostic opioid durability
([P 1868880](#)).
- Resolution:** Run the He-2005 design with heroin/fentanyl SA + intra-VTA anti-GDNF.
- Durability mechanism (self-amplifying GDNF loop) unverified here.**
- Scope:** What sustains the effect for weeks after drug clearance?
- Why it matters:** Durability is the model's central selling point vs receptor occupancy.
- Checked:** Multiple queries for a VTA GDNF positive-feedback loop returned no retrievable record via this endpoint.

10. *Resolution:* Longitudinal VTA GDNF/phospho-Ret time-course after single ibogaine in opioid-experienced animals; curator should verify the alcohol-domain autoregulation literature (Barak et al., Nat Neurosci 2011) directly.

11. **Locus specificity: GDNF vs broader neurotrophic response.**

12. *Scope:* Is durability attributable *specifically* to VTA GDNF, or to distributed BDNF (NAcc/SN/PFC) and cortical 5-HT2A plasticity?

13. *Why it matters:*

P 30890941

(broad BDNF) and

P 33299186

(TBG cortical 5-HT2A) dilute VTA-GDNF specificity.

14. *Checked:* Yes; both surfaced.

15. *Resolution:* Region- and factor-specific knockdown (VTA-GDNF vs NAcc-BDNF vs PFC-5-HT2A) with opioid SA readout.

16. **Human GDNF biomarker is null.**

17. *Scope:* No human evidence links GDNF (central or peripheral) to opioid dependence or ibogaine response; the human craving-tracking neurotrophin is BDNF.

18. *Why it matters:* Source-level absence for the GDNF-specific claim in humans.

19. *Checked:*

P 21890593

(only human GDNF-in-opioid datapoint found; null).

20. *Resolution:* CSF/plasma GDNF & BDNF time-course in ibogaine-treated OUD patients correlated with abstinence.

21. **Competing-mechanism arbitration.**

22. *Scope:* κ -agonism

(

P 39304653

)

P 26302653

and habenular $\alpha 3\beta 4$

(
P 16626688)

each produce durable/robust anti-opioid effects; their contribution to *ibogaine*'s effect is unpartitioned.

23. *Resolution*: Factorial perturbation (κ antagonist, $\alpha 3\beta 4$ rescue, anti-GDNF) in the same ibogaine opioid-SA paradigm.

24. **Source/dataset absences (curation-relevant).**

25. No human genetic (GenCC/ClinGen-type) association of *GDNF/RET* with OUD surfaced.

26. No omics/cohort dataset tying VTA GDNF to human OUD surfaced.

27. Clinical evidence base is thin: essentially 3 RCTs total across iboga alkaloids

(
P 42117212),

none mechanistic.

Alternative Models

Model	Relationship to seed	Basis	Opioid-domain evidence
κ-opioid agonist model (noribogaine/oxa-iboga as biased κ agonists driving durable suppression)	Alternative (parallel mechanism for the same durable phenotype)	P 26302653 P 39304653 P 9668680	Strong & opioid-specific (durable single-dose ↓ heroin/fentanyl)
Habenular $\alpha 3\beta 4$ nicotinic model (18-MC blocks morphine SA via MHb/IPN)	Alternative (different circuit node; explicitly <i>not</i> VTA)	P 16626688 P 17447255 P 16289028	Strong & opioid-specific ; VTA-negative
Distributed BDNF / cortical 5-HT_{2A} psychoplastogen model	Complementary / broader (superset of VTA-GDNF)	P 30890941 P 33299186	Moderate (heroin-seeking ↓ with TBG)
NMDA antagonism (ibogaine/noribogaine NMDA block)	Parallel (acute, may aid withdrawal/tolerance)	P 9668680	Mechanistic, contributory
Serotonergic (SERT/5-HT) module	Parallel; authors assign this more to <i>alcohol</i> than opioids	P 9668680	Weak for opioids
Polypharmacology (no single MoA)	Meta-alternative — durable effect may be emergent synergy	P 37937505	Review-level

The GDNF/VTA model is best positioned as **one node in a polypharmacological network**, strongest for alcohol, and — for opioids — competing with the κ and $\alpha 3\beta 4$ accounts that currently have *more direct opioid-domain data*.

Discriminating Tests

1. **Necessity test (definitive).** Heroin/fentanyl IV self-administration in rats → single ibogaine (or noribogaine) → intra-VTA anti-GDNF neutralizing antibody vs IgG control. *Predicted (seed)*: antibody abolishes the durable aftereffect. *Predicted ($\kappa/\alpha3\beta4$)*: aftereffect intact. Add κ -antagonist (norBNI) and $\alpha3\beta4$ rescue arms for full factorial dissociation.
2. **Circuit-specific molecular knockdown.** AAV-shRNA against *Gdnf/Gfra1/Ret* restricted to VTA vs *Bdnf* in NAcc vs *Htr2a* in PFC; opioid reinstatement readout. Distinguishes locus.
3. **Longitudinal biomarker cohort.** OUD patients receiving ibogaine (where legal) → serial plasma/CSF GDNF & BDNF + resting-state mesolimbic connectivity vs abstinence at 4/12/24 weeks. Tests whether human GDNF dynamics (not baseline) track response.
4. **Non-hallucinogenic analog head-to-head.** TBG (5-HT2A/cortical) vs oxa-noribogaine (κ) vs 18-MC ($\alpha3\beta4$) in identical opioid-SA/relapse assay with GDNF/Ret readout — partitions which durable-effect route requires GDNF.
5. **Responder stratification.** Glick 1991's marked inter-individual variability suggests a responder genotype/phenotype; screen *GDNF/RET/OPRK1* variants against aftereffect magnitude.

Curation Leads (require curator verification)

Candidate	evidence	references	+	snippets	to	verify:	-
P 10798408	— "Infusion of GDNF into the ventral tegmental area (VTA)... blocks certain biochemical adaptations to chronic cocaine or morphine..." → adds opioid bridge edge (GDNF→morphine adaptation).						-
P 1868880	— "the aftereffect occurred at a time when ibogaine should have been entirely eliminated from the body..." → adds direct opioid durability phenotype (mechanism-agnostic).						-
P 16626688	— "Local administration of 18-MC into the ventral tegmental area had no effect on morphine self-administration." → add as competing/site-challenge edge (habenular $\alpha3\beta4$, VTA-negative).						-

P 26302653

— "noribogaine was a G-protein biased kappa agonist..." → add **competing κ** node. -

P 28541119

— "At 1-month posttreatment follow-up, 15 subjects (50%) reported no opioid use..." → add **human durability** evidence. -

P 21890593

— "GDNF serum levels ... did not differ significantly ... whereas ... BDNF ... associated with craving" → add **human GDNF-null biomarker** discussion prompt.

Candidate pathophysiology nodes/edges: - Node: `habenulo-interpeduncular α3β4 nAChR (MHb/IPN)` → competing anti-opioid circuit. - Node: `κ-opioid receptor (biased G-protein signalling)` → competing durable-suppression route. - Edge (proposed, unconfirmed): `VTA GDNF → durable ↓ opioid self-administration` — mark **UNCONFIRMED / needs direct perturbation**.

Candidate ontology terms: - Cell type: dopaminergic neuron (VTA) [CL:0000700]; processes: GO glial cell line-derived neurotrophic factor receptor signaling pathway (GO:0035860), MAPK cascade (GO:0000165), regulation of synaptic plasticity (GO:0048167).

Candidate subtype restriction / status: - Restrict the *validated* GDNF/VTA necessity claim to **ethanol/alcohol use disorder**; keep OUD extension **EMERGING** with an explicit `reinforcer_transfer_inference` flag.

Candidate knowledge_gaps prompts: - "Intra-VTA anti-GDNF blockade of ibogaine's effect on opioid self-administration is unperformed." - "Human GDNF is not a demonstrated biomarker of opioid dependence or ibogaine response (BDNF is)." - "No GDNF/RET human genetic association with OUD identified as of 2026-08-28." - "Durability mechanism (self-amplifying VTA GDNF loop) unverified in this search; verify Barak et al. alcohol-domain literature."

Artifact: Mechanistic Diagram (competing routes to durable opioid-seeking suppression)

```

flowchart TD
    IBO["Single ibogaine dose"] --> NOR["Noribogaine (long-lived metabolite, ~20 µM brain)"]
    IBO --> ACUTE["Acute NMDA antagonism / motor effects (transient)"]

    subgraph SEED["SEED MODEL – GDNF/VTA neuroplasticity"]
        NOR --> GDNF["↑ GDNF transcription in VTA"]
        IBO --> GDNF
        GDNF --> RET["GFRα1/Ret → phospho-Ret → ERK1/2-MAPK"]
        RET --> REMOD["Mesolimbic (VTA DA) circuit remodeling"]
        REMOD --> DUR["Durable ↓ self-administration / relapse"]
    end

    subgraph KAPPA["COMPETING – κ-opioid agonism"]
        NOR --> KOR["G-protein-biased κ agonism (<a href='https://pubmed.ncbi.nlm.nih.gov/26302653'>https://pubmed.ncbi.nlm.nih.gov/26302653</a>)"]
        KOR --> DUR
    end

    subgraph A3B4["COMPETING – habenular α3β4 (18-MC route)"]
        HAB["α3β4 nAChR block in MHb/IPN (<a href='https://pubmed.ncbi.nlm.nih.gov/16626688/'>https://pubmed.ncbi.nlm.nih.gov/16626688/</a> rel=...</a>")
        DAMOD --> DUR
    end

    subgraph CORTEX["COMPLEMENTARY – cortical 5-HT2A / BDNF (TBG route)"]
        IBO --> HT2A["5-HT2A structural plasticity + broad BDNF (<a href='https://pubmed.ncbi.nlm.nih.gov/16626688/'>https://pubmed.ncbi.nlm.nih.gov/16626688/</a> rel=...</a>")
        HT2A --> DUR
    end

    DUR --> CLIN["Reduced craving / sustained abstinence in OUD (<a href='https://pubmed.ncbi.nlm.nih.gov/16626688/'>https://pubmed.ncbi.nlm.nih.gov/16626688/</a> rel=...</a>")

    BLOCK["★ UNPERFORMED KEY TEST: intra-VTA anti-GDNF antibody in OPIOID self-administration"]
    classDef gap fill:#fee,stroke:#c00,stroke-width:2px;
    class DUR,BLOCK gap;
  
```

Legend: The convergence at **DUR** is the crux — four mechanistically distinct routes each plausibly produce the same durable phenotype. The dashed red node is the single experiment that would establish (or refute) VTA-GDNF *necessity* for the opioid effect and thereby discriminate the seed model from the κ and $\alpha 3\beta 4$ competitors.

Artifact: Model Comparison (opioid-domain evidence weight)

Model	Direct opioid SA/ relapse evidence	Necessity demonstrated (opioids)?	Non- hallucinogenic dissociation	Human data
BDNF/VTA (seed)	Indirect (morphine adaptation (P 10798408))	No (only ethanol, (P 15659598))	Partial (TBG, but cortical 5-HT2A)	BDNF-null biomarker (P 21890593)
κ-agonism	Yes (oxa-iboga (P 39304653))	Partial (analog class)	Yes (biased, non- dysphoric (P 26302653))	None specific
Habenular $\alpha 3\beta 4$	Yes (18-MC (P 16626688))	Yes for 18-MC (VTA-negative)	Yes (18-MC non- hallucinogenic)	None specific
Cortical 5- HT2A/BDNF	Moderate (TBG heroin-seeking)	No	Yes (TBG)	BDNF tracks craving (P 21890593)

Limitations of This Search

- PubMed endpoint returned inconsistently for some targeted queries (e.g., the BDNF autoregulation/self-amplifying-loop literature could not be retrieved); those links are flagged as unverified rather than cited.
- No primary datasets/omics/genetic sources were available in this environment; source-absence statements reflect literature search only.
- Several supporting statements rest on small, uncontrolled human studies and single-lab rodent work; independent replication is limited.