

Developmental and Epileptic Encephalopathy 19 (DEE19) — Comprehensive Disease Report

Disease: Developmental and Epileptic Encephalopathy 19 (DEE19) **MONDO:** MONDO:0014328 | **OMIM:** 615744 | **Causal gene:** *GABRA1* (HGNC:4075) **Report type:** Disease-level aggregation from primary literature and ontology/database knowledge (not individual EHR-derived).

Evidence-source key: **[H]** human clinical, **[V]** in vitro/heterologous expression, **[M]** model organism, **[C]** computational. Frequencies for rare, ultra-rare disorders are qualitative unless a cohort number is given. Where a claim reflects general DEE/GABAA biology rather than a DEE19-specific study, this is stated explicitly.

1. Disease Information

Overview. DEE19 is a rare, genetically determined developmental and epileptic encephalopathy caused by heterozygous, usually *de novo*, pathogenic variants in *GABRA1*, the gene encoding the $\alpha 1$ subunit of the type A γ -aminobutyric acid (GABA-A) receptor. It presents in infancy with multiple, frequently drug-resistant seizure types accompanied by developmental delay/intellectual disability. The term "developmental **and** epileptic encephalopathy" denotes that impairment arises both from the underlying genetic lesion acting on brain development (the developmental encephalopathy) **and** from the epileptic activity itself worsening function (the epileptic encephalopathy).

Key identifiers. - **OMIM:** 615744 (Developmental and epileptic encephalopathy 19; formerly "Epileptic encephalopathy, early infantile, 19 / EIEE19") - **MONDO:** MONDO:0014328 - **Gene OMIM:** *GABRA1* 137160 - **Orphanet:** within "Non-syndromic genetic developmental and epileptic encephalopathy" (ORPHA:442835) and the *GABRA1*-related epilepsy spectrum; DEE has no single unique ORPHA number for the 19 subtype. - **ICD-10:** G40.4 (Other generalized epilepsy and epileptic syndromes) / G40.83 (in some coding sets). **ICD-11:** 8A61 / 8A6Z (Developmental and epileptic encephalopathies). - **MeSH:** "Spasms, Infantile" / "Epileptic

Syndromes" / "Epilepsy, Generalized" (no unique DEE19 MeSH; indexed under GABRA1 and epileptic encephalopathy). - **HGNC:** 4075; **NCBI Gene:** 2554; **Ensembl:** ENSG00000022355; **UniProt:** P14867 (GBRA1_HUMAN).

Synonyms / alternative names. Early infantile epileptic encephalopathy 19 (EIEE19); GABRA1-related epileptic encephalopathy; GABRA1-related developmental and epileptic encephalopathy; GABRA1 epilepsy. The *GABRA1* allelic spectrum also includes juvenile myoclonic epilepsy (JME, OMIM 611136) and childhood absence epilepsy susceptibility.

Data provenance. Content here is aggregated at the disease level from primary case series, functional studies, mechanistic reviews, and curated databases (OMIM, ClinVar, HPO, Orphanet); it is not derived from a single patient EHR.

2. Etiology

Primary cause — genetic (monogenic, dominant). DEE19 is caused by pathogenic variants in *GABRA1*. The first *GABRA1* epilepsy variant (p.Ala322Asp, A322D) was identified in a large French-Canadian JME family (Cossette et al. 2002 [H/V],

P 11992121

"an Ala322Asp mutation in GABRA1, encoding the alpha1 subunit of the gamma-aminobutyric acid receptor subtype A (GABA(A)), is found in affected individuals of a large French Canadian family with juvenile myoclonic epilepsy." Severe encephalopathic phenotypes arise chiefly from *de novo* variants (Hernandez et al. 2019 [H/V],

P 31056671

"two known GABRA1 mutations (c.335G>A, p.R112Q and c.343A>G, p.N115D) in six patients with intractable early onset epileptic encephalopathy."

Genetic risk factors. The causal variant itself is the risk factor. Most DEE19 variants are *de novo* dominant (see §9). No independent susceptibility loci or modifier genes are established for DEE19 specifically; general genetic-background modifiers are plausible but unproven.

Environmental risk factors. None established as causal. DEE19 is a monogenic disorder; environment does not initiate disease. General seizure-provoking factors (fever, sleep deprivation, missed medication, intercurrent illness) can precipitate seizures in an already-affected individual but are triggers, not causes.

Protective factors. No genetic protective alleles are defined. The only "protective" influence is functional: variants retaining greater residual GABA-A receptor function tend to produce milder phenotypes (Boßelmann et al. 2026 [H/C],

P 42546502

GABRA3 paralog paradigm, Johannesen et al. 2026 [H/M],

P 41289009

Effective early seizure control is the main modifiable protective factor against epileptic-encephalopathy worsening.

Gene–environment interactions. Not a major feature; disease is genetically determined. GxE is limited to environmental seizure triggers acting on the genetic substrate.

3. Phenotypes

DEE19 is clinically variable; the following are the characteristic features. HPO terms are suggested; frequencies are qualitative/cohort-based given rarity.

Core neurological phenotypes - Seizures, multiple types — the defining feature (obligate, ~100%). Types include generalized tonic-clonic, myoclonic, focal, atonic, absence, tonic, and epileptic spasms. HP:0001250 (Seizure), HP:0002133 (Status epilepticus), HP:0002123 (Generalized myoclonic seizure), HP:0002069 (Generalized tonic-clonic seizure), HP:0011153 (Focal-onset seizure), HP:0011097 (Epileptic spasm), HP:0002121 (Absence seizure). - **Global developmental delay / intellectual disability** (very frequent, ~most patients; severity mild→profound). HP:0001263 (Global developmental delay), HP:0001249 (Intellectual disability). - **Developmental regression / stagnation with seizure onset** (encephalopathy). HP:0002376 (Developmental regression). - **Hypotonia** (frequent). HP:0001252 (Hypotonia). - **Ataxia / movement abnormalities / dystonia** (subset). HP:0001251 (Ataxia), HP:0001332 (Dystonia). - **Abnormal EEG** — multifocal/generalized epileptiform discharges; hypsarrhythmia in those with spasms; sometimes photosensitivity. HP:0002353 (EEG abnormality), HP:0002521 (Hypsarrhythmia). - **Speech/language impairment**, absent or limited speech. HP:0002463 (Language impairment), HP:0001344 (Absent speech). - **Behavioral features** — autistic features, hyperactivity/ADHD-like behavior in a subset. HP:0000729 (Autistic behavior), HP:0000752 (Hyperactivity). - **Microcephaly** (subset, acquired or congenital). HP:0000252 (Microcephaly).

Onset: infantile, typically first months of life; in an 8-patient GABRA1 cohort onset was **3–8 months** (Zhang & Liu 2022 **[H]**);

P 35937053

"Epilepsy onset age was between 3 and 8 months of age." Milder allelic phenotypes (JME) present in adolescence.

Severity/progression: variable severity (mild epilepsy → severe DEE); seizure course is often chronic and drug-resistant with an epileptic-encephalopathy plateau; developmental trajectory is impaired and largely static-to-slowly-progressive rather than neurodegenerative.

Quality-of-life impact: severe. Refractory seizures, intellectual disability, communication and motor impairment produce high dependency, need for caregiving, and reduced QoL for patient and family. No DEE19-specific EQ-5D/SF-36 data exist; QoL burden is inferred from severe DEE literature.

4. Genetic / Molecular Information

Causal gene. *GABRA1* (HGNC:4075; NCBI Gene 2554; OMIM 137160; UniProt P14867), on chromosome **5q34**, encoding the GABA-A receptor $\alpha 1$ subunit (a Cys-loop ligand-gated ion channel subunit with a large extracellular N-terminal GABA-binding domain and four transmembrane helices M1–M4; M2 lines the chloride pore).

Pathogenic variants. - Variant type/class: predominantly heterozygous **missense** variants; also nonsense/frameshift (haploinsufficiency) and splice variants. Reported pathogenic missense residues cluster in (i) the **N-terminal GABA-binding domain** (e.g., p.Arg112Gln/R112Q, p.Asn115Asp/N115D) and (ii) **transmembrane domains** (e.g., p.Pro260Ser/P260S, p.Leu296Ser/L296S, p.Trp315Leu/W315L, p.Ala322Asp/A322D) (Hernandez et al. 2019 **[H/V]**);

P 31056671

Krampf et al. 2005 **[V]**;

P 16029191

"The $\alpha 1$ (R112Q and N115R) subunit residue substitutions were in the N-terminal GABA binding domain." - **Classification (ACMG/AMP):** most recurrent DEE19 variants are Pathogenic/Likely pathogenic in ClinVar; PS2 (*de novo*), PM2 (absent from gnomAD), PS3 (functional studies), PP2 (missense-intolerant gene) commonly apply. VUS exist and benefit from functional/

computational LoF-vs-GoF classification (Boßelmann et al. 2026 [H/C],).

P 42546502

- **Allele frequency:** pathogenic DEE19 variants are absent/ultra-rare in population databases (gnomAD); *GABRA1* is highly constrained (high missense/LoF intolerance). - **Somatic vs germline:** germline (constitutional). *De novo* germline events dominate severe cases; parental germline mosaicism is possible (see §9). No somatic-mosaicism disease role established. - **Functional consequence:** predominantly **loss-of-function** (reduced surface expression, impaired biogenesis, reduced GABA sensitivity/gating), but a subset — particularly pore-lining M2 variants — are **gain-of-function** (Boßelmann 2026,

P 42546502

paralog GABRA3,

P 41289009

Dominant-negative effects on assembled receptors are described for some variants.

Modifier genes. None validated for DEE19. Genetic background likely contributes to variable expressivity.

Epigenetic information. No DEE19-specific DNA-methylation "episignature" is established. (Some other DEE genes have episignatures; *GABRA1* is not a recognized episignature disorder to date.)

Chromosomal abnormalities. DEE19 is typically a single-nucleotide/small-indel disorder. Larger 5q34 deletions encompassing *GABRA1* (contiguous *GABRA1/GABRB2/GABRG2* cluster) can occur and are detectable by chromosomal microarray, but classic DEE19 is not a copy-number syndrome.

5. Environmental Information

- **Environmental factors:** none causal. Not applicable as an etiology.
- **Lifestyle factors:** not applicable to disease causation. Standard seizure hygiene (sleep, adherence) modulates seizure frequency.
- **Infectious agents:** none. DEE19 is genetic, not infectious. (In differential diagnosis, acquired/infectious encephalopathies must be excluded — see §10.)

6. Mechanism / Pathophysiology

Ordered causal chain (initiating lesion → clinical manifestation)

1. A heterozygous (usually *de novo*) **pathogenic variant in *GABRA1*** alters the $\alpha 1$ subunit protein sequence. (*demonstrated*)
2. This **leads to** abnormal $\alpha 1$ subunit biogenesis: impaired folding, endoplasmic-reticulum retention, and **excessive ER-associated degradation (ERAD)** of the misfolded subunit at the expense of forward trafficking, reducing oligomerization/trafficking and **cell-surface expression** of $\alpha 1$ -containing GABA-A receptors (Krampfl 2005 [V],

P 16029191

Macdonald 2010 [V],

P 20308251

Hernandez 2016 [V],

P 27622563

for the archetype A322D, Fu 2018 [V],

P 30481215

"the A322D mutation in the $\alpha 1$ subunit... causes its extensive misfolding and expedited degradation in the endoplasmic reticulum (ER)". (*demonstrated in vitro*) — This ERAD/proteostasis defect is pharmacologically reversible (see §12), implicating GO:0036503 (ERAD pathway) and GO:0030968 (ER unfolded protein response).

3. In parallel, surface-expressed mutant receptors show **reduced GABA-binding affinity and impaired channel gating** (reduced current amplitude, altered activation/deactivation) (Krampfl 2005; Cossette 2002). Branch: rare M2/pore variants instead **increase** channel activity → **gain-of-function** (Boßelmann 2026 [H/C],

P 42546502

(*demonstrated*)

4. Reduced/altered $\alpha 1\beta 2\gamma 2$ receptor function **results in** decreased fast phasic GABAergic inhibitory postsynaptic currents (reduced mIPSC amplitude/frequency) at inhibitory synapses (mouse VB thalamus, Zhou 2015 [M],

P 25447232

(demonstrated in model; inferred in human)

5. Loss of inhibition **leads to** a cortical/thalamocortical **excitation–inhibition imbalance and neuronal hyperexcitability**, with disinhibition of thalamic relay nuclei promoting abnormal thalamocortical oscillations (Zhou 2015 [M]). *(demonstrated in model)*
6. Hyperexcitability **results in** recurrent, multifocal/generalized **seizures** (clinical). *(demonstrated)*
7. Because $\alpha 1$ -GABA-A signaling also shapes **early neurodevelopment** (during the developmental period GABA is depolarizing/excitatory and guides proliferation, migration, differentiation, synaptogenesis; Stojanovic 2016 [H],

P 26518133

receptor dysfunction during a **critical infantile window** — compounded by the deleterious effect of ongoing seizures on the developing network — **leads to developmental delay/intellectual disability and encephalopathy** (the "developmental" + "epileptic" encephalopathy). *(partly inferred)*

8. For gain-of-function variants, excess/altered GABAergic signaling similarly **results in** severe, treatment-resistant epilepsy and profound impairment, by analogy to the GABRA3 paralog paradigm (Johannesen 2026 [H/M],

P 41289009

(inferred by paralogy)

Detail by category

- **Molecular pathways:** GABAergic inhibitory neurotransmission via the ionotropic GABA-A receptor–chloride channel; no canonical growth-factor signaling cascade. GO:0007214 (GABA receptor signaling pathway); GO:1902476 (chloride transmembrane transport); Reactome "GABA A receptor activation."
- **Cellular processes:** loss of fast synaptic inhibition; E/I imbalance; during development, altered neuronal proliferation/migration/synapse maturation (GO:0060078 regulation of postsynaptic membrane potential; GO:0007268 chemical synaptic transmission).
- **Protein dysfunction:** misfolding, ER retention, impaired assembly/trafficking → reduced surface receptor; and/or altered ligand binding and gating. Loss-of-function predominant; gain-of-function and dominant-negative subsets.

- **Metabolic changes:** none primary. (Ketogenic diet may help empirically via network effects, not a defined metabolic defect.)
- **Immune involvement:** none; not autoimmune/inflammatory.
- **Tissue-damage mechanisms:** not degenerative; injury is functional (network dysfunction) plus potential secondary excitotoxic effects of prolonged seizures/status epilepticus.
- **Biochemical abnormality:** ligand-gated **ion channel (chloride) dysfunction** — a channelopathy of inhibitory neurotransmission.
- **Epigenetic changes:** none established.
- **Molecular profiling:** no disease-specific transcriptomic/proteomic/metabolomic signature; mechanistic data derive from heterologous electrophysiology, cryo-EM structure ($\alpha 1\beta 3\gamma 2$; Laverty 2019 [V],

 30602789

and mouse models.

Cell types (CL): GABAergic inhibitory interneuron (CL:0000617), neuron (CL:0000540), pyramidal/glutamatergic neuron (postsynaptic target; CL:0000598), thalamic relay neuron. **Subcellular (GO CC):** postsynaptic membrane GO:0045211; GABA-A receptor complex GO:1902711; integral component of plasma membrane GO:0005887; endoplasmic reticulum GO:0005783 (mislocalization site). **CHEBI:** GABA (CHEBI:16865), chloride (CHEBI:17996).

7. Anatomical Structures Affected

- **Organ / system:** the **brain** and central nervous system (nervous system, UBERON:0001016). Primary organ: brain (UBERON:0000955); cerebral cortex (UBERON:0000956), thalamus (UBERON:0001897, thalamocortical circuits implicated by mouse data), hippocampus (UBERON:0002421), cerebellum ($\alpha 1$ highly expressed; UBERON:0002037). No primary involvement of non-neural organs; systemic effects are secondary to disability (e.g., feeding/respiratory complications).
- **Tissue/cell level:** nervous tissue; GABAergic inhibitory synapses on cortical and thalamic neurons. Affected cell populations: inhibitory interneurons and their postsynaptic partners (CL:0000617 GABAergic neuron; CL:0000540 neuron).

- **Subcellular level:** the **inhibitory postsynaptic membrane** (GO:0045211) and the **GABA-A receptor–chloride channel complex** (GO:1902711); with variant protein mislocalized to the **endoplasmic reticulum** (GO:0005783).
- **Localization / lateralization:** **bilateral, diffuse** cortical involvement (generalized encephalopathy); EEG discharges may be generalized or multifocal. Not a focal/lateralized lesional disorder, though asymmetric epileptiform features can occur.

8. Temporal Development

- **Onset:** congenital genetic lesion with clinical onset typically in **infancy (first year, commonly 3–8 months)** for DEE19; milder allelic phenotypes (JME) present in adolescence. Onset pattern: subacute/insidious emergence of seizures with developmental slowing.
- **Progression:** chronic, lifelong. Seizures are frequently drug-resistant; course is often an early "stormy" phase followed by a plateau (epileptic-encephalopathy pattern). Not classically neurodegenerative; developmental impairment is largely static-to-slowly-evolving.
- **Course pattern:** chronic with episodic seizure clusters and possible status epilepticus; some patients improve in seizure control over time while cognitive impairment persists.
- **Remission:** true remission is uncommon in severe DEE19; seizure freedom is treatment-dependent and variable. Milder allelic phenotypes can be well-controlled.
- **Critical period:** the **infantile window** of synaptic maturation is both the period of maximal vulnerability and the key window for early, effective seizure control and developmental intervention (rationale from developmental GABA biology,

P 26518133

9. Inheritance and Population

- **Epidemiology:** DEE19 is an **ultra-rare** disorder; no precise prevalence/incidence figures are established (individual GABAA-DEE subtypes each account for a small fraction of DEEs). DEEs collectively affect on the order of ~1 in 2,000 children. *GABRA1* is a recognized but minority cause among genetic DEEs.

- **Inheritance:** **autosomal dominant**, most often **de novo** in severe DEE19; **inherited autosomal-dominant** transmission occurs for milder alleles (e.g., the A322D JME family, Cossette 2002,

P 11992121

- **Penetrance:** high for severe *de novo* variants; **incomplete/variable penetrance and expressivity** for some inherited alleles (same variant can yield JME in one relative and more severe epilepsy in another).
- **Expressivity: highly variable** — a hallmark of *GABRA1* (mild IGE ↔ severe DEE), partly explained by variant functional class (LoF vs GoF;

P 42546502

- **Genetic anticipation:** not applicable (not a repeat-expansion disorder).
- **Germline mosaicism:** possible; recurrence in siblings of unaffected parents has been reported for DEE genes generally, warranting counseling.
- **Founder effects:** the A322D JME variant is notable in a French-Canadian pedigree but is family-specific, not a population founder allele.
- **Consanguinity:** not relevant (dominant mechanism).
- **Carrier frequency:** not applicable (dominant, typically *de novo*; pathogenic alleles absent from population databases).
- **Population demographics:** no ethnic predilection; reported worldwide. **Sex ratio ~1:1** (autosomal; the 8-patient cohort was 4 male/4 female,

P 35937053

Age distribution skews to pediatric onset.

10. Diagnostics

- **Genetic testing (definitive): exome or genome sequencing**, or a **multigene epilepsy/DEE panel** including *GABRA1*, is the diagnostic mainstay; **trio** testing establishes *de novo* status. Single-gene *GABRA1* testing is appropriate when the phenotype is highly suggestive. WES/WGS diagnostic yield in early-onset DEE is high (e.g., ~40–72% for onset <3 months; Thanuja & Kamate 2025 [H],

P 40088508

"WES gave an overall yield of 61.9%... and 71.4%... in cases with epilepsy onset before three months."). Early genetic diagnosis frequently changes management (Elkhateeb 2024 [H],

P 38221827

Esterhuizen 2023 [H],

P 36480001

- **Chromosomal microarray (CMA):** to detect 5q34 deletions involving the *GABRA* cluster; usually normal in point-variant DEE19.
- **Karyotype/FISH/mtDNA/repeat-expansion testing:** generally not informative; used to exclude alternative etiologies.
- **Variant interpretation:** ACMG/AMP criteria; functional and computational LoF-vs-GoF classification refines VUS and guides therapy

P 42546502

- **Electrophysiology:** EEG is central — multifocal/generalized epileptiform discharges; hypsarrhythmia if spasms; used for seizure classification and monitoring.
- **Neuroimaging: brain MRI** is typically normal or shows nonspecific findings (e.g., mild atrophy/delayed myelination); primarily excludes structural/malformative causes.
- **Laboratory/biomarkers:** no specific blood/CSF biomarker; metabolic workup (including CSF neurotransmitters, lactate) is used to exclude treatable metabolic epilepsies. The genetic variant is the definitive molecular marker.
- **Clinical criteria:** diagnosis follows the **ILAE** framework for developmental and epileptic encephalopathy (early-onset refractory seizures + developmental impairment) plus a confirmed pathogenic *GABRA1* variant.
- **Differential diagnosis:** other genetic DEEs — *SCN1A* (Dravet), *KCNQ2*, *STXBP1*, *CDKL5*, *SCN2A*, *SCN8A*, *KCNT1*, other GABAA genes (*GABRB3*, *GABRG2*, *GABRB2*, *GABRA5*); structural/metabolic/hypoxic-ischemic encephalopathies; pyridoxine-dependent epilepsy. Distinguished by gene testing and electroclinical pattern.
- **Screening:** not part of newborn screening; **cascade testing** of relatives and prenatal/preimplantation testing available once the familial variant is known.

11. Outcome / Prognosis

- **Survival/mortality:** no DEE19-specific survival figures. Severe DEEs carry elevated mortality including **SUDEP** and status epilepticus (Klein 2025 **[H]**,

([P 40105713](#))

"...refractory to treatment and to have high mortality rates."). Life expectancy is reduced in severely affected, non-ambulant, refractory patients; milder cases have near-normal survival.

- **Morbidity/function:** high — intellectual disability, motor impairment, communication deficits, dependency; frequent comorbid autism/ADHD-like behavior. Long-term disability is the norm in severe cases.
- **Disease course/complications:** drug-resistant epilepsy is common; DEEs show the **highest drug-resistance rate among childhood epilepsy syndromes (77.7%)** (Ayoub et al. 2024 **[H]**,

([P 39150742](#))

Complications: status epilepticus, injury from seizures, aspiration, feeding difficulty, SUDEP.

- **Recovery potential:** limited for cognitive deficits; seizure control may improve with tailored therapy but developmental impairment usually persists.
- **Prognostic factors:** earlier onset, multiple seizure types, myoclonus, and greater developmental delay predict drug resistance/poorer outcome

([P 39150742](#)).

Variant functional class (GoF associated with more severe, treatment-resistant disease) is an emerging molecular prognostic marker

([P 42546502](#)),

([P 41289009](#)).

12. Treatment

General principle: symptomatic seizure control plus developmental support; increasingly **genotype-guided** based on LoF vs GoF variant classification.

- **Pharmacotherapy (antiseizure medications, NCIT: Antiepileptic Agent):** broad-spectrum ASMs are used empirically — **valproate** (NCIT:C935), **levetiracetam**, **clobazam** and other **benzodiazepines** (NCIT:C29075; GABA-A positive modulators), **topiramate**, **lamotrigine**, **zonisamide**, and for spasms **vigabatrin**/ACTH/corticosteroids. **Cenobamate** is a newer ASM with efficacy in refractory epilepsy/DEE (Klein 2025 [H],

P 40105713

Because most DEE19 variants are LoF, drugs that **enhance residual GABA-A function** (benzodiazepines) are mechanistically rational; conversely, GABA-enhancing drugs may be counterproductive for GoF variants — underscoring functional classification.

- **Precision/personalized medicine: vinpocetine** is an emerging targeted therapy for **loss-of-function** GABAA/*GABRA1* DEE — adjunctive vinpocetine reduced seizure frequency and improved comorbidities in a case series of 9 patients with 8 GABAA-receptor variants (Gjerulfsen et al. 2026 [H/V],

P 42227896

"Adjunctive vinpocetine shows promise as a targeted therapy for patients with GABA[A receptor variants]." LoF-vs-GoF prediction models directly inform drug selection (Boßelmann 2026 [H/C],

P 42546502

- **Non-pharmacologic: ketogenic diet** (NCIT:C62735) is used empirically for refractory DEE; **vagus nerve stimulation**; epilepsy surgery is generally not applicable (non-lesional, generalized).
- **Advanced / experimental targeted therapeutics:** For loss-of-function variants that impair receptor **biogenesis/trafficking** (e.g., A322D), **proteostasis-modulating** strategies rescue mutant surface expression and chloride current in cell models — the BiP/HSP70 activator **BIX** and modest activation of the **ATF6/IRE1 unfolded-protein-response** pathways (Fu 2018 [V],

P 30481215

and **VCP/p97 inhibition** (Eeyarestatin I) plus the folding enhancer SAHA/vorinostat

(Han 2015 [V],

P 25406314

Ongoing work maps the GABAA proteostasis network to find drug-correctable variants

(

P 36030824

,

P 40112516

,

P 40161784

).

P 41178115

These are **preclinical/experimental**, not yet clinical. No approved gene therapy, ASO, or cell therapy for DEE19; conceptually attractive for a dominant channelopathy and under preclinical consideration.

- **Supportive/rehabilitative:** physical, occupational, and speech therapy; developmental/educational support; nutritional and respiratory care; management of behavioral comorbidities.
- **Treatment outcomes/adverse events:** response is variable and often partial; polytherapy is common; standard ASM adverse-event profiles apply (sedation, hepatotoxicity/valproate, behavioral effects/levetiracetam, visual-field effects/vigabatrin).
- **Strategy:** individualized, guided by seizure type, EEG, and increasingly variant functional class; combination therapy is frequent in refractory disease.

13. Prevention

- **Primary prevention:** not possible for *de novo* genetic disease. **Genetic counseling** for families with a known variant; **prenatal diagnosis / preimplantation genetic testing (PGT)** can prevent recurrence when a familial pathogenic variant is identified.
- **Secondary prevention:** early genetic diagnosis enables **early, targeted seizure control** to mitigate epileptic-encephalopathy worsening (rationale: developmental critical period,

P 26518133

value of early diagnosis,

P 38221827

- **Tertiary prevention:** optimize seizure control to reduce status epilepticus and SUDEP risk; early developmental therapies; treat comorbidities.
- **Immunization / infectious control:** not applicable (non-infectious); routine vaccination is encouraged as intercurrent illness/fever can trigger seizures.
- **Counseling:** recurrence risk generally low for *de novo* cases but with a caveat for **germline mosaicism**; 50% transmission risk for an affected parent carrying a dominant allele.

14. Other Species / Natural Disease

- **Taxonomy / orthologs:** *GABRA1* is highly conserved across vertebrates. Mouse *Gabra1* (NCBI Gene 14394; **Mus musculus**, NCBITaxon:10090); rat *Gabra1* (NCBITaxon:10116); zebrafish *gabra1* (**Danio rerio**, NCBITaxon:7955). Strong evolutionary conservation of the GABA-A receptor and its inhibitory function underlies cross-species modeling.
- **Natural disease in other species:** no well-characterized naturally occurring *GABRA1* epileptic-encephalopathy analog documented in companion animals (OMIA has no established *GABRA1* DEE entry); GABAergic mechanisms are broadly implicated in animal seizure disorders. Veterinary relevance is chiefly as experimental models rather than spontaneous disease.
- **Comparative biology:** the conserved role of $\alpha 1$ -GABA-A receptors in fast inhibition means mechanistic findings translate across mammals; evolutionary conservation supports use of mouse/zebrafish models.
- **Transmission:** not applicable (non-communicable genetic disease; no zoonotic potential).

15. Model Organisms

- **Mouse (*Mus musculus*, NCBITaxon:10090) — primary model. Heterozygous *Gabra1* deletion produces absence seizures and reduced thalamic inhibition (Zhou et al. 2015 [M],):**

P 25447232

"heterozygous deletion of *Gabra1*, the mouse homolog of the human absence epilepsy gene that encodes the GABAA receptor (*GABAAR*) $\alpha 1$ subunit, causes absence seizures." Demonstrates reduced synaptic $\alpha 1$, reduced mIPSC amplitude/frequency, ventrobasal-

thalamus disinhibition, and partial cortical compensation ($\alpha 3$ upregulation). Knock-in of specific human variants and paralog GoF knock-ins (e.g., *Gabra3*^{Q242L/+} showing increased seizure susceptibility, early death, cortical hyperexcitability;

P 41289009

[M]) model genotype-specific mechanisms.

- **Model types available:** knockout, heterozygous null, and knock-in (point-variant "humanized") mice; conditional alleles feasible via MGI/IMPC resources.
- **In vitro / cellular models:** heterologous expression (HEK293, *Xenopus* oocytes) of recombinant $\alpha 1\beta 2/\beta 3\gamma 2$ receptors for electrophysiology and trafficking assays (Cossette 2002; Krampfl 2005; Hernandez 2016) — the workhorse for LoF/GoF functional classification; patient-derived iPSC neurons are an emerging platform.
- **Phenotype recapitulation:** mouse *Gabra1* models reproduce **seizures and reduced inhibition** well; they capture the core channelopathy and network disinhibition.
- **Limitations:** heterozygous-null mice model absence-type seizures better than the full severe human DEE19 encephalopathy; compensatory subunit upregulation and species differences in subunit expression limit full phenotype capture; cognitive/developmental phenotypes are only partially modeled.
- **Resources:** MGI (mouse), ZFIN (zebrafish), RGD (rat), IMPC/IMSR for strains; Alliance of Genome Resources for orthology.

Summary of Findings, Supported and Refuted Hypotheses

Answer to the research question. DEE19 (OMIM 615744; MONDO:0014328) is a rare autosomal-dominant, usually *de novo*, developmental and epileptic encephalopathy caused by pathogenic variants in *GABRA1* (GABA-A receptor $\alpha 1$ subunit). Variants impair fast GABAergic inhibition — predominantly by loss of function (defective receptor biogenesis, surface expression, GABA sensitivity, and gating), with a gain-of-function subset — producing cortical/thalamocortical excitation–inhibition imbalance that manifests as infantile-onset, often drug-resistant multi-type seizures with developmental delay/intellectual disability.

Supported hypotheses: (1) *GABRA1* is the causal gene [strong: human genetics + functional data]; (2) loss-of-function/impaired inhibition is the dominant mechanism [strong: in vitro + mouse]; (3) variant functional class (LoF vs GoF) drives severity and treatment direction [emerging, strong]; (4) genotype-guided therapy (e.g., vinpocetine for LoF) is rational and clinically promising [emerging].

Refuted / not-applicable: environmental, infectious, autoimmune, metabolic, and epigenetic-etiological etiologies are **not** primary causes; DEE19 is a monogenic channelopathy.

Limitations & future directions. Precise epidemiology, natural-history/QoL metrics, and controlled treatment-outcome data specific to DEE19 are lacking; the pubmed corpus queried here was limited, so some claims lean on GABAA-DEE-wide and paralog evidence. Priorities: variant-specific functional maps (LoF vs GoF), prospective genotype-stratified trials (vinpocetine and GABA-A enhancers for LoF; caution/antagonism strategies for GoF), **clinical translation of proteostasis-corrector approaches** for biogenesis-defective variants (BiP/UPR modulators, VCP inhibition + HDAC inhibitors), iPSC/knock-in disease modeling, and development of allele-selective gene-based therapies (ASO/gene replacement) for this dominant channelopathy.

Key references (PMID)

- 11992121 — Cossette 2002, first *GABRA1* (A322D) epilepsy variant.
- 16029191 — Krampfl 2005, A322D severe loss-of-function mechanisms.
- 20308251 — Macdonald 2010, GABAA subunit epilepsy mechanisms review.
- 27622563 — Hernandez 2016, GABR variant functional risk.
- 31056671 — Hernandez 2019, *de novo* *GABRA1* in early-onset epileptic encephalopathy.
- 35937053 — Zhang & Liu 2022, *GABRA1* pediatric cohort (onset 3–8 mo).
- 25447232 — Zhou 2015, *Gabra1* mouse absence-seizure model.
- 26518133 — Stojanovic 2016, developmental $\alpha 1$ expression / excitatory GABA.
- 30602789 — Lavery 2019, cryo-EM $\alpha 1\beta 3\gamma 2$ structure.
- 42546502 — Boßelmann 2026, LoF/GoF prediction (505 individuals).
- 42227896 — Gjerulfsen 2026, vinpocetine precision therapy for LoF GABAA DEE.
- 41289009 — Johannesen 2026, *GABRA3* paralog LoF/GoF paradigm.
- 30481215 — Fu 2018, ERAD/proteostasis rescue (BIX, ATF6/IRE1) of misfolded $\alpha 1$ (A322D).

- 25406314 — Han 2015, VCP inhibition + SAHA restores trafficking-deficient $\alpha 1$ (A322D).
- 36030824 / 40112516 / 40161784 / 41178115 — GABAA proteostasis network and correctors (experimental).
- 39150742 — Ayoub 2024, drug-resistance in DEE (77.7%).
- 40105713 — Klein 2025, refractory DEE, high mortality, cenobamate.
- 40088508 / 38221827 / 36480001 — WES yield & value of early genetic diagnosis in DEE.

Generated by OpenScientist — Scientific Hypothesis Agent for Novel Discovery