

# **JB Special Review—Fundamental Roles of Glycans in Eukaryotes**

## **The cytoplasmic peptide:N-glycanase (Ngly1)—basic science encounters a human genetic disorder**

Received September 5, 2014; accepted September 30, 2014; published online November 13, 2014

### **Tadashi Suzuki\***

Glycometabolome Team, Systems Glycobiology Research Group, RIKEN-Max Planck Joint Research Center for Systems Chemical Biology, RIKEN Global Research Cluster, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan

\*Tadashi Suzuki, Glycometabolome Team, Systems Glycobiology Research Group, RIKEN-Max Planck Joint Research Center, RIKEN Global Research Cluster, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan. Tel: +81-48-467-9614, Fax: +81-48-462-4692, email: tsuzuki\_gm@riken.jp

**Peptide:N-glycanase (PNGase) is a de-N-glycosylating enzyme that cleaves intact N-glycans from glycoproteins/glycopeptides. The activity of the cytoplasmic PNGase in several mammalian-derived cultured cells was first reported in 1993, and 7 years later, the gene encoding the enzyme was identified in budding yeast. Although the gene—*PNG1* in budding yeast and *NGLY1*/*Ngly1* in mammalian cells—appears to be well conserved throughout eukaryotes, the biological significance of this enzyme has remained elusive until recently. However, discovery of a human genetic disorder involving the *NGLY1* gene clearly indicates that this enzyme plays a critical role in human biology. This review summarizes the research history of cytoplasmic PNGase. The importance of curiosity-driven, pure ‘basic science’ will also be discussed.**

**Keywords:** Ngly1/PNGase/ERAD/genetic disorder/basic science.

**Abbreviations:** ENGase, endo- $\beta$ -N-acetylglucosaminidase; ER, endoplasmic reticulum; ERAD, ER-associated degradation; fOS, free oligosaccharide; GH, glycoside hydrolase; PAW, PNGases and other worm; PNGase, peptide:N-glycanase; PUB, PNGase and ubiquitin; RTA $\Delta$ , ricin toxin A chain mutant; RTL, RTA $\Delta$ -transmembrane-Leu2; TGase, transglutaminase.


Progress in glycobiology over the past few decades has been explosive, and more and more researchers now appreciate the functional importance of glycan chains on glycoproteins. For example, the glycosylation of proteins has been shown to play critical roles in modulating the physiological/physicochemical properties of cognate proteins (1, 2).

Peptide:N-glycanase (PNGase; peptide-N<sup>4</sup>-(N-acetyl- $\beta$ -D-glucosaminyl) asparagine amidase; also referred to as glycoamidase or N-glycanase; EC 3.5.1.52) catalyzes the hydrolysis of the amide bond

between the proximal N-acetylglucosamine (GlcNAc) residue and the asparagine residue of a protein, generating an aspartic acid-containing polypeptide chain and a 1-amino GlcNAc-containing oligosaccharide (Fig. 1). Ammonia is then spontaneously released from the 1-amino GlcNAc at physiological pH (<8), giving rise to a free oligosaccharide (fOS) with an N,N'-diacetylchitobiose structure at the reducing end. PNGase activity was first described in almonds by Dr Noriko Takahashi in 1977 (3), followed by the identification of PNGase F from bacteria (4). Since then, those PNGases from bacteria/plant origin have been extensively used for analysing structures/functions of N-glycans on cognate glycoproteins (5, 6). In my opinion, it is not an overstatement to say that PNGase is a critical reagent for the advancement of glycobiology, as this enzyme greatly facilitates structure/function analyses of N-glycans because glycans can be released from glycoproteins under much milder conditions, compared with chemical deglycosylation, which may result in damaging the protein. Interestingly, however, the physiological function of PNGase itself had not been rigorously investigated in any organisms when I started my career as a researcher.

### **The Cytoplasmic PNGase—Early History**

In 1991, I was an undergraduate student in the laboratory of Prof. Yasuo Inoue, then the Department of Biochemistry and Biophysics, Faculty of Science, the University of Tokyo. That was the monumental year for PNGase research as Dr Akira Seko, then a graduate student in Prof. Inoue's laboratory, discovered the first occurrence of PNGase activity of animal origin, which prefers an acidic pH for optimal activity, in Medaka fish embryos (7). My initial project therefore was to search for similar PNGase activity in mammalian cells. Unfortunately, we were not able to identify PNGase activity similar to that found in Medaka embryos, but after a year of struggle, we unexpectedly identified the PNGase activity with totally distinct enzymatic properties in various mammalian-derived cultured cells (8, 9). The optimal activity of the typical mammalian PNGase is at a neutral pH, and a reducing reagent such as dithiothreitol is required for the *in vitro* detection of activity. The most striking (and puzzling) observation was that most of the enzyme activity was recovered from the cytosol fraction (8, 10). Initially, this observation obscured the functional importance of this enzyme, as not many researchers believed that the N-glycoproteins, potential substrates for the enzyme, would be localized in the cytosol. Why would one expect PNGase to be localized in the cytosol, where no substrates are there?



**Fig. 1 Reaction scheme of PNGase.** PNGase catalyzes the cleavage of the amide bond between the proximal GlcNAc residue and the asparagine residue, thereby producing a deglycosylated peptide in which the glycosylated asparagine is converted to aspartic acid and a free glycan with a 1-amino GlcNAc at the reducing end. The glycan is spontaneously hydrolyzed at physiological pH (<8) to generate ammonia and the corresponding free glycan with *N,N*-diacetylchitobiose at the reducing end.

Although we confirmed that the soluble PNGase activities occur ubiquitously in mammalian cells and tissues (8, 10), the functional role for this enzyme was very uncertain. Accordingly, I conducted an extensive literature survey to look for a clue for the potential function of PNGase or other deglycosylation enzymes in various cellular events, and several possibilities arose (11, 12). One of those hypotheses was that this enzyme could be involved in ‘non-lysosomal’ processing or the degradation of proteins (12). This was about the time when the emerging concept of ‘endoplasmic reticulum (ER)-associated degradation’ rocketed to fame.


### ER-Associated Degradation—Cytosolic Connection

The ER functions to synthesize soluble/membrane-spanning proteins that pass through the secretory pathway in eukaryotic cells. Proteins that are translocated into the lumen of the ER are then co- and post-translationally modified in various ways, including *N*-glycosylation, and eventually exit the ER to be delivered for their respective destinations via vesicular transport. On the other hand, it has long been known that this organelle appears to retain a ‘quality control’ or ‘homeostasis’ system, so that especially non-functional or malfunctioning proteins that fail to fold or oligomerize in a correct manner do not follow the similar transport pathway and are instead degraded rapidly. This degradation process was

independent of lysosomal degradation and was initially assumed to occur in the ER (13). However, it was subsequently found that the cytoplasmic proteasome plays a central role in degrading such proteins (14–20). The process, now called ER-associated degradation or ERAD, constitutes one of the protein quality control systems in eukaryotes (21, 22). In this system, misfolded glycoproteins in the luminal side are released into the cytosol, a process referred to as retrotranslocation or dislocation, after which the cytoplasmic PNGase gains an access to the substrates, *N*-glycoproteins, during their proteasomal degradation (Fig. 2). Soon after the cytoplasmic connection associated with the ERAD process was revealed, several glycoproteins were proposed to be deglycosylated, presumably by the cytoplasmic PNGase, during the degradation process (19, 23–32). These results prompted me to identify the gene encoding the cytoplasmic PNGase to gain additional insights into the functional importance of this enzyme.

### Identification of Gene Encoding the Cytoplasmic PNGase

After completing the PhD degree in 1997, I moved to the laboratory of Prof. William J. Lennarz at the State University of New York at Stony Brook as a post-doctoral fellow. I started to work with budding yeast, *Saccharomyces cerevisiae*, and after successful detection of cytoplasmic PNGase activity with similar enzymatic properties (neutral optimal pH and the



**Fig. 2 The involvement of cytoplasmic PNGase in ERAD.** The glycoproteins destined for degradation are translocated from the ER lumen to the cytosol. There, proteins are degraded mainly by proteasomes. PNGase acts on misfolded glycoproteins in the cytosol during the proteasomal degradation. This reaction is believed to enhance the efficiency of proteasomal degradation, while the precise order of the deglycosylation and proteasomal degradation under physiological conditions remains unclear. Note that other protein modification reactions, such as ubiquitination, are omitted in this scheme for the sake of simplification.

requirement of a reducing reagent for an enzyme assay) (33), we finally identified the gene encoding the cytoplasmic PNGase, which is referred to as *PNG1* (34). The process involved in the identification of the gene was very unique; first, we screened over 400 of Hartwell's temperature-sensitive strain collections (35) and isolated strain no. 352 as the one that was defective in PNGase activity. Our initial expectation was that the PNGase might be an essential gene in yeast, and therefore mutants with defective PNGase would exhibit a signature phenotype, if isolated. To our disappointment, however, after several backcrosses with wild-type strains, no detectable phenotypes co-segregated with a defect of PNGase activity. We then made full use of the combination of classical genetics and the most recent genome information of *S. cerevisiae* to identify the causative gene responsible for the loss of enzyme activity. The gene, *PNG1*, was indeed the one encoding the cytoplasmic PNGase (34). The lack of significant phenotypes in *png1* mutant in budding yeast, however, prevented us from clarifying the biological significance of this enzyme. The mouse orthologue, referred to as *Ngly1* (36), was found to be widely expressed in all tissues examined, which is consistent with the results of an *in vitro* PNGase activity assay (10).

### Structural Divergence of the Cytoplasmic PNGase—Evolutional Consequence?

When examined in the DNA database, it was found that the *PNG1* gene orthologue was widely distributed throughout eukaryotes (34, 37). Consistent with this finding, the soluble PNGase activity with a neutral optimal pH has also been widely reported among eukaryotes (Table I). An interesting feature of the cytoplasmic PNGase is that, while the core transglutaminase (TGase) domain critical for the PNGase activity is conserved, additional domains can be found in


**Table I. Reported occurrence of cytoplasmic PNGase activity.**

Source	References
Mammalian cells	(8, 10, 40)
Hen	(28)
Medaka Fish	(41)
<i>C. elegans</i> <sup>a</sup>	(38, 39)
<i>D. discoideum</i>	(42)
<i>S. cerevisiae</i>	(33)
<i>Schizosaccharomyces pombe</i> <sup>a</sup>	(43)
<i>A. thaliana</i> <sup>a</sup>	(44, 45)

<sup>a</sup>PNGase activity was found in the *Pngl*-orthologue protein expressed in a heterologous system.

the orthologues of higher eukaryotes (Fig. 3). These results most likely suggest that this protein acquired those additional domains as an evolutionary consequence. For example, it is interesting to note that in *Caenorhabditis elegans*, the cytoplasmic PNGase is a dual functional enzyme as thioredoxin, an oxidoreductase, and PNGase (38, 39). This domain structure may reflect the evolutionary requirement linking redox potential and PNGase activity, especially as this enzyme has a catalytic Cys residue. The contribution of those domains outside the catalytic core to the function of the cytoplasmic PNGase represents a key issue that remains to be fully elucidated.

Among the additional domains found in the orthologues of higher eukaryotes, the PUB (PNGase- and ubiquitin related) domain was first identified through a bioinformatics analysis, as I noted that this domain was also observed in various ubiquitin-proteasome system-related proteins (46, 47). Although it was initially hypothesized that it may serve as a protein-protein interaction domain (46), experimental evidence supporting the hypothesis is now accumulating (48–51). On the other hand, the C-terminal PAW domain (domain present in PNGases and other



**Fig. 3 Schematic representation of the primary structure of various eukaryotic PNGase orthologues.** Accession numbers for the protein sequences in Universal Protein Resource (<http://www.uniprot.org>) are as follows: *S. cerevisiae*, Q02890; *N. crassa*, D1MY48; *A. thaliana*, Q9FGY9; *D. discoideum*, Q55FC8; *C. elegans*, Q9TW67; *Dr. melanogaster*, Q7KRR5; *Mus musculus*, Q9JI78; *Homo sapiens*, Q96IV0. The domain structure is based on the conserved domain database in National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/cdd>) and common region aligned in all PNGase orthologues are shown: PUB domain (domain ID: cd10459; residue 11–93); TGase-superfamily domain (domain ID: pfam01841; residue 37–107); PAW (mannose binding) domain (domain ID: smart00613; residue 1–89); Thioredoxin-family domain (domain ID: cd02947; residue 2–92); L-type lectin domain (domain ID: cd01951; residue 11–223).

worm proteins) (47) has now been shown to be involved in the oligosaccharide binding of PNGase (52). The cytoplasmic PNGase from mammalian cells was previously found to retain lectin-like properties (53) and it was later suggested that certain types of fOSs can inhibit the enzyme activity in a non-competitive manner (54). The PAW domain, as an additional carbohydrate-binding domain besides the catalytic site, may be involved in this interesting biochemical property found in mammalian cytoplasmic PNGases. With regard to the sugar-binding specificity of the PAW domain, it was reported that the domain preferably binds larger high-mannose glycans, especially with those retaining the intact C-arm and lack the terminal  $\alpha$ 1,2-linked mannose in B-arm (Fig. 4) (55). How these specificities correlate with the properties of the cytoplasmic PNGase remains to be investigated.


Besides the mannose-binding PAW domain, the core TGase domain has an intrinsic ability to bind to *N*, *N*-diacetylchitobiose-containing glycans (54, 56, 57). Haloacetamidyl derivatives of oligosaccharides containing an *N*, *N*-diacetylchitobiose structure are potent inhibitors as well as a labelling reagent of the cytoplasmic PNGase, as they specifically label the catalytic Cys residue (56, 58). Those probes were shown to bind to other PNGase orthologues (59, 60), indicating that *N*, *N*-diacetylchitobiose-binding properties are a common feature of the cytoplasmic PNGase. The *N*, *N*-diacetylchitobiose-modified yeast PNGase has

been structurally characterized, and the residues responsible for carbohydrate binding have been identified (57). It was also shown that Z-VAD-fmk, a general caspase inhibitor, is a potent inhibitor of cytoplasmic PNGase (61, 62). Although Z-VAD-fmk may be used as an inhibitor of cytoplasmic PNGase, caution may be taken in its use, since it has a broad inhibition profile.

### Protein–Protein Interaction Involving the Cytoplasmic Peptide:*N*-glycanase

It should be noted that the cytoplasmic PNGase can participate in various protein–protein interactions with other proteins. In 2001, Rad23 was identified as a PNGase-binding protein through yeast two-hybrid screening (63). Although Rad23 was first identified as a protein involved in nucleotide excision repair (64), evidence shows that this protein binds to the 26S proteasome through the N-terminal ubiquitin-like domain (65) and also plays a role in proteasomal degradation (66, 67). Although the interaction between Rad23 and the PNGase appears to be conserved, the complex interphase is fundamentally different between yeast and mouse proteins, as revealed by the structure analysis of the PNGase–Rad23 complex (68, 69). It is also worth mentioning that Rad23 forms a complex with another DNA repair-related protein, Rad4, which has a similar core ‘TGase fold’ (70). It is therefore tempting to speculate that PNGase–Rad23 and





**Fig. 4** Structures of *N*-glycans bound with relatively higher affinity ( $K_a > 4.0 \times 10^4$ ) to the PAW domain of mouse Ngly1, as determined by frontal affinity chromatography analysis (55). The definition of the arms (A/B/C arm) described in the text is also indicated.

Rad4–Rad23 may have evolved from a common ancestor protein complex.

In addition to Rad23, various PNGase-binding proteins have been isolated for mouse Ngly1 protein (71). It is interesting to note that, among the proteins identified, most bind through the N-terminal region of Ngly1, which contains the PUB domain. One should also note that, upon subcellular fractionation, a minor fraction of the PNGase activity was associated with the membrane-bound fraction (8, 10, 33). Interestingly, the cytoplasmic PNGase in mammals is able to bind to p97/VCP/Cdc48, a key ATPases associated with diverse cellular activities (AAA) adenosine triphosphate (ATPase) for the ERAD pathway, as well as other ERAD or ubiquitin–proteasome pathway-related proteins, some of which are intrinsic membrane proteins (71–75). Those protein–protein interactions may also be mainly mediated by the PUB domain. Given the minor distribution of Ngly1 in the membrane fraction, however, binding to membrane proteins should be a sub-stoichiometric interaction. How such minor interactions contribute to the critical function of Ngly1 will need to be clarified via further studies. Recent findings also suggest that, at least *in vitro*, Fbs1, a glycoprotein-specific ubiquitin ligase (76, 77), protects misfolded glycoproteins from the action of cytoplasmic PNGase (78). These protein–protein interactions may be critical for establishing complex protein networks and for regulating the sequential reactions in a concerted manner, as the formation of such so-called ‘glycoprotein degradation complexes’ may greatly enhance the efficiency of ERAD by carrying out a number of reactions, *i.e.* ubiquitination, deglycosylation, deubiquitination and proteolysis in one location (79, 80).

### PNGase-Dependent and -Independent Pathway for Non-Lysosomal Degradation of Free Oligosaccharides

In yeast cells, the absence of cytoplasmic PNGase (Png1) results in significant reduction of the levels of

fOSs found in the cytosol, suggesting that the majority, if not all, of the fOSs in yeast are generated from misfolded glycoproteins in a PNGase-dependent manner (81–83). Therefore, these results indicate that analyses of the structure/amount of fOSs could serve as a read-out of overall glycan processing during the ERAD process (82, 83). Interestingly, this does not appear to be the case in mammalian cells, as the knockdown or knockout of the *Ngly1* gene has only a negligible effect on the structure/amount of fOSs (84) (Harada, Y. and Suzuki, T., unpublished data). These results clearly suggest that, in mammalian cells, the fOSs are derived from the cytoplasmic PNGase-independent reactions, the majority of which are, most likely, due to the hydrolytic activity of oligosaccharyltransferase (85). Therefore, one should evaluate the origin of fOSs carefully when analysed in different organisms (86). There are also cytosolic glycosidases involved in the catabolism of fOSs in mammals, such as endo- $\beta$ -*N*-acetylglucosaminidase (ENGase) (87, 88) or the cytoplasmic  $\alpha$ -mannosidase, Man2C1 (89, 90). For more details on the non-lysosomal process of glycan catabolism, readers are directed to review articles (80, 86, 91, 92).

### Functional Role of the Cytoplasmic PNGase in Various Organisms

Since the gene encoding the cytoplasmic PNGase was identified, much progress has been made in clarifying its role in the ERAD process. In budding yeast, a ricin toxin A chain mutant (RTA $\Delta$ ) was identified as an ERAD substrate that is degraded in a Png1-dependent manner (67, 93). Moreover, we recently established an *in vivo* system designed to evaluate the Png1-dependent ERAD using a new substrate, RTL (RTA $\Delta$ -transmembrane-Leu2), which consists of a luminal RTA $\Delta$  and the Leu2 cytoplasmic protein that is involved in the biosynthesis of leucine connected with a transmembrane domain (44, 94). RTL can be used to detect strains with defective ERAD by assessing the growth of yeast strains bearing a *leu2* mutation, as cells with

Table II. Reported phenotypes/pathological conditions caused by mutations in gene orthologues of cytoplasmic PNGase.

Source	Phenotypes observed	References
<i>S. cerevisiae</i>	Defect in ERAD; no growth/viability defects	(34, 67, 94)
<i>N. crassa</i>	Temperature-sensitive growth with strong polarity defects	(60, 105, 106)
<i>A. thaliana</i>	No obvious defect	(45)
<i>D. discoideum</i>	Slow growth and development; defect in cell aggregation during multicellular development	(42, 107)
<i>C. elegans</i>	Abnormal axon branching of VC4/VC5 egg-laying neurons, egg-laying behaviour defect	(108)
<i>Dr. melanogaster</i>	Severe developmental delay	(59)
<i>H. sapiens</i>	Global developmental delay, movement disorder, hypotonia	(109–111)

defective RTL degradation, when RTL is expressed, can supply sufficient Leu2 (as a part of RTL protein) to survive in a media lacking leucine. It should be noted that the deglycosylation of CPY\*, the best-studied glycoprotein ERAD substrate in yeast, was not observed in *png1Δ* cells, while a delay in degradation of CPY\* was observed (34). Quite recently, however, we observed the PNGase-mediated deglycosylation of intact CPY\* as well as its proteolytic degradation intermediates when proteasomal activity was compromised (Hosomi, A. and Suzuki, T., unpublished data). This result clearly indicates that Png1 is indeed involved in the degradation of CPY\*. It was also suggested that degradation of 6Myc-Hmg2, an ERAD substrate with a lesion in the transmembrane domain (ERAD-M), was attenuated in *png1Δ* cells (95), while the presence of deglycosylated 6Myc-Hmg2 was not detected in the presence of Png1.

In sharp contrast to the case of yeast, the functional importance of the cytoplasmic PNGase in mammalian ERAD process remains relatively unknown, other than the proposed critical role for class I major histocompatibility complex antigen presentation (29, 96–100). Initial attempts to inhibit PNGase activity by RNA interference or the use of inhibitors resulted in little or no effect on the efficiency of degradation of model substrates (61, 101, 102). However, recently quite a reliable assay system was developed for detecting *in vivo* PNGase activity, using a green fluorescent protein (GFP)-derived model ERAD substrate that exhibits fluorescence in a deglycosylation-dependent manner (103). This assay promises to be a very facile system for assessing *in vivo* PNGase activity in mammalian cells. More recently, it was also reported that EDEM1 protein, a key component of the glycoprotein ERAD process, was stabilized on the inhibition of cytoplasmic PNGase, implying that the EDEM1 may be an endogenous substrate for cytoplasmic PNGase (104). This result may also shed light on the functional importance of the cytoplasmic PNGase in mammalian cells. It should be noted that the inhibition of PNGase activity appeared to have little effect on the deglycosylation of EDEM1. Further studies will be needed to clarify the functional importance of PNGase on the stability of the EDEM1 protein.

Although the mutant of the gene orthologue for the cytoplasmic PNGase in budding yeast (34) or in *Arabidopsis thaliana* (45) did not show any significant phenotypic consequences, more recent studies suggest

that it may play important physiological roles in other eukaryotes (Table II).

### Deglycosylation-Dependent and -Independent Function of the Cytoplasmic PNGase: Which Is Important?

It was shown that mutants of PNG1 in *Neurospora crassa* showed temperature-sensitive growth with strong polarity defects (105) (Table II). Interestingly, the PNG1 in *N. crassa* contains intrinsic mutations in two of the three essential amino acids in the catalytic triad (*i.e.* Cys-to-Ala and His-to-Tyr mutations; (60)) that consists of Cys, His and Asp (Fig. 5) (112, 113). It is therefore predicted that, at least in some organisms, PNGase orthologues will also have a deglycosylation-independent function. It is also interesting to note here that the phenotypes observed in *N. crassa* (hyphal growth defect) and *C. elegans* (abnormal axon branching) are both temperature sensitive (105, 108). As hyphal growth and other cell migration processes in animal cells such as axon branching may share, at least in part, a common machinery, it will be important to see whether the phenotype observed in *C. elegans* may somehow be related to the deglycosylation-independent function of the PNGase orthologue found in *N. crassa*.

It should be noted here that cytoplasmic PNGase possesses two well-conserved CXXC motifs (Fig. 5). Initial mutagenesis study revealed that a mutation of the Cys residues in these CXXC motifs results in the loss of enzyme activity, suggesting that those motifs are critical for enzyme activity (113). As one of the CXXC motifs in *Drosophila melanogaster* PNGase orthologue (Pngl) is missing, and initial attempts to detect PNGase activity in the fly failed, it was proposed that the fly Pngl protein may also have an important enzyme-independent function (59). However, we now have evidence to show that Pngl also appears to have deglycosylating activity (Huang, C. and Suzuki, T., unpublished data), and those CXXC motifs are also missing in the *Dictyostelium discoideum* orthologue (Fig. 5), while having deglycosylating activity (42). These results imply that the CXXC motif may not be an absolute requirement for enzyme activity.

Quite recently, it was reported that, in filamentous fungi where no active cytoplasmic PNGase has been found, fungi-specific cytoplasmic ENGase appears to

	1	11	21	31	41	51	61
Consensus	llleLLhWfk	eefFtWvNnp	pCs--kCggp	t----srd-s	glPtdeElkh	g-akrVEvyr	CdaCgtetRF
Conservation	:*: :*	:* :*	* *		*	**	*. :*
M.musculus	LLLELLHWFk	EEFFRWVNNI	VCS--KCGGE	T---RSRDEA	LLPNDDDELKW	G-AKNVENHY	CDACQLSNRF
H.sapiens	LLLELLHWFk	EEFFHWVNNV	LCS--KCGGQ	T---RSRDRS	LLPSDDDELKW	G-AKEVEDHY	CDACQFSNRF
A.thaliana	FLQLLLWFk	KS-FRWVNEP	PCD--FCGNK	T---IGQG-M	GNPLTSELAY	G-ANRVEIYR	CTMCPTTTRF
C.elegans	ILKDLLHWFk	TQFFTWFDPR	TCF--KCTLK	C---STDGLQ	GTPTREEQKE	GGASRVEVYR	CDGCNTEMRF
D.discoideum	KMLMLLDWFk	NEYFTWTNSP	ECSDIKCGTP	S---TSSVGS	DRPTFEEQSH	Q-VSIVEVYR	C-ASNHVTRF
S.cerevisiae	LVKELLRYFK	QDFFKWCNKP	DCN--HCGQN	TSENMTPLGS	QGPNGEESKF	N-CGTVEIYK	CNRCGNITRF
D.melanogaster	LLVELLVNWFN	TQFFQWVNNI	PCR--VCGSE	E-----	SRLRRTEREG	D--IRVEVTV	C--CGQESKPF
N.crassa	MVRALLRWFR	RSFFTfVNNP	PCS--ECLSP	T---NKIRN	VAPTPEERAH	S-ATWVELYA	CVTCGAYERF
	71	81	91	101	111	121	131
Consensus	PRYNdpekLL	etrcGRCGEW	ANcFTllcra	lgfearyvwd	ftDHVWtEvY	sesqqRWlhv	DpCEdvcDkP
Conservation	**.. :*	:**.*:	* * * :	* * *	*: * * *	* * *	*. * *
M.musculus	PRYNNPEKLL	ETRCGRCGEW	ANCFTLCCRA	LGFEARYVWD	YTDHVWTEVY	SPSQQRWLHC	DACEDEVCDKP
H.sapiens	PRYNNPEKLL	ETRCGRCGEW	ANCFTLCCRA	VGFEARYVWD	YTDHVWTEVY	SPSQQRWLHC	DACEDEVCDKP
A.thaliana	PRYNDPLKLV	ETKKGRCGEW	ANCFPLYCRT	FGYDSRLIMD	FTDHVWTECY	SHSLKRWIHL	DPCEGVYDKP
C.elegans	PRYNNPAKLL	QTRTGRCGEW	ANCFGLLLAA	LNLESRFIYD	TTDHVWNEVY	LLAEQRWCHV	DPCEINTMDRP
D.discoideum	PRYNSVEKLL	STKCGRCGEW	ANAFTLFSIA	LGFTTRYILD	FTDHVWNEVY	IDG--RWIHW	DSCEATYDSP
S.cerevisiae	PRYNDPIKLL	ETRKGRCEW	CNLFRTLILKS	FGLDVRYVWN	REDHVWCEYF	SNFLNRWVHV	DSCEQSFDQP
D.melanogaster	YRYNDISQLL	VSRKGRCGEY	ANCFTLFCRA	LDYDARIVHS	HFDHVWTEVY	SEAQMRLWHV	DPSENVIDSP
N.crassa	PRYTEAWQLL	RVKRGRAADF	ANVFMTLCRA	LDIRARVWVC	QEDYLWTEIY	SEHQQRWVHV	DSCEAWDMF
	141	151	161				
Consensus	llYeigWgKk	lsYvIAfskd	gvvDVTWRY				
Conservation	*. * :	:. * :.	*: * *				
M.musculus	LLYEIGWGKK	LSYIIAFSKD	EVVDVTWRY				
H.sapiens	LLYEIGWGKK	LSYVIAFSKD	EVVDVTWRY				
A.thaliana	MLYEKGWNKK	LNIVIAISKD	GVCDVTKRY				
C.elegans	LLYTRGWGKT	LGVCIGYGSD	HVVDVTWRY				
D.discoideum	LYTEGGWGKQ	LSYVFAFEFN	GIYDVTSTRY				
S.cerevisiae	YIYSINWNKK	MSYCIAFGKD	GVVDVSKRY				
D.melanogaster	LMYQHGWKRH	IDYILAYSRD	DIQDVTWRY				
N.crassa	HMYKKNWGKK	MSYVIAFSRE	GAVDVTTRY				

**Fig. 5 Comparison of amino acid sequences around the catalytic residues between the cytoplasmic PNGase orthologues.** The sequences of the TGase domain (Fig. 3) and its surrounding regions well conserved throughout the PNGase orthologues were aligned. The amino acid residues aligned are as follows; *M. musculus* (Q9JI78), 226–388; *H. sapiens* (Q96IV0), 229–391; *A. thaliana* (Q9FGY9), 173–333; *C. elegans* (Q9TW67), 170–333; *D. discoideum* (Q55FC8), 129–290; *S. cerevisiae* (Q02890), 108–273; *Dr. melanogaster* (Q7KRR5), 225–378; *N. crassa* (D1MY48), 129–290. The two CXXC motifs conserved in some orthologues are underlined. The catalytic residues (Cys, His and Asp) are indicated by number signs (#). Sequence alignment was carried out using ClustalW (114) and the alignment result is shown using UCSF Chimera software (115) (courtesy of Mr Masaki Kato, Structural Glycobiology Team, RIKEN). Note that two of the three catalytic residues are not conserved in the PNGase orthologue from *N. crassa*.

be present. This fungi ENGase belongs to the glycoside hydrolase (GH) family 18 in the CAZy database (116), which is different from the GH family 85 cytosolic ENGase ubiquitously found in other eukaryotes (87). It was found that the fungi-specific cytoplasmic ENGase can promote the degradation of RTL in budding yeast *png1Δ* cells (117). It is tempting to speculate that filamentous fungi may have acquired the fungi-specific GH 18 cytosolic ENGase to substitute for the deglycosylation function of the cytoplasmic PNGase.

## Discovery of Human Patients of *NGLY1* Human Genetic Disorder

Recently, an exome analysis identified a human patient with mutations in the *NGLY1* gene (109) and an increasing number of the patients harbouring mutations in *NGLY1* alleles have been reported since then (110, 111). The patients exhibited multiple symptoms that include global developmental delay, multifocal epilepsy, involuntary movement, abnormal liver function and the absence of tears (109–111). The mechanistic insight into these phenotypic consequences remains elusive. Whatever the mechanism, however, these observations clearly indicate the functional importance of the cytoplasmic PNGases in human biology. This rapidly developing *NGLY1* story has enjoyed

exposure from major media such as CNN (<http://edition.cnn.com/2014/03/20/health/ngly1-genetic-disorder/>) or the New Yorker magazine ([http://www.newyorker.com/reporting/2014/07/21/140721fa\\_fact\\_mnookin](http://www.newyorker.com/reporting/2014/07/21/140721fa_fact_mnookin)) because the research was actually not at all led by researchers; it was patients' families who were the driving force (118). I am hopeful that this example will serve as a new paradigm for conducting medical research, as we now know that concerted efforts among patients' families, clinicians and researchers can make a huge difference.

## Beauty of 'Basic Science'

As I stated above, many unexpected, serendipitous findings have arisen during my 20-plus year scientific journey dealing with research on cytoplasmic PNGase (its discovery was already a completely unexpected result). Who would have thought that human genetic disorders could exist because of mutations in *NGLY1* gene, when no obvious phenotype is seen in mutants of the yeast orthologous gene? These unexpected findings continue as we analyse *ngly1*-KO mice, on which we hope to report in the near future. The bottom line is that my research on PNGase/Ngly1 is all about curiosity-driven science. If one can only be allowed to carry out 'important science', we would not have been able to continue our studies of this enzyme.



In the beginning, I could not have cared less about how 'important' this enzyme will be. I just wanted to find out what this enzyme is doing, irrespective of whether it is important or not.

My scientific journey, which represents quite a turnaround from pure basic science to identify human genetic diseases, reminds me of the old words by Prof. Yasuo Inoue, my very first mentor; He always emphasized that both basic research and applied research are important; while everyone understands the importance of the latter, the former is sometimes not so much appreciated. Whichever the direction will be, however, there was one important motto on research (in the original words of Prof. Fujio Egami, Prof. Inoue's mentor):

'There is no "important" research subject.

If there is, it must have already been made important by others.

What the most important is to make our own research be important by ourselves!'

After all this time, I came to realize that I simply followed my mentor's (or my mentor's mentor's) words. In this connection, it has been just 20 years since we first wrote reviews describing the potential functions of deglycosylating enzymes in nature (11, 12), some of which still remain as attractive hypotheses. I just feel fortunate that, in our research field, there are so many outstanding questions that still remain to be explored.

## Acknowledgements

I am deeply grateful to the following former mentors/advisors, who generously allowed me to pursue my genuine interests: the late Prof. Yasuo Inoue, Prof. Sadako Inoue, Prof. Ken Kitajima (Nagoya University), Prof. Yasufumi Emori (University of Tokyo), Prof. William J. Lennarz (Stony Brook University), the late Prof. Yoshitaka Nagai and Prof. Naoyuki Taniguchi (RIKEN). I also wish to thank former/current collaborators and laboratory members for sharing my excitement and enthusiasm. Special thanks go to Masaki Kato (Structural Glycobiology Team, RIKEN) for his kind help on preparing Fig. 5.

## Funding

This research in the Suzuki laboratory is supported by the Yamada Science Foundation as well as Grant-In-Aids (25291030, 25650041, 26110725) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

## Conflict of Interest

None declared.

## References

- Varki, A. (1993) Biological roles of oligosaccharides: all of the theories are correct. *Glycobiology* **3**, 97–130
- Haltiwanger, R.S. and Lowe, J.B. (2004) Role of glycosylation in development. *Annu. Rev. Biochem.* **73**, 491–537
- Takahashi, N. (1977) Demonstration of a new amidase acting on glycopeptides. *Biochem. Biophys. Res. Commun.* **76**, 1194–1201
- Plummer, T.H. Jr., Elder, J.H., Alexander, S., Phelan, A.W., and Tarentino, A.L. (1984) Demonstration of peptide:N-glycosidase F activity in endo-beta-N-acetylglucosaminidase F preparations. *J. Biol. Chem.* **259**, 10700–10704
- Maley, F., Trimble, R.B., Tarentino, A.L., and Plummer, T.H. Jr. (1989) Characterization of glycoproteins and their associated oligosaccharides through the use of endoglycosidases. *Anal. Biochem.* **180**, 195–204
- Wang, T. and Voglmeir, J. (2014) PNGases as valuable tools in glycoprotein analysis. *Protein Pept. Lett.* **21**, 976–985
- Seko, A., Kitajima, K., Inoue, Y., and Inoue, S. (1991) Peptide:N-glycosidase activity found in the early embryos of *Oryzias latipes* (Medaka fish). The first demonstration of the occurrence of peptide:N-glycosidase in animal cells and its implication for the presence of a de-N-glycosylation system in living organisms. *J. Biol. Chem.* **266**, 22110–22114
- Suzuki, T., Seko, A., Kitajima, K., Inoue, Y., and Inoue, S. (1993) Identification of peptide:N-glycanase activity in mammalian-derived cultured cells. *Biochem. Biophys. Res. Commun.* **194**, 1124–1130
- Suzuki, T., Seko, A., Kitajima, K., Inoue, Y., and Inoue, S. (1994) Purification and enzymatic properties of peptide:N-glycanase from C3H mouse-derived L-929 fibroblast cells. Possible widespread occurrence of post-translational remodeling of proteins by N-deglycosylation. *J. Biol. Chem.* **269**, 17611–17618
- Kitajima, K., Suzuki, T., Kouchi, Z., Inoue, S., and Inoue, Y. (1995) Identification and distribution of peptide:N-glycanase (PNGase) in mouse organs. *Arch. Biochem. Biophys.* **319**, 393–401
- Suzuki, T., Kitajima, K., Inoue, S., and Inoue, Y. (1994) Occurrence and biological roles of proximal glycanases in animal cells. *Glycobiology* **4**, 777–789
- Suzuki, T., Kitajima, K., Inoue, S., and Inoue, Y. (1995) N-glycosylation/deglycosylation as a mechanism for the post-translational modification/remodification of proteins. *Glycoconj. J.* **12**, 183–193
- Klausner, R.D. and Sitia, R. (1990) Protein degradation in the endoplasmic reticulum. *Cell* **62**, 611–614
- Ward, C.L., Omura, S., and Kopito, R.R. (1995) Degradation of CFTR by the ubiquitin-proteasome pathway. *Cell* **83**, 121–127
- Biederer, T., Volkwein, C., and Sommer, T. (1996) Degradation of subunits of the Sec61p complex, an integral component of the ER membrane, by the ubiquitin-proteasome pathway. *EMBO J.* **15**, 2069–2076
- Hiller, M.M., Finger, A., Schweiger, M., and Wolf, D.H. (1996) ER degradation of a misfolded luminal protein by the cytosolic ubiquitin-proteasome pathway. *Science* **273**, 1725–1728
- Qu, D., Teckman, J.H., Omura, S., and Perlmutter, D.H. (1996) Degradation of a mutant secretory protein, alpha1-antitrypsin Z, in the endoplasmic reticulum requires proteasome activity. *J. Biol. Chem.* **271**, 22791–22795
- Werner, E.D., Brodsky, J.L., and McCracken, A.A. (1996) Proteasome-dependent endoplasmic reticulum-associated protein degradation: an unconventional route to a familiar fate. *Proc. Natl. Acad. Sci. U S A.* **93**, 13797–13801
- Wiertz, E.J., Jones, T.R., Sun, L., Bogoy, M., Geuze, H.J., and Ploegh, H.L. (1996) The human cytomegalovirus US11 gene product dislocates MHC class I heavy chains from the endoplasmic reticulum to the cytosol. *Cell* **84**, 769–779



20. Kopito, R.R. (1997) ER quality control: the cytoplasmic connection. *Cell* **88**, 427–430
21. Needham, P.G. and Brodsky, J.L. (2013) How early studies on secreted and membrane protein quality control gave rise to the ER associated degradation (ERAD) pathway: the early history of ERAD. *Biochim. Biophys. Acta* **1833**, 2447–2457
22. Merulla, J., Fasana, E., Solda, T., and Molinari, M. (2013) Specificity and regulation of the endoplasmic reticulum-associated degradation machinery. *Traffic* **14**, 767–777
23. Wiertz, E.J., Tortorella, D., Bogoy, M., Yu, J., Mothes, W., Jones, T.R., Rapoport, T.A., and Ploegh, H.L. (1996) Sec61-mediated transfer of a membrane protein from the endoplasmic reticulum to the proteasome for destruction. *Nature* **384**, 432–438
24. Hughes, E.A., Hammond, C., and Cresswell, P. (1997) Misfolded major histocompatibility complex class I heavy chains are translocated into the cytoplasm and degraded by the proteasome. *Proc. Natl. Acad. Sci. U S A* **94**, 1896–1901
25. Huppa, J.B. and Ploegh, H.L. (1997) The alpha chain of the T cell antigen receptor is degraded in the cytosol. *Immunity* **7**, 113–122
26. Yu, H., Kaung, G., Kobayashi, S., and Kopito, R.R. (1997) Cytosolic degradation of T-cell receptor alpha chains by the proteasome. *J. Biol. Chem.* **272**, 20800–20804
27. Halaban, R., Cheng, E., Zhang, Y., Moellmann, G., Hanlon, D., Michalak, M., Setaluri, V., and Hebert, D.N. (1997) Aberrant retention of tyrosinase in the endoplasmic reticulum mediates accelerated degradation of the enzyme and contributes to the dedifferentiated phenotype of amelanotic melanoma cells. *Proc. Natl. Acad. Sci. U S A* **94**, 6210–6215
28. Suzuki, T., Kitajima, K., Emori, Y., Inoue, Y., and Inoue, S. (1997) Site-specific de-N-glycosylation of diglycosylated ovalbumin in hen oviduct by endogenous peptide:N-glycanase as a quality control system for newly synthesized proteins. *Proc. Natl. Acad. Sci. U S A* **94**, 6244–6249
29. Mosse, C.A., Meadows, L., Luckey, C.J., Kittlesen, D.J., Huczko, E.L., Slingluff, C.L., Shabanowitz, J., Hunt, D.F., and Engelhard, V.H. (1998) The class I antigen-processing pathway for the membrane protein tyrosinase involves translation in the endoplasmic reticulum and processing in the cytosol. *J. Exp. Med.* **187**, 37–48
30. Bebok, Z., Mazzochi, C., King, S.A., Hong, J.S., and Sorscher, E.J. (1998) The mechanism underlying cystic fibrosis transmembrane conductance regulator transport from the endoplasmic reticulum to the proteasome includes Sec61beta and a cytosolic, deglycosylated intermediary. *J. Biol. Chem.* **273**, 29873–29878
31. de Virgilio, M., Weninger, H., and Ivessa, N.E. (1998) Ubiquitination is required for the retro-translocation of a short-lived luminal endoplasmic reticulum glycoprotein to the cytosol for degradation by the proteasome. *J. Biol. Chem.* **273**, 9734–9743
32. Johnston, J.A., Ward, C.L., and Kopito, R.R. (1998) Aggresomes: a cellular response to misfolded proteins. *J. Cell. Biol.* **143**, 1883–1898
33. Suzuki, T., Park, H., Kitajima, K., and Lennarz, W.J. (1998) Peptides glycosylated in the endoplasmic reticulum of yeast are subsequently deglycosylated by a soluble peptide:N-glycanase activity. *J. Biol. Chem.* **273**, 21526–21530
34. Suzuki, T., Park, H., Hollingsworth, N.M., Sternglanz, R., and Lennarz, W.J. (2000) *PNG1*, a yeast gene encoding a highly conserved peptide:N-glycanase. *J. Cell. Biol.* **149**, 1039–1052
35. Hartwell, L.H. (1967) Macromolecule synthesis in temperature-sensitive mutants of yeast. *J. Bacteriol.* **93**, 1662–1670
36. Suzuki, T., Kwofie, M.A., and Lennarz, W.J. (2003) *Ngly1*, a mouse gene encoding a deglycosylating enzyme implicated in proteasomal degradation: expression, genomic organization, and chromosomal mapping. *Biochem. Biophys. Res. Commun.* **304**, 326–332
37. Suzuki, T., Park, H., and Lennarz, W.J. (2002) Cytoplasmic peptide:N-glycanase (PNGase) in eukaryotic cells: occurrence, primary structure, and potential functions. *FASEB J.* **16**, 635–641
38. Suzuki, T., Tanabe, K., Hara, I., Taniguchi, N., and Colavita, A. (2007) Dual enzymatic properties of the cytoplasmic peptide:N-glycanase in *C. elegans*. *Biochem. Biophys. Res. Commun.* **358**, 837–841
39. Kato, T., Kawahara, A., Ashida, H., and Yamamoto, K. (2007) Unique peptide:N-glycanase of *Caenorhabditis elegans* has activity of protein disulphide reductase as well as of deglycosylation. *J. Biochem.* **142**, 175–181
40. Romisch, K. and Ali, B.R. (1997) Similar processes mediate glycopeptide export from the endoplasmic reticulum in mammalian cells and *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. U S A* **94**, 6730–6734
41. Seko, A., Kitajima, K., Iwamatsu, T., Inoue, Y., and Inoue, S. (1999) Identification of two discrete peptide:N-glycanases in *Oryzias latipes* during embryogenesis. *Glycobiology* **9**, 887–895
42. Gosain, A., Lohia, R., Shrivastava, A., and Saran, S. (2012) Identification and characterization of peptide:N-glycanase from *Dictyostelium discoideum*. *BMC Biochem.* **13**, 9
43. Xin, F., Wang, S., Song, L., Liang, Q., and Qi, Q. (2008) Molecular identification and characterization of peptide:N-glycanase from *Schizosaccharomyces pombe*. *Biochem. Biophys. Res. Commun.* **368**, 907–912
44. Masahara-Negishi, Y., Hosomi, A., Della Mea, M., Serafini-Fracassini, D., and Suzuki, T. (2012) A plant peptide:N-glycanase orthologue facilitates glycoprotein ER-associated degradation in yeast. *Biochim. Biophys. Acta* **1820**, 1457–1462
45. Diepold, A., Li, G., Lennarz, W.J., Nurnberger, T., and Brunner, F. (2007) The *Arabidopsis* AtPNG1 gene encodes a peptide:N-glycanase. *Plant J.* **52**, 94–104
46. Suzuki, T., Park, H., Till, E.A., and Lennarz, W.J. (2001) The PUB domain: a putative protein-protein interaction domain implicated in the ubiquitin-proteasome pathway. *Biochem. Biophys. Res. Commun.* **287**, 1083–1087
47. Doerks, T., Copley, R.R., Schultz, J., Ponting, C.P., and Bork, P. (2002) Systematic identification of novel protein domain families associated with nuclear functions. *Genome Res.* **12**, 47–56
48. Allen, M.D., Buchberger, A., and Bycroft, M. (2006) The PUB domain functions as a p97 binding module in human peptide N-glycanase. *J. Biol. Chem.* **281**, 25502–25508
49. Madsen, L., Seeger, M., Semple, C.A., and Hartmann-Petersen, R. (2009) New ATPase regulators—p97 goes to the PUB. *Int. J. Biochem. Cell Biol.* **41**, 2380–2388
50. Kamiya, Y., Uekusa, Y., Sumiyoshi, A., Sasakawa, H., Hirao, T., Suzuki, T., and Kato, K. (2012) NMR characterization of the interaction between the PUB domain

- of peptide:N-glycanase and ubiquitin-like domain of HR23. *FEBS Lett.* **586**, 1141–1146
51. Schaeffer, V., Akutsu, M., Olma, M.H., Gomes, L.C., Kawasaki, M., and Dikic, I. (2014) Binding of OTULIN to the PUB domain of HOIP controls NF-kappaB signaling. *Mol. Cell.* **54**, 349–361
  52. Zhou, X., Zhao, G., Truglio, J.J., Wang, L., Li, G., Lennarz, W.J., and Schindelin, H. (2006) Structural and biochemical studies of the C-terminal domain of mouse peptide-N-glycanase identify it as a mannose-binding module. *Proc. Natl. Acad. Sci. U S A.* **103**, 17214–17219
  53. Suzuki, T., Kitajima, K., Inoue, S., and Inoue, Y. (1994) Does an animal peptide: N-glycanase have the dual role as an enzyme and a carbohydrate-binding protein? *Glycoconj. J.* **11**, 469–476
  54. Suzuki, T., Kitajima, K., Inoue, Y., and Inoue, S. (1995) Carbohydrate-binding property of peptide: N-glycanase from mouse fibroblast L-929 cells as evaluated by inhibition and binding experiments using various oligosaccharides. *J. Biol. Chem.* **270**, 15181–15186
  55. Kamiya, Y., Kamiya, D., Urade, R., Suzuki, T., and Kato, K. (2009) Sophisticated modes of sugar recognition by intracellular lectins involved in quality control of glycoproteins in *Glycobiology Research Trends* (Powell, G. and McCabe, O., eds.) pp. 27–40, Nova Science Publishers, Hauppauge, NY
  56. Suzuki, T., Hara, I., Nakano, M., Zhao, G., Lennarz, W.J., Schindelin, H., Taniguchi, N., Totani, K., Matsuo, I., and Ito, Y. (2006) Site-specific labeling of cytoplasmic peptide:N-glycanase by *N,N'*-diacetylchitobiose-related compounds. *J. Biol. Chem.* **281**, 22152–22160
  57. Zhao, G., Li, G., Zhou, X., Matsuo, I., Ito, Y., Suzuki, T., Lennarz, W.J., and Schindelin, H. (2009) Structural and mutational studies on the importance of oligosaccharide binding for the activity of yeast PNGase. *Glycobiology* **19**, 118–125
  58. Hagihara, S., Miyazaki, A., Matsuo, I., Tatami, A., Suzuki, T., and Ito, Y. (2007) Fluorescently labeled inhibitor for profiling cytoplasmic peptide:N-glycanase. *Glycobiology* **17**, 1070–1076
  59. Funakoshi, Y., Negishi, Y., Gergen, J.P., Seino, J., Ishii, K., Lennarz, W.J., Matsuo, I., Ito, Y., Taniguchi, N., and Suzuki, T. (2010) Evidence for an essential deglycosylation-independent activity of PNGase in *Drosophila melanogaster*. *PLoS One* **5**, e10545
  60. Mierz, S., Funakoshi, Y., Negishi, Y., Suzuki, T., and Seiler, S. (2010) The *Neurospora* peptide:N-glycanase ortholog PNG1 is essential for cell polarity despite its lack of enzymatic activity. *J. Biol. Chem.* **285**, 2326–2332
  61. Misaghi, S., Pacold, M.E., Blom, D., Ploegh, H.L., and Korbel, G.A. (2004) Using a small molecule inhibitor of peptide: N-glycanase to probe its role in glycoprotein turnover. *Chem. Biol.* **11**, 1677–1687
  62. Misaghi, S., Korbel, G.A., Kessler, B., Spooner, E., and Ploegh, H.L. (2006) z-VAD-fmk inhibits peptide:N-glycanase and may result in ER stress. *Cell Death Differ.* **13**, 163–165
  63. Suzuki, T., Park, H., Kwofie, M.A., and Lennarz, W.J. (2001) Rad23 provides a link between the Png1 deglycosylating enzyme and the 26 S proteasome in yeast. *J. Biol. Chem.* **276**, 21601–21607
  64. Watkins, J.F., Sung, P., Prakash, L., and Prakash, S. (1993) The *Saccharomyces cerevisiae* DNA repair gene RAD23 encodes a nuclear protein containing a ubiquitin-like domain required for biological function. *Mol. Cell. Biol.* **13**, 7757–7765
  65. Schaubert, C., Chen, L., Tongaonkar, P., Vega, I., Lambertson, D., Potts, W., and Madura, K. (1998) Rad23 links DNA repair to the ubiquitin/proteasome pathway. *Nature* **391**, 715–718
  66. Heessen, S., Masucci, M.G., and Dantuma, N.P. (2005) The UBA2 domain functions as an intrinsic stabilization signal that protects Rad23 from proteasomal degradation. *Mol. Cell.* **18**, 225–235
  67. Kim, I., Ahn, J., Liu, C., Tanabe, K., Apodaca, J., Suzuki, T., and Rao, H. (2006) The Png1-Rad23 complex regulates glycoprotein turnover. *J. Cell. Biol.* **172**, 211–219
  68. Lee, J.H., Choi, J.M., Lee, C., Yi, K.J., and Cho, Y. (2005) Structure of a peptide:N-glycanase-Rad23 complex: insight into the deglycosylation for denatured glycoproteins. *Proc. Natl. Acad. Sci. U S A.* **102**, 9144–9149
  69. Zhao, G., Zhou, X., Wang, L., Li, G., Kisker, C., Lennarz, W.J., and Schindelin, H. (2006) Structure of the mouse peptide N-glycanase-HR23 complex suggests co-evolution of the endoplasmic reticulum-associated degradation and DNA repair pathways. *J. Biol. Chem.* **281**, 13751–13761
  70. Anantharaman, V., Koonin, E.V., and Aravind, L. (2001) Peptide-N-glycanases and DNA repair proteins, Xp-C/Rad4, are, respectively, active and inactivated enzymes sharing a common transglutaminase fold. *Hum. Mol. Genet.* **10**, 1627–1630
  71. Park, H., Suzuki, T., and Lennarz, W.J. (2001) Identification of proteins that interact with mammalian peptide:N-glycanase and implicate this hydrolase in the proteasome-dependent pathway for protein degradation. *Proc. Natl. Acad. Sci. U S A.* **98**, 11163–11168
  72. Katiyar, S., Li, G., and Lennarz, W.J. (2004) A complex between peptide:N-glycanase and two proteasome-linked proteins suggests a mechanism for the degradation of misfolded glycoproteins. *Proc. Natl. Acad. Sci. U S A.* **101**, 13774–13779
  73. Katiyar, S., Joshi, S., and Lennarz, W.J. (2005) The retrotranslocation protein Derlin-1 binds peptide:N-glycanase to the endoplasmic reticulum. *Mol. Biol. Cell* **16**, 4584–4594
  74. Li, G., Zhou, X., Zhao, G., Schindelin, H., and Lennarz, W.J. (2005) Multiple modes of interaction of the deglycosylation enzyme, mouse peptide N-glycanase, with the proteasome. *Proc. Natl. Acad. Sci. U S A.* **102**, 15809–15814
  75. Li, G., Zhao, G., Zhou, X., Schindelin, H., and Lennarz, W.J. (2006) The AAA ATPase p97 links peptide N-glycanase to the endoplasmic reticulum-associated E3 ligase autocrine motility factor receptor. *Proc. Natl. Acad. Sci. U S A.* **103**, 8348–8353
  76. Yoshida, Y., Chiba, T., Tokunaga, F., Kawasaki, H., Iwai, K., Suzuki, T., Ito, Y., Matsuoka, K., Yoshida, M., Tanaka, K., and Tai, T. (2002) E3 ubiquitin ligase that recognizes sugar chains. *Nature* **418**, 438–442
  77. Yoshida, Y. and Tanaka, K. (2010) Lectin-like ERAD players in ER and cytosol. *Biochim. Biophys. Acta* **1800**, 172–180
  78. Yamaguchi, Y., Hirao, T., Sakata, E., Kamiya, Y., Kurimoto, E., Yoshida, Y., Suzuki, T., Tanaka, K., and Kato, K. (2007) Fbs1 protects the misfolded glycoproteins from the attack of peptide:N-glycanase. *Biochem. Biophys. Res. Commun.* **362**, 712–716
  79. Suzuki, T. and Lennarz, W.J. (2003) Hypothesis: a glycoprotein-degradation complex formed by protein-protein

- interaction involves cytoplasmic peptide:N-glycanase. *Biochem. Biophys. Res. Commun.* **302**, 1–5
80. Suzuki, T. (2007) Cytoplasmic peptide:N-glycanase and catabolic pathway for free N-glycans in the cytosol. *Semin. Cell Dev. Biol.* **18**, 762–769
  81. Chantret, I., Frenoy, J.P., and Moore, S.E. (2003) Free-oligosaccharide control in the yeast *Saccharomyces cerevisiae*: roles for peptide:N-glycanase (Png1p) and vacuolar mannosidase (Ams1p). *Biochem. J.* **373**, 901–908
  82. Hirayama, H., Seino, J., Kitajima, T., Jigami, Y., and Suzuki, T. (2010) Free oligosaccharides to monitor glycoprotein endoplasmic reticulum-associated degradation in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **285**, 12390–12404
  83. Chantret, I., Kodali, V.P., Lahmouich, C., Harvey, D.J., and Moore, S.E. (2011) Endoplasmic reticulum-associated degradation (ERAD) and free oligosaccharide generation in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **286**, 41786–41800
  84. Chantret, I., Fasseu, M., Zaoui, K., Le Bizec, C., Yaye, H.S., Dupre, T., and Moore, S.E. (2010) Identification of roles for peptide: N-glycanase and endo-beta-N-acetylglucosaminidase (Engase1p) during protein N-glycosylation in human HepG2 cells. *PLoS One* **5**, e11734
  85. Harada, Y., Buser, R., Ngwa, E.M., Hirayama, H., Aebi, M., and Suzuki, T. (2013) Eukaryotic oligosaccharyl-transferase generates free oligosaccharides during N-glycosylation. *J. Biol. Chem.* **288**, 32673–32684
  86. Suzuki, T. and Harada, Y. (2014) Non-lysosomal degradation pathway for N-linked glycans and dolichol-linked oligosaccharides. *Biochem. Biophys. Res. Commun.* **453**, 213–219
  87. Suzuki, T., Yano, K., Sugimoto, S., Kitajima, K., Lennarz, W.J., Inoue, S., Inoue, Y., and Emori, Y. (2002) Endo-beta-N-acetylglucosaminidase, an enzyme involved in processing of free oligosaccharides in the cytosol. *Proc. Natl. Acad. Sci. U S A.* **99**, 9691–9696
  88. Kato, T., Fujita, K., Takeuchi, M., Kobayashi, K., Natsuka, S., Ikura, K., Kumagai, H., and Yamamoto, K. (2002) Identification of an endo-beta-N-acetylglucosaminidase gene in *Caenorhabditis elegans* and its expression in *Escherichia coli*. *Glycobiology* **12**, 581–587
  89. Suzuki, T., Hara, I., Nakano, M., Shigeta, M., Nakagawa, T., Kondo, A., Funakoshi, Y., and Taniguchi, N. (2006) Man2C1, an alpha-mannosidase, is involved in the trimming of free oligosaccharides in the cytosol. *Biochem. J.* **400**, 33–41
  90. Costanzi, E., Balducci, C., Cacan, R., Duvet, S., Orlacchio, A., and Beccari, T. (2006) Cloning and expression of mouse cytosolic alpha-mannosidase (Man2c1). *Biochim. Biophys. Acta* **1760**, 1580–1586
  91. Chantret, I. and Moore, S.E. (2008) Free oligosaccharide regulation during mammalian protein N-glycosylation. *Glycobiology* **18**, 210–224
  92. Suzuki, T. (2009) Introduction to “glycometabolome”. *Trends Glycosci. Glycotechnol.* **21**, 219–227
  93. Tanabe, K., Lennarz, W.J., and Suzuki, T. (2006) A cytoplasmic peptide: N-glycanase. *Methods Enzymol.* **415**, 46–55
  94. Hosomi, A., Tanabe, K., Hirayama, H., Kim, I., Rao, H., and Suzuki, T. (2010) Identification of an Htm1 (EDEM)-dependent, Mns1-independent endoplasmic reticulum-associated degradation (ERAD) pathway in *Saccharomyces cerevisiae*: application of a novel assay for glycoprotein ERAD. *J. Biol. Chem.* **285**, 24324–24334
  95. Kario, E., Amar, N., Elazar, Z., and Navon, A. (2011) A new autophagy-related checkpoint in the degradation of an ERAD-M target. *J. Biol. Chem.* **286**, 11479–11491
  96. Skipper, J.C., Hendrickson, R.C., Gulden, P.H., Brichard, V., Van Pel, A., Chen, Y., Shabanowitz, J., Wolfel, T., Slingsluff, C.L. Jr., Boon, T., Hunt, D.F., and Engelhard, V.H. (1996) An HLA-A2-restricted tyrosinase antigen on melanoma cells results from post-translational modification and suggests a novel pathway for processing of membrane proteins. *J. Exp. Med.* **183**, 527–534
  97. Altrich-VanLith, M.L., Ostankovitch, M., Polefrone, J.M., Mosse, C.A., Shabanowitz, J., Hunt, D.F., and Engelhard, V.H. (2006) Processing of a class I-restricted epitope from tyrosinase requires peptide N-glycanase and the cooperative action of endoplasmic reticulum aminopeptidase 1 and cytosolic proteases. *J. Immunol.* **177**, 5440–5450
  98. Kario, E., Tirosh, B., Ploegh, H.L., and Navon, A. (2008) N-linked glycosylation does not impair proteasomal degradation but affects class I major histocompatibility complex presentation. *J. Biol. Chem.* **283**, 244–254
  99. Ostankovitch, M., Altrich-Vanlith, M., Robila, V., and Engelhard, V.H. (2009) N-glycosylation enhances presentation of a MHC class I-restricted epitope from tyrosinase. *J. Immunol.* **182**, 4830–4835
  100. Dalet, A., Robbins, P.F., Stroobant, V., Vigneron, N., Li, Y.F., El-Gamil, M., Hanada, K., Yang, J.C., Rosenberg, S.A., and Van den Eynde, B.J. (2011) An antigenic peptide produced by reverse splicing and double asparagine deamidation. *Proc. Natl. Acad. Sci. U S A.* **108**, E323–E331
  101. Hirsch, C., Blom, D., and Ploegh, H.L. (2003) A role for N-glycanase in the cytosolic turnover of glycoproteins. *EMBO J.* **22**, 1036–1046
  102. Blom, D., Hirsch, C., Stern, P., Tortorella, D., and Ploegh, H.L. (2004) A glycosylated type I membrane protein becomes cytosolic when peptide: N-glycanase is compromised. *EMBO J.* **23**, 650–658
  103. Grotzke, J.E., Lu, Q., and Cresswell, P. (2013) Deglycosylation-dependent fluorescent proteins provide unique tools for the study of ER-associated degradation. *Proc. Natl. Acad. Sci. U S A.* **110**, 3393–3398
  104. Park, S., Jang, I., Zuber, C., Lee, Y., Cho, J.W., Matsuo, I., Ito, Y., and Roth, J. (2014) ERADication of EDEM1 occurs by selective autophagy and requires deglycosylation by cytoplasmic peptide N-glycanase. *Histochem. Cell Biol.* **142**, 153–169
  105. Seiler, S. and Plamann, M. (2003) The genetic basis of cellular morphogenesis in the filamentous fungus *Neurospora crassa*. *Mol. Biol. Cell.* **14**, 4352–4364
  106. Dieterle, M.G., Wiest, A.E., Plamann, M., and McCluskey, K. (2010) Characterization of the temperature-sensitive mutations un-7 and png-1 in *Neurospora crassa*. *PLoS One* **5**, e10703
  107. Gosain, A., Srivastava, A., and Saran, S. (2014) Peptide: N-glycanase is expressed in prestalk cells and plays a role in the differentiation of prespore cells during development of *Dictyostelium discoideum*. *Indian J. Exp. Biol.* **52**, 197–206
  108. Habibi-Babadi, N., Su, A., de Carvalho, C.E., and Colavita, A. (2010) The N-glycanase png-1 acts to limit axon branching during organ formation in *Caenorhabditis elegans*. *J. Neurosci.* **30**, 1766–1776
  109. Need, A.C., Shashi, V., Hitomi, Y., Schoch, K., Shianna, K.V., McDonald, M.T., Meisler, M.H., and



- Goldstein, D.B. (2012) Clinical application of exome sequencing in undiagnosed genetic conditions. *J. Med. Genet.* **49**, 353–361
110. Enns, G.M., Shashi, V., Bainbridge, M., Gambello, M.J., Zahir, F.R., Bast, T., Crimian, R., Schoch, K., Platt, J., Cox, R., Bernstein, J.A., Scavina, M., Walter, R.S., Bibb, A., Jones, M., Hegde, M., Graham, B.H., Need, A.C., Oviedo, A., Schaaf, C.P., Boyle, S., Butte, A.J., Chen, R., Chen, R., Clark, M.J., Haraksingh, R., Cowan, T.M., He, P., Langlois, S., Zoghbi, H.Y., Snyder, M., Gibbs, R.A., Freeze, H.H., Goldstein, D.B., Chen, R., and Consortium, F.C. (2014) Mutations in *NGLY1* cause an inherited disorder of the endoplasmic reticulum-associated degradation pathway. *Genet. Med.* **16**, 751–758
  111. Caglayan, A.O., Comu, S., Baranoski, J.F., Parman, Y., Kaymakcalan, H., Akgumus, G.T., Caglar, C., Dolen, D., Erson Omay, E.Z., Harmanci, A.S., Mishra, K., Freeze, H.H., Yasuno, K., Bilguvar, K., and Gunel, M. (2014) *NGLY1* mutation causes neuromotor impairment, intellectual disability, and neuropathy. *Eur. J. Med. Genet.*, doi: 10.1016/j.ejmg.2014.08.008
  112. Makarova, K.S., Aravind, L., and Koonin, E.V. (1999) A superfamily of archaeal, bacterial, and eukaryotic proteins homologous to animal transglutaminases. *Protein Sci.* **8**, 1714–1719
  113. Katiyar, S., Suzuki, T., Balgobin, B.J., and Lennarz, W.J. (2002) Site-directed mutagenesis study of yeast peptide:*N*-glycanase. Insight into the reaction mechanism of deglycosylation. *J. Biol. Chem.* **277**, 12953–12959
  114. Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H., Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., Thompson, J.D., Gibson, T.J., and Higgins, D.G. (2007) Clustal W and clustal X version 2.0. *Bioinformatics* **23**, 2947–2948
  115. Meng, E.C., Pettersen, E.F., Couch, G.S., Huang, C.C., and Ferrin, T.E. (2006) Tools for integrated sequence-structure analysis with UCSF Chimera. *BMC Bioinformatics*. **7**, 339
  116. Cantarel, B.L., Coutinho, P.M., Rancurel, C., Bernard, T., Lombard, V., and Henrissat, B. (2009) The Carbohydrate-Active EnZymes database (CAZy): an expert resource for glycogenomics. *Nucleic Acids Res.* **37**, D233–D238
  117. Tzelepis, G., Hosomi, A., Hossain, T.J., Hirayama, H., Dubey, M., Jensen, D.F., Suzuki, T., and Karlsson, M. (2014) Endo-beta-*N*-acetylglucosamidases (ENGases) in the fungus *Trichoderma atroviride*: possible involvement of the filamentous fungi-specific cytosolic ENGase in the ERAD process. *Biochem. Biophys. Res. Commun.* **449**, 256–261
  118. Might, M. and Wilsey, M. (2014) The shifting model in clinical diagnostics: how next-generation sequencing and families are altering the way rare diseases are discovered, studied, and treated. *Genet. Med.* **16**, 736–737