PATIENT DEMOGRAPHIC FORM 10/27/2026

YANEZ FREITEZ, ROSANGELA ANA

DOB: 07/09/1986
Sex: Female
DOA: 09/07/2026
Case type: No Fault
Address: 725 10TH ST, New York, NY, 10019

Cell Phone: 347-758-1902
Home phone:
Social Security Number:

Primary Insurance: PROGRESSIVE (NY) - NYP
Address: PO BOX 2930, Clinton, IA, 52733
Claim Number: 26-599343106
Policy Number: 875571455
Policy Holder: YANEZ FREITEZ, ROSANGELA ANAIS
Phone Number:
Fax:
Carrier Case:
WCB Case:

Secondary Insurance:
Address:
Claim Number:
Policy Number:
Policy Holder:
Phone Number:
Fax:

BI Attorney: MORRISON & WAGNER LLP
Address: 49 W 38TH STREET, New York, NY, 10018
Phone number: 212-343-8000
WC Attorney:
Address:
Phone number:

Referring Doctor:

Referring Office:

ATLANTIC MEDICAL & DIAGNOSTIC, P.C.

41-12 77th Street 3rd Floor Elmhurst NY11373 - 1996
Phone#:929 813-1399 , Fax#: 866 300-4168, Email: info@atlanticmdpc.com

Administrative
48 Wall Street, Suite 1100, New York, NY 10005
Phone#: (844) 425-8637; 1212-918-4669, Fax#: (866) 300-4168; 212-918-4801
Email: info@atlanticmdpc.com

PROGRESS NOTE

Patient First Name:	Patient Last Name:	Date of Birth:	Sex:
Rosangela	Yanez Freitez	07-09-1986	Female
Patient Address:		**SSN:**	
725 10th Street, New York NY 10019			
Attending Provider:	**Referring Provider:**	**Visit Date:**	**Chart No.:**
Jerry Samedi, PA-C		09-16-2026	SCL74562
Insurance Name:	**Insurance Subscriber's ID:**	**Date of Injury:**	**Claim Id:**
Progressive Insurance Co-NF	875571455	09-07-2026	NO NUMBER
Appointment Location Address:	**Encounter Type:**		
41-12 77th Street , 3rd Floor, Elmhurst NY 11373 - 1996	New Patient - Auto-Accident		

Chief Complaint: Headache, Cervical Spine Pain, Shoulder Pain, Thoracic Spine Pain, Knee Pain, Lumbar Spine Pain

Motor Vehicle Accident

Rosangela Yanez Freitez is a 40 year old female who was involved in an accident on 09-07-2026. Patient current employment status is Cook.Working Reduced hours due to accident At the time of accident patient was wearing seatbelt. Patient was a back seat passenger. Patient was in the car. Three other people were in the vehicle. she suffered a rear impact to the vehicle. moving she suffered no loss of consciousness.
This was the first car accident of the patient. She was immediately admitted to the ER. Brooklyn Hospital Center In the ER, a physical exam, X-rays and pain killers was done. Cray for Neck, Shoulder and Wrist was then sent home
Patient does not have history of back problems in the past.

Headache

She states that her headache is constant. Location of headache at onset was Frontal and Bilateral. Patient reports that she has same episode(s) 3-6 hours. **She reports that she has photophobia.** There is no vomiting associated with her headache. **There is neck stiffness.** She does not report pain in her eyes.

Cervical Spine Pain
Patient reported pain is bilateral. The patient describes her pain as constant. The pain is aching, pressure like, pulling and tight . Patient says, at its worse her pain is 8/10. Right now she describes her pain as 8/10 on a pain scale of 0-10. The pain is made worse by bending, such as cleaning, moving items, laundry, movement, sitting a long time and turning side to side. Other associated symptoms/problems are as follows: difficulty staying asleep due to pain .

Shoulder Pain
Patient reported pain is on right side. The patient describes her pain as constant. The pain is aching, numbing, pressure like, sharp, pulling and tight . **The pain radiates to the right upper extremity. Numbness** Patient says, at its worse her pain is 9/10. Right now she describes her pain as 9/10 on a pain scale of 0-10. The pain is made worse by bending, such as cleaning, moving items, laundry, lifting, movement and laying on area. Other associated symptoms/problems are as follows: difficulty staying asleep due to pain .

Thoracic Spine Pain
Patient reported pain is bilateral. The patient describes her pain as constant. The pain is aching, pressure like, sharp, pulling and tight . Patient says, at its worse her pain is 8/10. Right now she describes her pain as 8/10 on a pain scale of 0-10. The pain is made worse by bending, such as cleaning, moving items, laundry, lifting, movement and standing a long time. Other associated symptoms/problems are as follows: difficulty staying asleep due to pain .

Knee Pain
Patient reported pain is on left side. The patient describes her pain as constant. The pain is aching, pressure like, sharp, tight and feels weak . Patient says, at its worse her pain is 9/10. Right now she describes her pain as 9/10 on a pain scale of 0-10. The pain is made worse by bending, going up stairs, going down stairs, such as cleaning, moving items, laundry, movement, standing a long time and standing straight up. Other associated symptoms/problems are as follows: difficulty staying asleep due to pain .

Lumbar Spine Pain
Patient reported pain is bilateral. The patient describes her pain as constant. The pain is aching, pressure like, sharp, pulling and tight . Patient says, at its worse her pain is 9/10. Right now she describes her pain as 8/10 on a pain scale of 0-10. The pain is made worse by bending, such as cleaning, moving items, laundry, lifting, sitting a long time, standing a long time and standing straight up. Other associated symptoms/problems are as follows: difficulty staying asleep due to pain .

Past Medical History
No Known Past Medical History

Surgical History
Tummy Tuck: 2026
Breast Augmentation : 2026
NO METALS

Allergy
No Known Drug Allergies.

Social History
Use of Drugs/Alcohol/Tobacco: Patient states that she never drinks any alcohol. Current some day smoker

Review of Systems

Musculoskeletal: Reports **muscle pain**, **muscle cramp**, **muscle pain or tenderness**, **neck pain**, **shoulder pain**, **back pain** and **joint pain**. Denies muscle twitches, muscle wasting, muscle weakness, loss of muscle bulk, joint stiffness, joint swelling, morning stiffness, swollen joints, night cramps, atrophy and postural abnormalities. Denies any joint problem or abnormalities. Denies any limitation of joint movements. No history of fractures.

Neurological: Denies seizures, blackouts, trouble with memory, trouble concentrating, stroke, loss of strength, fainting spells, memory loss, involuntary movements, poor coordination, numbness, spasticity, weakness and tremors. The patient does not observe any gait disturbance. Reports **headache**.

Psychiatric: Denies anxiety, depression and mood swings. Reports **nervousness** and **sleeping difficulty**.

Vitals

Weight: 168.00 lbs. **Handedness:** right-handed. **Height:** 66.00 inches.

Physical Examination

Cervical Spine Exam: Inspection reveals normal curvature of the cervical spine. The cervical spine musculature is supple. Palpation of the cervical facets reveals no pain. Cervical spine is noted to be stable. **Anterior flexion is noted to be 30 degrees (ROM range 0-60 degrees). There is pain noted when neck is flexed anteriorly. Extension of cervical spine noted to be 40 degrees (ROM range 0-75 degrees). There is pain noted with extension of cervical spine. Left lateral rotation noted to be 50 degrees (ROM range 0-80 degrees). There is pain noted with left lateral rotation of C-spine. Left lateral flexion noted to be 20 degrees (ROM range 0-45 degrees). There is pain noted with left lateral flexion of C-spine. Right lateral rotation of the C-spine is noted to be 40 degrees (ROM range 0-80 degrees). There is pain noted with right lateral rotation. Right lateral flexion of the C-spine is noted to be 20 degrees (ROM range 0-45 degrees). There is pain noted with right lateral flexion.** There are no palpable trigger points in the muscles of the head and neck.

Thoracic Spine: There is no evidence of atrophy or asymmetry noted in the thoracic spine. **There is tenderness noted at paraspinal muscles, facet joint lines, levator scapulae muscle and supraspinatus.** Range of motion of the thoracic spine is normal with both flexion and extension without pain. There is no evidence of crepitation, laxity or instability noted. **Hyperextension of thoracic spine caused increased pain.**

Lumbar Spine: Inspection of the lumbar spine reveals no scoliosis. Palpation of the lumbar facet reveals no pain. There is no pain noted over the lumbar intervertebral spaces (discs) on palpation. Palpation of the bilateral sacroiliac joint area reveals tenderness Palpation of the bilateral sacroiliac joint area reveals no pain. Trochanteric Bursa - Tenderness Palpation of the greater trochanteric bursa reveals no tenderness bilaterally. **Anterior flexion of lumbar spine is noted to be Pain at 45 degrees. Anterior lumbar flexion causes pain. Extension of lumbar spine is noted to be 5 degrees (ROM range 0-30 degrees). There is pain noted with lumbar extension. Left lateral flexion of the lumbar spine is noted to be 5 degrees (ROM range 0-25 degrees). Left lateral flexion causes pain. Right lateral flexion of the lumbar spine is noted to be 5 degrees (ROM range 0-25 degrees). There is pain noted with right lateral flexion. Trigger Point Tenderness Assessment: Erector Spinae muscle, Lumbar Multifidus Muscle, Latissimus Dorsi Muscle, Positive Jump Sign, Positive Local Twitch response and Positive Referred Pain.**

Joints-Shoulder: **On palpation, there is tenderness noted over the right anterior acromioclavicular joint, anterior acromial border, posterior glenoid rim, trapezius and supraspinatus. Right shoulder impingement: Positive. Right shoulder O'brien's test: Positive.** Right shoulder popeye sign: Negative.

Joints-Knee: **Tenderness is present in the vastus medialis, patella and posterior knee of the left knee.**

Neurology - Coordination: Diadochokinesis is found to be normal. Finger-to-nose testing is normal. The

patients gait appears to be normal. The patient was able to do heel walk. The patient was able to do toe walk.
Neurology - Straight Leg Raising: Right: Positive at 35 degrees. Left: Positive at 40 degrees.
Neurology - Motor Strength: Right UE strength - Flexors: 5/5. Left UE strength - Flexors: 5/5. Right UE strength - Extensors: 5/5. Left UE strength - Extensors: 5/5. Right UE Tone: Normal. Left UE Tone: Normal. Right LE strength - Flexors: 5/5. Left LE strength - Flexors: 5/5. Right LE strength - Extensors: 5/5. Left LE strength - Extensors: 5/5. Right LE Tone: Normal. Left LE Tone: Normal.

Pain Management

- Cluneal Nerve Block
- ManaSport+ Device
- Physical Therapy Prescription
- PNEUMATIC COMPRESSION DEVICE
- Sciatic Nerve Injection
- Transcutaneous Electrical Joint Stimulation Device
- Trigger Point injection

Labs Ordered
Order No: LAB0113138 Dated: 09-16-2026 Lab: DME VENDOR
DME: CERVICAL COLLAR, CERVICAL PILLOW, LUMBO SACRAL SUPPORT, LUMBAR BACK CUSHION, BED BOARD/EGG CRATE MATTRESS, HEATING PAD (THERMOPHORE), COLD/HOT WATER CIRCULATION UNIT
Order No: LAB0113139 Dated: 09-16-2026 Lab: DME VENDOR
DME: Orthopedic Positioning Seat

Radiology Ordered
Order No: RAD0055831 Dated: 09-16-2026 Rad: HIGHLINE RADIOLOGY
Magnetic Resonance Imaging: MRI Cervical Spine without CM, MRI Lumbar Spine without CM, MRI Thoracic without CM
Order No: RAD0055832 Dated: 09-16-2026 Rad: FAST CARE MEDICAL DIAGNOSTICS, PLLC
Magnetic Resonance Imaging: MRI Left Knee without contrast, MRI Right Shoulder without CM

Assessment and Plan
ICD: Cervicogenic headache (R51.9)
ICD: Acute cervical myofascial strain (S16.1XXA)
ICD: Acute thoracic myofascial strain (S29.019A)
ICD: Acute lumbosacral myofascial strain (S39.012A)
ICD: Derangement of right shoulder joint (M24.9)
ICD: Derangement of knee, left (M23.92)

Diagnosis
ICD: Cervicogenic headache (R51.9)
ICD: Acute cervical myofascial strain (S16.1XXA)
ICD: Acute thoracic myofascial strain (S29.019A)
ICD: Acute lumbosacral myofascial strain (S39.012A)
ICD: Derangement of right shoulder joint (M24.9)
ICD: Derangement of knee, left (M23.92)
Elements of Medical Decision Making

Assessment: Lumbar Posttraumatic Sprain and Strain Syndrome, Lumbar disc displacement
Patient presents in the office with worsening symptoms of lower back pain. Patient is status post trauma/motor vehicle accident. Patient was asymptomatic prior to the accident.
MRI: PENDING R/O derangement
EMG/NCV: PENDING R/O Nerve compression
As a direct result of the injury patient also endorses difficulties with activities of daily living and a negative impact on functional capacity and range of motion.
Ultrasound of lumbar spine conducted using Butterfly IQ+ transducer.
Muscle patterns of lumbar spine are consistent with diffuse and focal echogenic deflections at paraspinal musculature. This pattern is highly suggestive of muscle spasms of musculature at bilateral L4-L5, L5-S1.

Therapeutic Pain Management injections (ie: trigger point injections, etc) shortly after a motor vehicle accident are being recommended because they are medically necessary and extremely helpful as detailed below.

When a patient experiences whole body trauma associated with a motor vehicle accident, their bodies are subjected to physical forces of acceleration and deceleration which are transmitted throughout their entire body, and typically manifest as more problematic in those areas which are more vulnerable. This often includes the spine, along with shoulder, knee and wrist joints and the muscles and other soft tissues (ligaments, tendons and cartilage) which are in close proximity to these areas.

According to the Cleveland Clinic, many providers use trigger point injections alongside physical therapy and stretching exercises to alleviate myofascial pain. There are significant benefits to speed up recovery by initiating TPI therapy as early as possible after injury or trauma. This strategy is particularly beneficial for those patients who are unable to perform effective physical therapy or stretching exercises due to intense pain and helps amplify and expedite the impact of conservative care. Many of our patients are dealing with acute worsening or persistent lower back, neck and or extremity pain. A motor vehicle accident or mechanical trauma is often deadly or life threatening. After an injury the body's enormous demand to drive forward the healing process is largely hindered by the patient's limited range of motion with body mechanics, acute pain and stiffness. The sequela of these symptoms is further exacerbated by the patients inability to perform and carry out the necessary requirement of physical therapy and stretching exercises. If these steps mentioned above are not initiated as early as possible, the patient is at a higher risk for developing scar tissue and adhesions within the injured muscle groups and tendons. This often leads to permanent decreased range of motion, difficulty with activities of daily living, human suffering and increased disability. Patients who do not receive prompt intervention often end up having a higher incidence of surgical intervention, which we are trying to avoid.

Deploying TPI therapy as early as possible will yield the best possible outcome for recovery utilizing the least number of resources. The efficacy of trigger point injections is intensified when combined with conservative care. This multidimensional pain management technique is often beneficial to decrease both the need and use of strong habit-forming oral pain medications such as Oxycodone, Oxycontin and Morphine.

Plan
- RTC 4 weeks, review TPI/NBI Therapy and plan of care.
- Patient verbally expressed understanding The Following plan has been discussed with the patient in detail:
- U/S guided Trigger point injection therapy to be performed to improve functional capacity and mitigate symptoms of pain to increase capacity for PT/exercises.
- Patient would benefit from consultation for Cervical and Lumbar EMG/NCV testing to ensure no significant nerve damage is present.
- Patient would benefit from functional capacity testing and range of motion testing to gauge progress/improvement.

- MRI without contrast of the Cervical, Thoracic, Lumbar, R Shoulder, L Knee regions Ordered.

- Patient will continue Chiropractic, Physical Therapy, Acupuncture 3 to 4 times/week for 4-6 weeks.

-Psychiatric evaluation requested in an effort to diagnose and treat persistent HA, anxiety, depression or sleep disorders.

Rx Prescribed today:
-cyclobenzaprine 7.5 mg tablet 1 Tablet Once A Day PRN for 30 Days
-diclofenac 3 % topical gel
-lidocaine 5 % topical ointment
-Ibuprofen 800 mg oral tablet

The following DME's were ordered:
-ManaSport+ Device
-PNEUMATIC COMPRESSION DEVICE
-Transcutaneous Electrical Joint Stimulation Device

-CERVICAL COLLAR
-CERVICAL PILLOW
-LUMBAR BACK CUSHION
-LUMBO SACRAL SUPPORT
-BED BOARD
-EGG CRATE MATTRESS
-HEATING PAD (THERMOPHORE)
-COLD/HOT WATER CIRCULATION UNIT
-Orthopedic Positioning Seat

Referral sent for functional capacity test, range of motion , acupuncture , chiropractor. Neurology, orthopedic , interventional management as indicated

- RTC 4 weeks, review TPI, IAI, NBI Therapy and plan of care.
- Patient verbally expressed understanding and verbally agreed to plan.

Procedure
TODAYS VISIT: TYPE: (Ultrasound Guided Trigger Point Injection Therapy Lumbar Spine)
Procedure/Interventions
Bilateral CLUNEAL NERVE BLOCK injection was performed on this visit.
- I prepped and filled syringe with 1cc Lidocaine, 0.5cc Bupivacaine, 0.5cc Dexamethasone and administered to the areas listed above.
- Informed Consent obtained
- Risks of NBI discussed with patient, including but not limited to infection and allergic reaction discussed
- Sterile prep and 30G needle were used.
- Patient tolerated procedure well and endorses pain relief, no complications.
- Will continue to manage and follow consistently, will evaluate and assess for level of improvement on upcoming visits.

TODAYS VISIT: TYPE: (Ultrasound Guided Trigger Point Injection Therapy Lumbar Spine)
Procedure/Interventions

Bilateral Erector Spinae muscles Ultrasound Guided Trigger Point injection was performed on this visit.
- I prepped and filled syringe with 1cc Lidocaine, 0.5cc Bupivacaine, 0.5cc Dexamethasone and administered to the areas listed above.

- Informed Consent obtained
- Risks of TPI discussed with patient, including but not limited to infection and allergic reaction discussed
- Sterile prep and 30G needle were used.
- Patient tolerated procedure well and endorses pain relief, no complications.
- Will continue to manage and follow consistently, will evaluate and assess for level of improvement on upcoming visits.

TODAYS VISIT: TYPE: (Ultrasound Guided Sciatic Nerve Block Injection Therapy)
- Informed Consent obtained
- Sterile prep and 30G needle were used.
- Patient tolerated procedure well and endorses pain relief.
-U/S guided Bilateral Sciatic Nerve Block Injection

Medications used:
1cc of lidocaine, 0.5cc of bupivacaine, 0.5cc of dexamethasone

Multiple sites for trigger point injections are indicated due to the complex nature of whole-body trauma and the body's natural tendency to resist the applied forces during an accident. Clinically significant trigger points or knots are identified with ultrasound guidance carefully noting echogenic deflections within specified muscle groups that are so often characteristic of muscle spasms. Muscles that are in proximity or adjacent need to be treated due to physical mechanics innervation and function of these muscle groups. Patients may receive up to six trigger point injections, although many suffer from more than ten clinically significant myofascial knots that would clearly benefit from treatment.
The diagnosis of trigger points is not subjective, and their presence is confirmed two different ways, by physical exam and by ultrasound. Diagnosis of a trigger point is NOT adequate by itself. One must know the precise anatomic location so that medication can be delivered by needle injection to exact area which requires treatment in a safe and effective way.

Safety: Many therapeutic injections are performed in areas with associated nerves, arteries or veins. Avoiding these structures is very important from a safety perspective.

Efficacy: We utilize dexamethasone sodium phosphate, bupivacaine and lidocaine medications when performing therapeutic injections. If they are not introduced to the precise location, it will not be effective and can possibly create other problems. Visualization with ultrasound helps ensure clinical efficacy is optimized.


CPT/HCPC Codes:

Name	Code	Units	Comments
Office O/p New Mod 45 Min	99204	1.00	
Prolong Service W/o Contact	99358	1.00	
Njx 1/Mlt Trigger Point 1/2	20552	1.00	
Njx Aa&/strd Other Pn/branch	64450	2.00	
Njx Aa&/strd Sciatic Nrv Img	64445	2.00	
Injection Dexamethosone Sodium Phosphate 1 Mg	J1100	6.00	
Inj, bupivacaine, nos, 0.5mg	J0665	15.00	
Injection, Lidocaine 1 mg	J3490	30.00	
Echo Guide For Biopsy	76942	1.00	
Echo Guide For Biopsy	76942	1.00	
Echo Guide For Biopsy	76942	1.00	
Echo Guide For Biopsy	76942	1.00	

Echo Guide For Biopsy	76942	1.00	
Echo Guide For Biopsy	76942	1.00	

Prescription
lidocaine 5 % topical ointment Apply 2-3 grams to affected area every 8 Hours PRN Pain for 30 Days , Prescribe 250 Gram
diclofenac 3 % topical gel 2-3 Gram Every 8 Hours PRN for painful areas, may alternate with Lidocaine ointment, por dolor musculo, Prescribe 200 Gram
ibuprofen 800 mg tablet 1 Tablet Every 8 Hours FOR SYSTEMIC INFLAMMATION And Pain for 15 Days , Prescribe 90 Tablet
cyclobenzaprine 7.5 mg tablet Take 1 tablet by mouth once daily at bed time (MUSLE RELAXER-MAY CAUSE DROWSINESS) for 30 Days , Prescribe 30 Tablet

Follow up: 4 Weeks



Jerry Samedi, PA-C
NPI: 1104784701
This has been electronically signed by Jerry Samedi, PA-C on 09-16-2026.

Ultrasound-guided trigger point injections in the cervicothoracic musculature: a new and unreported technique.
Botwin KP, et al. Pain Physician. 2008. PMID: 19057634
Ultrasound-Guided 5-in-1 Injection: Trigger Point Injections and Nerve Hydrodissections for Nonspecific Upper Back Pain.
Tang TY, et al. Am J Phys Med Rehabil. 2019. PMID: 30461431 No abstract available.
Ultrasound-guided transversus abdominis plane block vs trigger point injections for chronic abdominal wall pain: a randomized clinical trial.
Moeschler SM, et al. Pain. 2021. PMID: 33433147 Clinical Trial.

Ultrasound-guided interventional procedures for myofascial trigger points: a systematic review.
Diep D, Chen KJQ, Kumbhare D.Reg Anesth Pain Med. 2021 Jan;46(1):73-80. doi: 10.1136/rapm-2020-101898. Epub 2020 Nov 6.PMID: 33159004 Review

Ultrasound-Guided Musculoskeletal Injections.
Lin JS, Gimarc DC, Adler RS, Beltran LS, Merkle AN.Semin Musculoskelet Radiol. 2021 Dec;25(6):769-784. doi: 10.1055/s-0041-1740349. Epub 2021 Dec 22.PMID: 34937117

Ultrasound-guided transversus abdominis plane block vs trigger pointinjections for chronic abdominal wall pain: a randomized clinical trial.
Moeschler SM, Pollard EM, Pingree MJ, Pittelkow TP, Bendel MA, Mauck WD, Watson JC, Eldrige JS, Loftus CG, Hooten WM.Pain. 2021 Jun 1;162(6):1800-1805. doi: 10.1097/j.pain.0000000000002181.PMID: 33433147 Clinical Trial.

Trigger Point Injection.
Hammi C, Schroeder JD, Yeung B.2021 Sep 13. In: StatPearls . Treasure Island (FL): StatPearls Publishing; 2022 Jan-.PMID: 31194353 Free Books & Documents

Sonography-guided trigger point injections in abdominal myofascial pain syndrome.
Rhim HC, Cha JH, Cha J, Kim DH.Medicine (Baltimore). 2020 Dec 4;99(49):e23408. doi: 10.1097/MD.0000000000023408.PMID: 33285730 Free PMC article.

Zhang WB, Yao DW, Wu WX.Zhongguo Zhen Jiu. 2019 Aug 12;39(8):867-70. doi: 10.13703/j.0255-2930.2019.08.017.PMID: 31397133 Clinical Trial. Chinese.

Ultrasound-Guided Three-in-One Trigger Point Injection for Muscles of the Posterior Axillary Fold.
Chang MC, Boudier-Revéret M, Hsiao MY.Pain Pract. 2020 Jan;20(1):110-111. doi: 10.1111/papr.12819. Epub 2019 Sep 13.PMID: 31310698 No abstract available.

Ultrasound-guided trigger point injection for piriformis syndrome in the emergency department
Aquino-Jose VM, Blinder V, Johnson J, Havryliuk T.J Am Coll Emerg Physicians Open. 2020 Jun 24;1(5):876-879. doi: 10.1002/emp2.12153. eCollection 2020 Oct.PMID: 33145535

Feasibility of Ultrasound-Guided Trigger Point Injection in Patients with Myofascial Pain Syndrome.
Kang JJ, Kim J, Park S, Paek S, Kim TH, Kim DK.Healthcare (Basel). 2019 Oct 15;7(4):118. doi: 10.3390/healthcare7040118.PMID: 31618922

A novel, nonopiod-based treatment approach to men with urologic chronic pelvic pain syndrome using ultrasound-guided nerve hydrodissection and pelvic floor musculature trigger point injections.
Hui J, Seko K, Shrikhande G, Ahmed T, Hill C, Reutter C, Shrikhande A.Neurourol Urodyn. 2020 Feb;39(2):658-664. doi: 10.1002/nau.24242. Epub 2019 Dec 2.PMID: 31793050

Ultrasound-guided interfascial blocks of the trapezius muscle for cervicogenic headache: A report of two cases.
Arc T.Agri. 2021 Oct;33(4):278-281. doi: 10.14744/agri.2020.08831.PMID: 34671952 English.

Clinical indications for image-guided interventional procedures in the musculoskeletal system: a Delphi-based consensus paper from the European Society of Musculoskeletal Radiology (ESSR)-Part II, elbow and wrist.
Sconfienza LM, Adriaensen M, Albano D, Aparisi Gómez MP, Bazzocchi A, Beggs I, Bignotti B, Chianca V, Corazza A, Dalili D, De Dea M, Del Cura JL, Di Pietto F, Drakonaki E, Facal de Castro F, Filippiadis D, Gielen J, Gitto S, Gupta H, Klauser AS, Lalam R, Martin S, Martinoli C, Mauri G, McCarthy C, McNally E, Melaki K, Messina C, Mirón Mombiela R, Neubauer B, Obradov M, Olchowy C, Orlandi D, Plagou A, Prada Gonzalez R, Rutkauskas S, Snoj Z, Tagliafico AS, Talaska A, Vasilevska-Nikodinovska V, Vucetic J, Wilson D, Zaottini F, Zappia M, Allen G; Ultrasound and Interventional Subcommittees of the European Society of Musculoskeletal Radiology (ESSR).Eur Radiol. 2020 Apr;30(4):2220-2230. doi: 10.1007/s00330-019-06545-6. Epub 2019 Dec 16.PMID: 31844963

Pain and functionality improved when underlying neuromuscular dysfunction addressed in chronic pelvic pain patients.
Natarajan J, Ahmed T, Patil S, Mamsaang M, Kapadia R, Tailor Y, Shrikhande A.Neurourol Urodyn. 2021 Aug;40(6):1609-1615. doi: 10.1002/nau.24726. Epub 2021 Jun 3.PMID: 34082473

Ultrasound guided serrates posterior superior muscle block relieves interscapular myofascial pain
Taketa Y, Irisawa Y, Fujitani T.J Clin Anesth. 2018 Feb;44:10-11. doi: 10.1016/j.jclinane.2017.10.009.PMID: 29078063 No abstract available.

An Original Approach for the Treatment of Varicose Veins of the Lower Limbs.
Sotiris D, Pallotta G, Nittari G, Amenta F.J Clin Aesthet Dermatol. 2020 Aug;13(8):E59-E64. Epub 2020 Aug 1.PMID: 33178385

Expansion of 1 mL of Solution by Ultrasound-Guided Injection Between the Trapezius and Rhomboid Muscles: A Cadaver Study.
Kimura H, Kobayashi T, Zenita Y, Kurosawa A, Aizawa S.Pain Med. 2020 May 1;21(5):1018- 1024. doi: 10.1093/pm/pnz170.PMID: 31504919

Low Back Pain Caused by Iliopsoas Tendinopathy Treated with UltrasoundGuided Local Injection of Anesthetic and Steroid: A Retrospective Study.
Zhu Z, Zhang J, Sheng J, Zhang C, Xie Z.J Pain Res. 2020 Nov 20;13:3023-3029. doi: 10.2147/JPR.S281880. eCollection 2020.PMID: 33244260

Oliván-Gonzalvo G, Ruiz-Serrano J, Uclés-Moreno P, Fernández-Sanz A.An Sist Sanit Navar. 2021 Aug 20;44(2):303-307. doi: 10.23938/ASSN.0952.PMID: 34132249 Spanish.

Anterior Cutaneous Nerve Entrapment Syndrome in Children: A Prospective Observational Study
Kifer T, Mišak Z, Jadrešin O, Hojsak I.Clin J Pain. 2018 Jul;34(7):670-673. doi: 10.1097/AJP.0000000000000573.PMID: 29189217

Ultrasound-Guided Intra-articular Injection of the Radio-ulnar and Radio-humeral Joints and Ultrasound-Guided Dry Needling of the Affected Limb Muscles to Relieve Fixed Pronation Deformity and Myofascial Issues around the Shoulder, in a Case of Complex Regional Pain Syndrome Type 1.
Pai RS, Vas L.Pain Pract. 2018 Feb;18(2):273-282. doi: 10.1111/papr.12596. Epub 2017 Jun 14.PMID: 28434187

Ultrasound-Guided Botulinum Toxin Injection for the Treatment of Teres Major Fasciculations.
Cushman DM, Carefoot A, Rees MR.Am J Phys Med Rehabil. 2022 Jul 1;101(7):e112-e114. doi: 10.1097/PHM.0000000000002011. Epub 2022 Mar 15.PMID: 35302525

Interventions for myofascial pain syndrome in cancer pain: recent advances: why, when, where and how.
Thottungal A, Kumar P, Bhaskar A.Curr Opin Support Palliat Care. 2019 Sep;13(3):262-269. doi: 10.1097/SPC.0000000000000446.PMID: 31348012 Review.

Identifying the Dorsal Scapular Artery Optimizes the Safety and Precision of the UltrasoundGuided 5-in-1 Injection. Tang TY, Wu CH.Am J Phys Med Rehabil. 2019 Jul;98(7):e80-e81. doi: 10.1097/PHM.0000000000001137.PMID: 30640288 No abstract available.

Prospective Audit of a Pathway for In-Patient Pain Management of Chronic Abdominal Pain: A Novel and Cost-Effective Strategy
Niraj G, Chaudhri S.Pain Med. 2018 Mar 1;19(3):589-597. doi: 10.1093/pm/pnx118.PMID: 28521010

Ultrasound-Guided Intermediate Cervical Plexus Block With Depot Steroids in the Management of Refractory Neck Pain Secondary to Cervicothoracic Myofascial Pain Syndrome: A Case Series
Thawale R, Alva S, Niraj G.A A Pract. 2019 Dec 15;13(12):446-449. doi: 10.1213/XAA.0000000000001102.PMID: 31592828

Organic Semiconducting Photoacoustic Nanodroplets for LaserActivatable Ultrasound Imaging and Combinational Cancer Therapy.
Tang W, Yang Z, Wang S, Wang Z, Song J, Yu G, Fan W, Dai Y, Wang J, Shan L, Niu G, Fan Q, Chen X.ACS Nano. 2018 Mar 27;12(3):2610-2622. doi: 10.1021/acsnano.7b08628. Epub 2018 Feb 21.PMID: 29451774

Neuromuscular treatment approach for women with chronic pelvic pain syndrome improving pelvic pain and

functionality
Patil S, Daniel G, Vyas R, Tailor Y, Howell M, Ahmed T, Reutter C, Shrikhande A.Neurourol Urodyn. 2022 Jan;41(1):220-228. doi: 10.1002/nau.24799. Epub 2021 Sep 16.PMID: 34529875

First Case of Occipital Neuralgia Treated by Fascial Hydrodissection
Kaga M.Am J Case Rep. 2022 May 17;23:e936475. doi: 10.12659/AJCR.936475.PMID: 35578561

Efficacy of extracorporeal shock waves in the treatment of myofascial pain syndrome: a systematic review and meta-analysis of controlled clinical studies.
Wu T, Li S, Ren J, Wang D, Ai Y.Ann Transl Med. 2022 Feb;10(4):165. doi: 10.21037/atm-22- 295.PMID: 35280414

Reviewed by:



Mark Filstein, M.D.
NPI No.: 1649396110

This has been electronically signed by Mark Filstein, M.D. on 09-21-2026 07:40:02.

Outcome assessment Test reviewed on 09-21-2026 07:40:02.

Extracorporeal Shock Wave & RPW Referral / Treatment

PATIENT NAME: X Rosangela

PATIENT SIGNATURE: X Rosangela Y.

DATE OF SERVICE: 10/27/26 DATE OF ACCIDENT: 09/07/26

LOCATION: _____

PROCEDURE # ____1 ____2 ____3 ____4 ____5

___NECK ✓LOW BACK ___THORACIC ___LT/RT SHOULDER ___LT/RT KNEE

___OTHER_____

SHOCKWAVE THERAPY

DX CODES:

☐ M54.2 Cervicalgia

☐ M54.6 Pain in Thoracic

☑ M54.5 Low Back Pain

☐ M54.41 Lumbago with Sciatica R side

☐ M54.42 Lumbago with Sciatica L side

☐ M25.561 Pain in Right Knee Joint

☐ M25.562 Pain in Left knee joint

☐ M25.511 Pain in Right Shoulder

☐ M25.512 Pain in Left shoulder

☐ M79.604 Pain in Right Leg

☐ M79.605 Pat in Left Leg

OTHER _____

Referring Physician: dr Marc Parnes _____

Name **Signature**

Consent for Shock Wave Therapy Treatment (ESWT) and Hold Harmless Agreement

I understand that I am a patient of _____

Cooperation with treatment:
In order, for Shock Wave Therapy treatment to be effective, I must come to scheduled appointments unless there are unforeseen circumstances. I understand and agree to cooperate with and perform the home physical therapy program intended for me. If I have trouble with any part of my treatment program, I will discuss it with my provider.

No warranty:
I understand that there are no guarantees regarding a cure for or improvement in my condition. understand that my provider will outline and discuss the goals of Shock Wave Therapy treatment for my condition and will discuss treatment options with me before I consent to treatment.

Informed consent for treatment:
The term "informed consent" means that the potential risks, benefits have been explained to me.

Potential risks:
Side effects from ESWT are limited to mild bruising, swelling, pain, numbness or tingling in the treated area, and the recovery is minimal compared with that of surgical intervention. This discomfort is usually temporary; if it does not subside in a reasonable time, I agree to contact my physician.

Potential benefits:
I may experience an improvement in my symptoms and an increase in my ability to perform daily activities. I may experience increased strength, awareness, flexibility, and endurance in my movements. I may experience decreased pain and discomfort. I should gain a greater knowledge about managing my condition and the resources available to me

Alternatives: If I do not wish to participate in the ESWT program, I will discuss my medical, surgical, or pharmacological alternatives with my physician or primary care provider.

Payment: I understand that I am responsible for any charges not covered by insurance.

I have read the above information and consent to ESWT.
I hereby release and hold harmless _____for any loss/judgement and/or harm which might come to me because of this Shock Wave Therapy treatment (ESWT). I waive all claims associated with this treatment which I may have against _____, his employees, officers, directors, and agents.

Patient Name (Print): X Rosangela

Patient Signature: X Rosangela

Parent/Guardian Signature: _____

Date: _____

NEW YORK MOTOR VEHICLE NO-FAULT INSURANCE LAW
ASSIGNMENT OF BENEFITS FORM

(FOR ACCIDENTS OCCURRING ON AND AFTER 3/1/02)

I, YANEZ ANA, ("Assignor") hereby assign to NORMAN MEDICAL, P.C. ("Assignee")
(Print patient's name) (Print hospital or health care provider name)
all rights privileges and remedies to payment for health care services provided by assignee to which I am entitled under Article 51 (the No-Fault statute) of the Insurance Law.

The Assignee hereby certifies that they have not received any payment from or on behalf of the Assignor and shall not pursue payment directly from the Assignor for services provided by said Assignee for injuries sustained due to the motor vehicle accident which occurred on 09/27/25, not withstanding any other agreement
(Print accident date)
to the contrary.

This agreement may be revoked by the assignee when benefits are not payable based upon the assignor's lack of coverage and/or violation of a policy condition due to the actions or conduct of the assignor.

ANY PERSON WHO KNOWINGLY AND WITH INTENT TO DEFRAUD ANY INSURANCE COMPANY OR OTHER PERSON FILES AN APPLICATION FOR COMMERCIAL INSURANCE OR A STATEMENT OF CLAIM FOR ANY COMMERCIAL OR PERSONAL INSURANCE BENEFITS CONTAINING ANY MATERIALLY FALSE INFORMATION, OR CONCEALS FOR THE PURPOSE OF MISLEADING, INFORMATION CONCERNING ANY FACT MATERIAL THERETO, AND ANY PERSON WHO, IN CONNECTION WITH SUCH APPLICATION OR CLAIM, KNOWINGLY MAKES OR KNOWINGLY ASSISTS, ABETS, SOLICITS OR CONSPIRES WITH ANOTHER TO MAKE A FALSE REPORT OF THE THEFT, DESTRUCTION, DAMAGE OR CONVERSION OF ANY MOTOR VEHICLE TO A LAW ENFORCEMENT AGENCY, THE DEPARTMENT OF MOTOR VEHICLES OR AN INSURANCE COMPANY, COMMITS A FRAUDULENT INSURANCE ACT, WHICH IS A CRIME, AND SHALL ALSO BE SUBJECT TO A CIVIL PENALTY NOT TO EXCEED FIVE THOUSAND DOLLARS AND THE VALUE OF THE SUBJECT MOTOR VEHICLE OR STATED CLAIM FOR EACH VIOLATION.

X Rosangela
(Print name of Patient)

X Rosangela
(Signature of Patient)

10/27/25
(Date of signature)

(Address of Patient)

NORMAN MEDICAL, P.C.
(Print name of Provider)

(Signature of Provider)

714 SENECA AVENUE

10/27/25
(Date of signature)

RIDGEWOOD NY 11385
(Address of Provider)

NYS FORM NF-AOB (Rev 1/2004)

Patient Name: YANEZ ANA

MR#: _____ Date of Accident: 09/07/26

MEDICAL LIEN

Attorney Name:

_____

Patient Name: YANEZ ANA

I hereby authorize and direct my attorney, to pay directly to _____ such sums as may be due and owing for professional services rendered to me both by reason of this accident and by reason of any other bills that are due to the provider and to withhold such sums from any settlement of judgment as is necessary to adequately protect the provider.
I hereby further give a lien to the provider on any proceeds to which I may become entitled as a result of any settlement of judgment in any claim or litigation arising out of the injuries for which I have been treated of injuries in connection therewith, whether such proceeds are remitted directly to me or to you, my attorney.

I fully understand that I am directly responsible to the provider for all professional bills submitted by the provider for services rendered to me by the provider and that this agreement is made solely for the providers' additional protection and in consideration of the provider awaiting payment. I further understand that such payment is not contingent on any settlement, judgment, or verdict by which I may eventually recover said fee

Attorney agrees to notify the doctors immediately of the name and contacting information of any attorney substituted in his or her place.

X Rosa

Print Patient Name

10/27/26

Date

X Rosangela Y

Patient's Signature

_____

Signature of Parent / Guardian

ACKNOWLEDGEMENT OF ASSIGNMENT & LIEN BY ATTORNEY

The undersigned being the attorney of record on his own behalf and on behalf of any other attorney or attorneys who are associated with the undersigned or who are substituted in his stead for the above patient, does hereby agree to observe all the terms of the above and agrees to withhold such sums from any settlement, judgment or verdict as may be necessary to adequately protect.

_____

Attorney's Signature

_____

Date

*NOTE TO ATTORNEY

Please sign and return one copy to the provider's office; keep a copy for your records.