

Math 109 HW3

Neo Lee

02/14/2023

Problem 6.5

(i)

Proposition 1. $A \subseteq B \Leftrightarrow A \cup B = B$.

Proof. (\Rightarrow ; $A \cup B \subseteq B$) $\forall x \in A \cup B, x \in B$ because $A \subseteq B$.

(\Rightarrow ; $B \subseteq A \cup B$) By definition, $\forall y \in B, y \in B \cup S$ for any arbitrary set S . Therefore, $B \subseteq A \cup B$.

Since $A \cup B \subseteq B$ and $B \subseteq A \cup B$, $A \cup B = B$, and (\Rightarrow) is proved.

(\Leftarrow) By definition, $\forall z \in A, z \in A \cup S$ for any arbitrary set S , which means $A \subseteq A \cup S$. Hence, $A \subseteq A \cup B$, which is equivalent to $A \subseteq B$. \square

(ii)

Proposition 2. $A \subseteq B \Leftrightarrow A \cap B = A$.

Proof. (\Rightarrow ; $A \cap B \subseteq A$) By definition, $\forall x \in A \cap B, x \in A$, thus $A \cap B \subseteq A$.

(\Rightarrow ; $A \subseteq A \cap B$) $\forall y \in A, y \in A \cap B$ because $A \subseteq B$.

Since $A \cap B \subseteq A$ and $A \subseteq A \cap B$, $A \cap B = A$, and (\Rightarrow) is proved.

(\Leftarrow) By definition, $(B \cap S) \subseteq B$ for any arbitrary set S . Hence, $A = A \cap B \subseteq B$. \square

Problem 6.6

Proposition 3. If $A \cap B \subseteq C$ and $x \in B$, then $x \notin A - C$.

Proof. Assume to the contrary that if $A \cap B \subseteq C$ and $x \in B$, then $x \in A - C$. It means that $x \in A$ and $x \notin C$. Since $A \cap B \subseteq C$, $x \notin C \Rightarrow x \notin A \cap B$. We know $x \in A$ and $x \notin A \cap B$, therefore, $x \in A \cap B^c$. It means $x \in B^c \Rightarrow x \notin B$, which contradicts that $x \in B$. \square

Problem 6.7

Proposition 4. For subsets of a universal set U , $A \subseteq B$ if and only if $B^c \subseteq A^c$.

Proof. $A \subseteq B$ means that for an arbitrary x , if $x \in A$, then $x \in B$. Logically, it is equivalent to its contrapositive, which states for an arbitrary x , if $x \notin B$, then $x \notin A$. $x \notin B$ can be written as $x \in B^c$, and $x \notin A$ can be written as $x \in A^c$. Therefore, the entire statement can be rewritten as for an arbitrary x , if $x \in B^c$, then $x \in A^c$, which is the definition of $B^c \subseteq A^c$. \square

Problem 7.1

(i) $m = \mathbb{Z}^+$

(ii) $m = \{1\}$

(iii) $m = \mathbb{Z}^+$

(iv) $n = \emptyset$

Problem 7.2

(i)

Proposition 5. *Disproving $\forall m, n \in \mathbb{Z}^+, m \leq n$ means proving $\exists m, n \in \mathbb{Z}^+, m > n$.*

Proof. Let $m = 3$ and $n = 2$, $m > n$. □

(ii)

Proposition 6. $\exists m, n \in \mathbb{Z}^+, m \leq n$.

Proof. Let $m = 2$ and $n = 3$, $m \leq n$. □

(iii)

Proposition 7. $\forall m \in \mathbb{Z}^+, \exists n \in \mathbb{Z}^+, m \leq n$.

Proof. Let $n = m$. $\forall m \in \mathbb{Z}^+, m = n \Rightarrow m \leq n$. □

(iv)

Proposition 8. $\exists m \in \mathbb{Z}^+, \forall n \in \mathbb{Z}^+, m \leq n$.

Proof. Let $m = 1$. $\forall n \in \mathbb{Z}^+, m \leq n$. □

(v)

Proposition 9. $\forall n \in \mathbb{Z}^+, \exists m \in \mathbb{Z}^+, m \leq n$.

Proof. Let $m = 1$. $\forall n \in \mathbb{Z}^+, m \leq n$. □

(vi)

Proposition 10. *Disproving $\exists n \in \mathbb{Z}^+, \forall m \in \mathbb{Z}^+, m \leq n$ means proving $\forall n \in \mathbb{Z}^+, \exists m \in \mathbb{Z}^+, m > n$.*

Proof. Let $m = n + 1$. $\forall n \in \mathbb{Z}^+, m > n$. □

Problem 7.4

(i)

Proposition 11. $\forall x \in \mathbb{R}, \exists y \in \mathbb{R}, x + y = 0$.

Proof. Let $y = -x$. $\forall x \in \mathbb{R}, x + y = x - x = 0$. □

(ii)

Proposition 12. *Disproving $\exists y \in \mathbb{R}, \forall x \in \mathbb{R}, x + y = 0$ mean proving $\forall y \in \mathbb{R}, \exists x \in \mathbb{R}, x + y \neq 0$.*

Proof. Let $x = -y + 1$. $\forall y \in \mathbb{R}, y + x = y - y + 1 = 1 \neq 0$. □

(iii)

Proposition 13. $\forall x \in \mathbb{R}, \exists y \in \mathbb{R}, xy = 0$.

Proof. Let $y = 0$. $\forall x \in \mathbb{R}, xy = x \cdot 0 = 0$. □

(iv)

Proposition 14. $\exists y \in \mathbb{R}, \forall x \in \mathbb{R}, xy = 0$.

Proof. Let $y = 0$. $\forall x \in \mathbb{R}, xy = x \cdot 0 = 0$. □

(v)

Proposition 15. *Disproving $\forall x \in \mathbb{R}, \exists y \in \mathbb{R}, xy = 1$ means proving $\exists x \in \mathbb{R}, \forall y \in \mathbb{R}, xy \neq 1$.*

Proof. Let $x = 0$. $\forall y \in \mathbb{R}, xy = 0 \cdot y = 0 \neq 1$. □

(vi)

Proposition 16. *Disproving $\exists y \in \mathbb{R}, \forall x \in \mathbb{R}, xy = 1$ means proving $\forall y \in \mathbb{R}, \exists x \in \mathbb{R}, xy \neq 1$.*

Proof. Let $x = 0$. $\forall y \in \mathbb{R}, xy = 0 \cdot y = 0 \neq 1$. □

(vii)

Proposition 17. $\forall n \in \mathbb{Z}^+, (n \text{ is even or } n \text{ is odd})$.

Proof. $\forall n \in \mathbb{Z}^+$, n is either even or n is not even. By definition, if n is not even, then n is odd, which logically means n is even or n is odd. □

(viii)

Proposition 18. *Disproving $(\forall n \in \mathbb{Z}^+, n \text{ is even})$ or $(\forall n \in \mathbb{Z}^+, n \text{ is odd})$ means proving $(\exists n \in \mathbb{Z}^+, n \text{ is odd})$ and $(\exists n \in \mathbb{Z}^+, n \text{ is even})$.*

Proof. For the first half of the statement, let $n = 1$, then n is odd, which proves $(\forall n \in \mathbb{Z}^+, n \text{ is odd})$. For the second half of the statement, let $n = 2$, then n is even, which proves $(\exists n \in \mathbb{Z}^+, n \text{ is even})$. □

Problem 7.7

Page 115 Problem 4

Proposition 19. $A \cap B = A \cap C$ and $A \cup B = A \cup C$ if and only if $B = C$.

Proof. (\Rightarrow)

$$B = B \cap (A \cup B) \tag{1}$$

$$= B \cap (A \cup C) \quad (\because A \cup B = A \cup C) \tag{2}$$

$$= (B \cap A) \cup (B \cap C) \tag{3}$$

$$= (A \cap B) \cup (B \cap C) \tag{4}$$

$$= (A \cap C) \cup (B \cap C) \quad (\because A \cap B = A \cap C) \tag{5}$$

$$= (A \cup B) \cap C \tag{6}$$

$$= (A \cup C) \cap C \quad (\because A \cup B = A \cup C) \tag{7}$$

$$= C \tag{8}$$

(\Leftarrow) This is apparent because we only need to substitute B with C , then we will get $A \cap B = A \cap C$ and $A \cup B = A \cup C$. □

Page 117 Problem 13