



# Goals For This Unit

- Introduce Graph Theory (Section 11.1) – **This PPT.**



# Goals For This Unit

- Introduce Graph Theory (Section 11.1) – This PPT.
- Introduce Trees (Section 12.1)



# Goals For This Unit

- Introduce Graph Theory (Section 11.1) – This PPT.
- Introduce Trees (Section 12.1)
- Discuss Rooted Trees (Section 12.2)



# Goals For This Unit

- Introduce Graph Theory (Section 11.1) – This PPT.
- Introduce Trees (Section 12.1)
- Discuss Rooted Trees (Section 12.2)
- Discuss Sorting (Section 12.3)



# Goals For This Unit

- Introduce Graph Theory (Section 11.1) – This PPT.
- Introduce Trees (Section 12.1)
- Discuss Rooted Trees (Section 12.2)
- Discuss Sorting (Section 12.3)
- **Discuss Weighted Trees and Prefix Codes (Section 12.3).**



# What is Graph Theory?

- The study of graphs.
  - Informally, graphs are mathematical structures used to model binary relations between elements of a given set.

# What is Graph Theory?

- The study of graphs.
  - Informally, graphs are mathematical structures used to model binary relations between elements of a given set.
- The terminology of Graph Theory is nicely summarized at Wikipedia  
[http://en.wikipedia.org/wiki/Glossary\\_of\\_graph\\_theory](http://en.wikipedia.org/wiki/Glossary_of_graph_theory)




# What is Graph Theory?

- The study of graphs.
  - Informally, graphs are mathematical structures used to model binary relations between elements of a given set.
- The terminology of Graph Theory is nicely summarized at Wikipedia  
[http://en.wikipedia.org/wiki/Glossary\\_of\\_graph\\_theory](http://en.wikipedia.org/wiki/Glossary_of_graph_theory)
- The origin of the theory is credited to Euler, 1736 – The Königsberg Bridge Problem.


# The Königsberg Bridge Problem

- Is there a route which crosses each of the seven bridges exactly once?





# The Königsberg Bridge Problem

- Is there a route which crosses each of the seven bridges exactly once?
- NO.



# The Königsberg Bridge Problem


- Schematic:







# The Königsberg Bridge Problem

- Schematic:





- Graph:



# The Königsberg Bridge Problem

- **Question:** Can one trace this graph starting and ending at the same point without lifting the pencil from the paper, traversing each edge exactly once?



# Aside: Terminology

- You must be familiar with the basic terminology – there's lots of it in Graph Theory.
- You are responsible for the nomenclature handout emailed to you today.
- Again, see Wikipedia.

## Basic Def<sup>n</sup>

Consider a graph  $G$ , and let  $v$  and  $w$  be vertices in  $G$ .

- A walk from  $v$  to  $w$  is a finite alternating sequence of adjacent vertices and edges of  $G$  and has the form

$$v_0 e_1 v_1 e_2 \dots v_{n-1} e_n v_n$$

where the  $v$ 's represent vertices, the  $e$ 's edges,  $v_0 = v$ ,  $v_n = w$ , and  $i = 1, 2, \dots, n$ ,  $v_{i-1}$  &  $v_i$  are the endpoints of  $e_i$ .

- The trivial walk from  $v$  to  $v$  consists of the single vertex  $v$ .
- A path from  $v$  to  $w$  is a walk from  $v$  to  $w$  that does not contain a repeated edge. (a walk s.t.  $e_i \neq e_j$   $\forall i \neq j$ ).
- A simple path from  $v$  to  $w$  is a path that does not contain a repeated vertex. (a walk s.t. either with  $v \neq w$  or  $v = w$  and  $n > 1$ ).
- A closed walk is a walk that starts and ends @ same vertex.
- A circuit is a closed walk that does not contain a repeated edge (a walk where  $v_0 = v_n$ ,  $e_i$  distinct).
- A simple circuit is a circuit w/ no repeated vertex except the 1<sup>st</sup> and last.

# The Königsberg Bridge Problem

**Informally :**

- Begin at vertex A.



# The Königsberg Bridge Problem

## Informally :

- Begin at vertex A.
- Each time you pass through any *other* vertex, you use up two edges (arrive and depart).

# The Königsberg Bridge Problem

## Informally:

- Begin at vertex A.
- Each time you pass through any *other* vertex, you use up two edges (arrive and depart).
- Thus, if it is possible to find such a route that uses all the edges of the graph and starts and ends at A, then the *total* number of arrival and departures from each vertex must be a multiple of two (*i.e.*,  $\deg(v_i) = 2n$ )

# The Königsberg Bridge Problem

## Informally :

- Begin at vertex A.
- Each time you pass through any *other* vertex, you use up two edges (arrive and depart).
- Thus, if it is possible to find such a route that uses all the edges of the graph and starts and ends at A, then the *total* number of arrival and departures from each vertex must be a multiple of two (*i.e.*,  $\deg(v_i) = 2n$ )
- But,  $\deg(B) = 5$ ,  $\deg(C) = \deg(D) = 3$  ( $5+3+3$  not even).

# The Königsberg Bridge Problem

## Informally :

- Begin at vertex A.
- Each time you pass through any other vertex, you use up two edges (arrive and depart).
- Thus, if it is possible to find such a route that uses all the edges of the graph and starts and ends at A, then the *total* number of arrival and departures from each vertex must be a multiple of two (*i.e.*,  $\deg(v_i) = 2n$ )
- But,  $\deg(B) = 5$ ,  $\deg(C) = \deg(D) = 3$  ( $5+3+3$  not even).
- Hence, there is no such route. Thus, it is impossible to travel around Königsberg crossing each bridge exactly once.



# Fundamental Concepts

## Definition: Graph

A graph  $G$  is an ordered pair of disjoint sets,  $(V, E)$  such that  $E$  is a subset of the set of unordered pairs of  $V$ .

# Fundamental Concepts

## Definition: Graph

A graph  $G$  is an ordered pair of disjoint sets,  $(V, E)$  such that  $E$  is a subset of the set of unordered pairs of  $V$ .

### NOTES:

- Graphs are formal mathematical structures that model pairwise relations among objects from a particular set.

# Fundamental Concepts

## Definition: Graph

A graph  $G$  is an ordered pair of disjoint sets,  $(V, E)$  such that  $E$  is a subset of the set of unordered pairs of  $V$ .

### NOTES:

- Graphs are formal mathematical structures that model pairwise relations among objects from a particular set.
- If  $G$  is a graph, then  $E = E(G)$  is the edge set of  $G$  and  $V = V(G)$  is the vertex set of  $G$ .

# Fundamental Concepts

## Definition: Graph

A graph  $G$  is an ordered pair of disjoint sets,  $(V, E)$  such that  $E$  is a subset of the set of unordered pairs of  $V$ .

## NOTES:

- Graphs are formal mathematical structures that model pairwise relations among objects from a particular set.
- If  $G$  is a graph, then  $E = E(G)$  is the edge set of  $G$  and  $V = V(G)$  is the vertex set of  $G$ .
- Clearly,  $E \subseteq V \times V$ .

# Fundamental Concepts

## Definition: Graph

A graph  $G$  is an ordered pair of disjoint sets,  $(V, E)$  such that  $E$  is a subset of the set of unordered pairs of  $V$ .

## NOTES:



- Graphs are formal mathematical structures that model pairwise relations among objects from a particular set.
- If  $G$  is a graph, then  $E = E(G)$  is the edge set of  $G$  and  $V = V(G)$  is the vertex set of  $G$ .
- Clearly,  $E \subseteq V \times V$ .
- Edges connect pairs of vertices.



# Fundamental Concepts

## Definition: Digraph (directed graph)

A graph  $D$  with ordered pairs of nodes called arcs.



# Fundamental Concepts


## **Definition: Simple Graph**

A graph with no multiple edges or loops.


# Fundamental Concepts

## Definition: Multigraph


A graph with multiple edges (two nodes connected by more than one edge).



*simple graph*




*nonsimple graph  
with multiple edges*



*nonsimple graph  
with loops*


# The Königsberg Bridge Problem

- This is a multigraph.




# The Königsberg Bridge Problem

- This is a multigraph.
- **FACT:** A path that uses all edges in a multigraph exactly once is called an **Eulerian path**.



# The Königsberg Bridge Problem


- This is a multigraph.
- **FACT:** A path that uses all edges in a multigraph exactly once is called an **Eulerian path**.
- **Theorem:** A multigraph exhibiting a path between any pair of vertices has an Eulerian path *iff* there are only 0 or 2 vertices that have an odd number of edges leaving them.






# The Königsberg Bridge Problem

- This is a multigraph.
- **FACT:** A path that uses all edges in a multigraph exactly once is called an **Eulerian path**.
- **Theorem:** A multigraph exhibiting a path between any pair of vertices has an Eulerian path *iff* there are only 0 or 2 vertices that have an odd number of edges leaving them.
- The multigraph to the right has 4 vertices with an odd number of edges leaving them, so it has no Eulerian path.



# The Königsberg Bridge Problem

- This is a multigraph.
- **FACT:** A path that uses all edges in a multigraph exactly once is called an **Eulerian path**.
- **Theorem:** A multigraph exhibiting a path between any pair of vertices has an Eulerian path *iff* there are only 0 or 2 vertices that have an odd number of edges leaving them.
- The multigraph to the right has 4 vertices with an odd number of edges leaving them, so it has no Eulerian path.



# A Related Problem

## Theorem: The Handshake Theorem

If  $G$  is any undirected graph, then the sum of the degrees of all the  $n$  vertices,  $v$ , of  $G$  equals twice the number of edges,  $m$ , of  $G$ . That is,

$$\sum_{v \in V(G)} \deg(v) = 2m$$

# A Related Problem

## Proof: The Handshake Theorem

- **Crux:** Each edge has two end vertices.

# A Related Problem

## **Proof: The Handshake Theorem**

- Let  $G$  be an arbitrarily chosen undirected graph.

# A Related Problem

## **Proof: The Handshake Theorem**

- Let  $G$  be an arbitrarily chosen undirected graph.
- Trivially true if there are no vertices (no edges).



# A Related Problem

## Proof: The Handshake Theorem

- Let  $G$  be an arbitrarily chosen undirected graph.
- Trivially true if there are no vertices (no edges).
- If  $G$  has  $n$  vertices,  $v_1, v_2, \dots, v_n$ , and  $m$  edges, where  $n \in \mathbf{Z}^+$ ,  $m \in \mathbf{N}$ , we claim that each edge of  $G$  contributes 2 to the total degree of  $G$ .

# A Related Problem

## Proof: The Handshake Theorem

- Let  $G$  be an arbitrarily chosen undirected graph.
- Trivially true if there are no vertices (no edges).
- If  $G$  has  $n$  vertices,  $v_1, v_2, \dots, v_n$ , and  $m$  edges, where  $n \in \mathbf{Z}^+$ ,  $m \in \mathbf{N}$ , we claim that each edge of  $G$  contributes 2 to the total degree of  $G$ .
- For suppose  $e$  is an arbitrarily chosen edge with endpoints  $v_i$  and  $v_j$ .

# A Related Problem

## Proof: The Handshake Theorem

- Let  $G$  be an arbitrarily chosen undirected graph.
- Trivially true if there are no vertices (no edges).
- If  $G$  has  $n$  vertices,  $v_1, v_2, \dots, v_n$ , and  $m$  edges, where  $n \in \mathbf{Z}^+$ ,  $m \in \mathbf{N}$ , we claim that each edge of  $G$  contributes 2 to the total degree of  $G$ .
- For suppose  $e$  is an arbitrarily chosen edge with endpoints  $v_i$  and  $v_j$ .
- This edge contributes 1 to the degree of  $v_i$  and 1 to the degree of  $v_j$ .

# A Related Problem

## Proof: The Handshake Theorem

- Let  $G$  be an arbitrarily chosen undirected graph.
- Trivially true if there are no vertices (no edges).
- If  $G$  has  $n$  vertices,  $v_1, v_2, \dots, v_n$ , and  $m$  edges, where  $n \in \mathbf{Z}^+$ ,  $m \in \mathbf{N}$ , we claim that each edge of  $G$  contributes 2 to the total degree of  $G$ .
- For suppose  $e$  is an arbitrarily chosen edge with endpoints  $v_i$  and  $v_j$ .
- This edge contributes 1 to the degree of  $v_i$  and 1 to the degree of  $v_j$ .
- This is true even if  $v_i = v_j$  (an edge that is a loop is counted twice in computing the degree on which it is incident).

# A Related Problem

## Proof: The Handshake Theorem

- Let  $G$  be an arbitrarily chosen undirected graph.
- Trivially true if there are no vertices (no edges).
- If  $G$  has  $n$  vertices,  $v_1, v_2, \dots, v_n$ , and  $m$  edges, where  $n \in \mathbf{Z}^+$ ,  $m \in \mathbf{N}$ , we claim that each edge of  $G$  contributes 2 to the total degree of  $G$ .
- For suppose  $e$  is an arbitrarily chosen edge with endpoints  $v_i$  and  $v_j$ .
- This edge contributes 1 to the degree of  $v_i$  and 1 to the degree of  $v_j$ .
- This is true even if  $v_i = v_j$  (an edge that is a loop is counted twice in computing the degree on which it is incident).
- Therefore,  $e$  contributes 2 to the total degree of  $G$ .

# A Related Problem


## Proof: The Handshake Theorem

- Let  $G$  be an arbitrarily chosen undirected graph.
- Trivially true if there are no vertices (no edges).
- If  $G$  has  $n$  vertices,  $v_1, v_2, \dots, v_n$ , and  $m$  edges, where  $n \in \mathbf{Z}^+$ ,  $m \in \mathbf{N}$ , we claim that each edge of  $G$  contributes 2 to the total degree of  $G$ .
- For suppose  $e$  is an arbitrarily chosen edge with endpoints  $v_i$  and  $v_j$ .
- This edge contributes 1 to the degree of  $v_i$  and 1 to the degree of  $v_j$ .
- This is true even if  $v_i = v_j$  (an edge that is a loop is counted twice in computing the degree on which it is incident).
- Therefore,  $e$  contributes 2 to the total degree of  $G$ .
- Since  $e$  was arbitrarily chosen, this shows that each edge of  $G$  contributes 2 to the total degree of  $G$ .

**QED**

# Example

- Apply the Handshake Theorem to the following:



$$\# \text{ edges} = 9$$

✓ OK

$$\sum \text{deg} = 18$$

# Corollary

- **Theorem:** The total degree of an undirected graph is even.



# Corollary

- **Theorem:** The total degree of an undirected graph is even.
  - Known as The Handshaking “Lemma”.

# Corollary

- **Theorem:** The total degree of an undirected graph is even.
  - Known as The Handshaking “Lemma”.
  - Actually, “The number of vertices of odd degree is even”.

# Corollary

- **Theorem:** The total degree of an undirected graph is even.
  - Known as The Handshaking “Lemma”.
  - Actually, “The number of vertices of odd degree is even”.
  - Example: Is the graph with 4 vertices of degrees 1, 1, 2, and 3 possible?

NO!



# Presentation Terminated