



MONIKA R 2024-CSE ▾

M2

Started on Wednesday, 8 October 2025, 3:27 PM

State Finished

Completed on Wednesday, 8 October 2025, 4:05 PM

Time taken 38 mins 26 secs

Marks 1.00/1.00

Grade 4.00 out of 4.00 (100%)

Question 1 | Correct | Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that $A[j] - A[i] = k$, $i \neq j$.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as $5 - 1 = 4$

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer: (penalty regime: 0 %)

```

1  #include <stdio.h>
2
3  int hasPairWithDifference(int arr[], int n, int k) {
4      int i = 0, j = 1;
5
6      while (i < n && j < n) {
7          if (i != j) {
8              int diff = arr[j] - arr[i];
9              if (diff == k)
10                 return 1;
11                 else if (diff < k)
12                     j++;
13                 else
14                     i++;
15             } else {
16                 j++;
17             }
18         }
19
20         return 0;
21     }
22
23     int main() {
24         int n, k;
25         scanf("%d", &n);
26
27         int arr[n];
28         for (int i = 0; i < n; i++)
29             scanf("%d", &arr[i]);
30
31         scanf("%d", &k);
32
33         printf("%d\n", hasPairWithDifference(arr, n, k));
34         return 0;
35     }
36

```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

[Back to Course](#)