

# Chapter 6 part 1

## Jointly Distributed Random Variables

Jingchen (Monika) Hu

Vassar College

MATH 241

# Outline

- 1 Joint distribution
- 2 Independent random variables

# Outline

- 1 Joint distribution
- 2 Independent random variables

# Joint cdf

## Definition

We have a pair of random variables (either discrete or continuous)  $X$  and  $Y$ . The *joint cumulative probability distribution function* of  $X$  and  $Y$  is defined by


$$F_{X,Y}(x, y) = P[X \leq x, Y \leq y]$$

# Joint cdf

## Definition

We have a pair of random variables (either discrete or continuous)  $X$  and  $Y$ . The *joint cumulative probability distribution function* of  $X$  and  $Y$  is defined by

$$\begin{aligned} F_{X,Y}(x,y) &= P[X \leq x, Y \leq y] \\ &= P[(X,Y) \text{ lies south-west of the point } (x,y)] \end{aligned}$$



# Properties of joint cdf

- For one random variable: marginal cdf

$$F_X(x) = F_{X,Y}(x, \infty)$$

$$F_X(x) = P(X \leq x) = P(X \leq x, Y \leq \infty) = F_{X,Y}(x, \infty)$$

# Properties of joint cdf

- For one random variable: marginal cdf

$$F_X(x) = F_{X,Y}(x, \infty)$$

$$F_X(x) = P(X \leq x) = P(X \leq x, Y \leq \infty) = F_{X,Y}(x, \infty)$$

$$F_Y(y) = F_{X,Y}(\infty, y)$$

$$F_Y(y) = P(Y \leq y) = P(X \leq \infty, Y \leq y) = F_{X,Y}(\infty, y)$$

# Properties of joint cdf

- For one random variable: marginal cdf

$$F_X(x) = F_{X,Y}(x, \infty)$$

$$F_X(x) = P(X \leq x) = P(X \leq x, Y \leq \infty) = F_{X,Y}(x, \infty)$$

$$F_Y(y) = F_{X,Y}(\infty, y)$$

$$F_Y(y) = P(Y \leq y) = P(X \leq \infty, Y \leq y) = F_{X,Y}(\infty, y)$$

- Joint probabilities

$$P(X > x, Y > y) = 1 - F_X(x) - F_Y(y) + F_{X,Y}(x, y)$$



## Example: two discrete random variables

Draw two socks at random, without replacement, from a drawer full of twelve colored socks:

6 black, 4 white, 2 purple

Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

Then the distributions of  $B$  and  $W$  are given by:

## Example: two discrete random variables

Draw two socks at random, without replacement, from a drawer full of twelve colored socks:

6 black, 4 white, 2 purple

Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

Then the distributions of  $B$  and  $W$  are given by:

	0	1	2
$P(B=k)$			
$P(W=k)$			

## Example: two discrete random variables

Draw two socks at random, without replacement, from a drawer full of twelve colored socks:

6 black, 4 white, 2 purple

Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

Then the distributions of  $B$  and  $W$  are given by:

	0	1	2
$P(B=k)$	$\frac{6}{12} \cdot \frac{5}{11} = \frac{15}{66}$		
$P(W=k)$			

## Example: two discrete random variables

Draw two socks at random, without replacement, from a drawer full of twelve colored socks:

6 black, 4 white, 2 purple

Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

Then the distributions of  $B$  and  $W$  are given by:

	0	1	2
$P(B=k)$	$\frac{6}{12} \cdot \frac{5}{11} = \frac{15}{66}$	$2 \cdot \frac{6}{12} \cdot \frac{6}{11} = \frac{36}{66}$	
$P(W=k)$			

## Example: two discrete random variables

Draw two socks at random, without replacement, from a drawer full of twelve colored socks:

6 black, 4 white, 2 purple

Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

Then the distributions of  $B$  and  $W$  are given by:

	0	1	2
$P(B=k)$	$\frac{6}{12} \cdot \frac{5}{11} = \frac{15}{66}$	$2 \cdot \frac{6}{12} \cdot \frac{6}{11} = \frac{36}{66}$	$\frac{6}{12} \cdot \frac{5}{11} = \frac{15}{66}$
$P(W=k)$			

## Example: two discrete random variables

Draw two socks at random, without replacement, from a drawer full of twelve colored socks:

6 black, 4 white, 2 purple

Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

Then the distributions of  $B$  and  $W$  are given by:

	0	1	2
$P(B=k)$	$\frac{6}{12} \cdot \frac{5}{11} = \frac{15}{66}$	$2 \cdot \frac{6}{12} \cdot \frac{6}{11} = \frac{36}{66}$	$\frac{6}{12} \cdot \frac{5}{11} = \frac{15}{66}$
$P(W=k)$	$\frac{8}{12} \cdot \frac{7}{11} = \frac{28}{66}$		

## Example: two discrete random variables

Draw two socks at random, without replacement, from a drawer full of twelve colored socks:

6 black, 4 white, 2 purple

Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

Then the distributions of  $B$  and  $W$  are given by:

	0	1	2
$P(B=k)$	$\frac{6}{12} \cdot \frac{5}{11} = \frac{15}{66}$	$2 \cdot \frac{6}{12} \cdot \frac{6}{11} = \frac{36}{66}$	$\frac{6}{12} \cdot \frac{5}{11} = \frac{15}{66}$
$P(W=k)$	$\frac{8}{12} \cdot \frac{7}{11} = \frac{28}{66}$	$2 \cdot \frac{4}{12} \cdot \frac{8}{11} = \frac{32}{66}$	

## Example: two discrete random variables

Draw two socks at random, without replacement, from a drawer full of twelve colored socks:

6 black, 4 white, 2 purple

Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

Then the distributions of  $B$  and  $W$  are given by:

	0	1	2
$P(B=k)$	$\frac{6}{12} \cdot \frac{5}{11} = \frac{15}{66}$	$2 \cdot \frac{6}{12} \cdot \frac{6}{11} = \frac{36}{66}$	$\frac{6}{12} \cdot \frac{5}{11} = \frac{15}{66}$
$P(W=k)$	$\frac{8}{12} \cdot \frac{7}{11} = \frac{28}{66}$	$2 \cdot \frac{4}{12} \cdot \frac{8}{11} = \frac{32}{66}$	$\frac{4}{12} \cdot \frac{3}{11} = \frac{6}{66}$



## Example: two discrete random variables

Draw two socks at random, without replacement, from a drawer full of twelve colored socks:

6 black, 4 white, 2 purple

Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

Then the distributions of  $B$  and  $W$  are given by:

	0	1	2
$P(B=k)$	$\frac{6}{12} \cdot \frac{5}{11} = \frac{15}{66}$	$2 \cdot \frac{6}{12} \cdot \frac{6}{11} = \frac{36}{66}$	$\frac{6}{12} \cdot \frac{5}{11} = \frac{15}{66}$
$P(W=k)$	$\frac{8}{12} \cdot \frac{7}{11} = \frac{28}{66}$	$2 \cdot \frac{4}{12} \cdot \frac{8}{11} = \frac{32}{66}$	$\frac{4}{12} \cdot \frac{3}{11} = \frac{6}{66}$

Note -  $P(B = k) = \frac{\binom{6}{k} \binom{6}{2-k}}{\binom{12}{2}}$  and  $P(W = k) = \frac{\binom{4}{k} \binom{8}{2-k}}{\binom{12}{2}}$

Draw two socks at random, without replacement, from a drawer full of twelve colored socks: 6 black, 4 white, 2 purple. Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

The *joint distribution* is given by:  $p_{B,W}(b, w) = P(B = b, W = w)$

		W			
		0	1	2	
B	0				$P(B = b, W = w) = \left\{ \right.$
	1				
	2				

Draw two socks at random, without replacement, from a drawer full of twelve colored socks: 6 black, 4 white, 2 purple. Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

The *joint distribution* is given by:  $p_{B,W}(b, w) = P(B = b, W = w)$

		W			
		0	1	2	
B	0	$\frac{1}{66}$			
	1				
	2				

$$P(B = b, W = w) = \begin{cases} 1/66 & \text{if } b=0, w=0 \\ 0 & \text{otherwise} \end{cases}$$

Draw two socks at random, without replacement, from a drawer full of twelve colored socks: 6 black, 4 white, 2 purple. Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

The *joint distribution* is given by:  $p_{B,W}(b, w) = P(B = b, W = w)$

		W			
		0	1	2	
B	0	$\frac{1}{66}$	$\frac{8}{66}$		
	1				
	2				

$$P(B = b, W = w) = \begin{cases} 1/66 & \text{If } b=0, w=0 \\ 8/66 & \text{If } b=0, w=1 \end{cases}$$

Draw two socks at random, without replacement, from a drawer full of twelve colored socks: 6 black, 4 white, 2 purple. Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

The *joint distribution* is given by:  $p_{B,W}(b, w) = P(B = b, W = w)$

		W			
		0	1	2	
B	0	$\frac{1}{66}$	$\frac{8}{66}$	$\frac{6}{66}$	
	1				
	2				

$$P(B = b, W = w) = \begin{cases} 1/66 & \text{If } b=0, w=0 \\ 8/66 & \text{If } b=0, w=1 \\ 6/66 & \text{If } b=0, w=2 \end{cases}$$

Draw two socks at random, without replacement, from a drawer full of twelve colored socks: 6 black, 4 white, 2 purple. Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

The *joint distribution* is given by:  $p_{B,W}(b, w) = P(B = b, W = w)$

		W			
		0	1	2	
B	0	$\frac{1}{66}$	$\frac{8}{66}$	$\frac{6}{66}$	
	1	$\frac{12}{66}$			
	2				

$$P(B = b, W = w) = \begin{cases} 1/66 & \text{If } b=0, w=0 \\ 8/66 & \text{If } b=0, w=1 \\ 6/66 & \text{If } b=0, w=2 \\ 12/66 & \text{If } b=1, w=0 \end{cases}$$

Draw two socks at random, without replacement, from a drawer full of twelve colored socks: 6 black, 4 white, 2 purple. Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

The *joint distribution* is given by:  $p_{B,W}(b, w) = P(B = b, W = w)$

		W			
		0	1	2	
B	0	$\frac{1}{66}$	$\frac{8}{66}$	$\frac{6}{66}$	
	1	$\frac{12}{66}$	$\frac{24}{66}$		
	2				

$$P(B = b, W = w) = \begin{cases} 1/66 & \text{If } b=0, w=0 \\ 8/66 & \text{If } b=0, w=1 \\ 6/66 & \text{If } b=0, w=2 \\ 12/66 & \text{If } b=1, w=0 \\ 24/66 & \text{If } b=1, w=1 \end{cases}$$

Draw two socks at random, without replacement, from a drawer full of twelve colored socks: 6 black, 4 white, 2 purple. Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

The *joint distribution* is given by:  $p_{B,W}(b, w) = P(B = b, W = w)$

		W			
		0	1	2	
B	0	$\frac{1}{66}$	$\frac{8}{66}$	$\frac{6}{66}$	
	1	$\frac{12}{66}$	$\frac{24}{66}$	0	
	2				

$$P(B = b, W = w) = \begin{cases} 1/66 & \text{If } b=0, w=0 \\ 8/66 & \text{If } b=0, w=1 \\ 6/66 & \text{If } b=0, w=2 \\ 12/66 & \text{If } b=1, w=0 \\ 24/66 & \text{If } b=1, w=1 \end{cases}$$



Draw two socks at random, without replacement, from a drawer full of twelve colored socks: 6 black, 4 white, 2 purple. Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

The *joint distribution* is given by:  $p_{B,W}(b, w) = P(B = b, W = w)$

		W			
		0	1	2	
B	0	$\frac{1}{66}$	$\frac{8}{66}$	$\frac{6}{66}$	
	1	$\frac{12}{66}$	$\frac{24}{66}$	0	
	2	$\frac{15}{66}$			

$$P(B = b, W = w) = \begin{cases} 1/66 & \text{If } b=0, w=0 \\ 8/66 & \text{If } b=0, w=1 \\ 6/66 & \text{If } b=0, w=2 \\ 12/66 & \text{If } b=1, w=0 \\ 24/66 & \text{If } b=1, w=1 \\ 15/66 & \text{If } b=2, w=0 \end{cases}$$

Draw two socks at random, without replacement, from a drawer full of twelve colored socks: 6 black, 4 white, 2 purple. Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

The *joint distribution* is given by:  $p_{B,W}(b, w) = P(B = b, W = w)$

		W			
		0	1	2	
B	0	$\frac{1}{66}$	$\frac{8}{66}$	$\frac{6}{66}$	
	1	$\frac{12}{66}$	$\frac{24}{66}$	0	
	2	$\frac{15}{66}$	0	0	

$$P(B = b, W = w) = \begin{cases} 1/66 & \text{If } b=0, w=0 \\ 8/66 & \text{If } b=0, w=1 \\ 6/66 & \text{If } b=0, w=2 \\ 12/66 & \text{If } b=1, w=0 \\ 24/66 & \text{If } b=1, w=1 \\ 15/66 & \text{If } b=2, w=0 \end{cases}$$

Draw two socks at random, without replacement, from a drawer full of twelve colored socks: 6 black, 4 white, 2 purple. Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

The *joint distribution* is given by:  $p_{B,W}(b, w) = P(B = b, W = w)$

		W			$P(B = b, W = w) = \begin{cases} 1/66 & \text{If } b=0, w=0 \\ 8/66 & \text{If } b=0, w=1 \\ 6/66 & \text{If } b=0, w=2 \\ 12/66 & \text{If } b=1, w=0 \\ 24/66 & \text{If } b=1, w=1 \\ 15/66 & \text{If } b=2, w=0 \end{cases}$
		0	1	2	
B	0	$\frac{1}{66}$	$\frac{8}{66}$	$\frac{6}{66}$	
	1	$\frac{12}{66}$	$\frac{24}{66}$	0	
	2	$\frac{15}{66}$	0	0	

$$P(B = b, W = w) = \frac{\binom{6}{b} \binom{4}{w} \binom{2}{2-b-w}}{\binom{12}{2}}, \text{ for } 0 \leq b, w \leq 2 \text{ and } b + w \leq 2$$

Draw two socks at random, without replacement, from a drawer full of twelve colored socks: 6 black, 4 white, 2 purple. Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

The *joint distribution* is given by:  $p_{B,W}(b, w) = P(B = b, W = w)$

		W			$P(B = b, W = w) = \begin{cases} 1/66 & \text{If } b=0, w=0 \\ 8/66 & \text{If } b=0, w=1 \\ 6/66 & \text{If } b=0, w=2 \\ 12/66 & \text{If } b=1, w=0 \\ 24/66 & \text{If } b=1, w=1 \\ 15/66 & \text{If } b=2, w=0 \end{cases}$
		0	1	2	
B	0	$\frac{1}{66}$	$\frac{8}{66}$	$\frac{6}{66}$	
	1	$\frac{12}{66}$	$\frac{24}{66}$	0	
	2	$\frac{15}{66}$	0	0	

$$P(B = b, W = w) = \frac{\binom{6}{b} \binom{4}{w} \binom{2}{2-b-w}}{\binom{12}{2}}, \text{ for } 0 \leq b, w \leq 2 \text{ and } b + w \leq 2$$

Draw two socks at random, without replacement, from a drawer full of twelve colored socks: 6 black, 4 white, 2 purple. Let  $B$  be the number of Black socks,  $W$  the number of White socks drawn.

The *joint distribution* is given by:  $p_{B,W}(b, w) = P(B = b, W = w)$

		W			
		0	1	2	
B	0	$\frac{1}{66}$	$\frac{8}{66}$	$\frac{6}{66}$	
	1	$\frac{12}{66}$	$\frac{24}{66}$	0	
	2	$\frac{15}{66}$	0	0	
		$\frac{28}{66}$	$\frac{32}{66}$	$\frac{6}{66}$	

$$P(B = b, W = w) = \begin{cases} 1/66 & \text{If } b=0, w=0 \\ 8/66 & \text{If } b=0, w=1 \\ 6/66 & \text{If } b=0, w=2 \\ 12/66 & \text{If } b=1, w=0 \\ 24/66 & \text{If } b=1, w=1 \\ 15/66 & \text{If } b=2, w=0 \end{cases}$$

$$P(B = b, W = w) = \frac{\binom{6}{b} \binom{4}{w} \binom{2}{2-b-w}}{\binom{12}{2}}, \text{ for } 0 \leq b, w \leq 2 \text{ and } b + w \leq 2$$

## Marginal Distributions

Note that the column and row sums are the distributions of  $B$  and  $W$  respectively.

$$P(B = b) = P(B = b, W = 0) + P(B = b, W = 1) + P(B = b, W = 2)$$

$$P(W = w) = P(B = 0, W = w) + P(B = 1, W = w) + P(B = 2, W = w)$$

## Marginal Distributions

Note that the column and row sums are the distributions of  $B$  and  $W$  respectively.

$$P(B = b) = P(B = b, W = 0) + P(B = b, W = 1) + P(B = b, W = 2)$$

$$P(W = w) = P(B = 0, W = w) + P(B = 1, W = w) + P(B = 2, W = w)$$

These are the marginal distributions of  $B$  and  $W$ . In general,

$$P(X = x) = \sum_y P(X = x, Y = y) = \sum_y P(X = x \mid Y = y)P(Y = y)$$

## Conditional Distribution

Conditional distributions are defined as we have seen previously with

$$P(X = x \mid Y = y) = \frac{P(X = x, Y = y)}{P(Y = y)} = \frac{\text{joint pmf}}{\text{marginal pmf}}$$



# Joint distribution of two continuous random variables

## Definition

Random variables  $X$  and  $Y$  are *jointly continuous* if there exists a function  $f(x, y)$  such that

❶ Non-negative  $f(x, y) \geq 0$ , for any  $x, y \in \mathbb{R}$ , and

❷  $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = 1$ .

$f_{X,Y}(x, y)$  is called the *joint probability density function* of  $X$  and  $Y$ .

# Joint distribution of two continuous random variables

## Definition

Random variables  $X$  and  $Y$  are *jointly continuous* if there exists a function  $f(x, y)$  such that

- ① Non-negative  $f(x, y) \geq 0$ , for any  $x, y \in \mathbb{R}$ , and
- ②  $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = 1$ .

$f_{X,Y}(x, y)$  is called the *joint probability density function* of  $X$  and  $Y$ .

- For any set  $C \subset \mathbb{R}^2$ ,

$$P[(X, Y) \in C] = \iint_{(x,y) \in C} f(x, y) \, dx \, dy$$

# Joint distribution of two continuous random variables

## Definition

Random variables  $X$  and  $Y$  are *jointly continuous* if there exists a function  $f(x, y)$  such that

- ① Non-negative  $f(x, y) \geq 0$ , for any  $x, y \in \mathbb{R}$ , and
- ②  $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = 1$ .

$f_{X,Y}(x, y)$  is called the *joint probability density function* of  $X$  and  $Y$ .

- For any set  $C \subset \mathbb{R}^2$ ,

$$P[(X, Y) \in C] = \iint_{(x,y) \in C} f(x, y) \, dx \, dy$$

- Connection between joint pdf and joint cdf

$$F(a, b) = P(X \leq a, Y \leq b) = \int_{-\infty}^b \int_{-\infty}^a f(x, y) \, dx \, dy$$

# Joint distribution of two continuous random variables

## Definition

Random variables  $X$  and  $Y$  are *jointly continuous* if there exists a function  $f(x, y)$  such that

- ❶ Non-negative  $f(x, y) \geq 0$ , for any  $x, y \in \mathbb{R}$ , and
- ❷  $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = 1$ .

$f_{X,Y}(x, y)$  is called the *joint probability density function* of  $X$  and  $Y$ .

- For any set  $C \subset \mathbb{R}^2$ ,

$$P[(X, Y) \in C] = \iint_{(x,y) \in C} f(x, y) \, dx \, dy$$

- Connection between joint pdf and joint cdf

$$F(a, b) = P(X \leq a, Y \leq b) = \int_{-\infty}^b \int_{-\infty}^a f(x, y) \, dx \, dy$$

$$f(x, y) = \frac{\partial^2}{\partial x \partial y} F(x, y)$$


## Marginal pdfs

Marginal probability density functions are defined in terms of “integrating out” one of the random variables.

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) \, dy$$

$$f_Y(y) = \int_{-\infty}^{\infty} f(x, y) \, dx$$


Let  $X$  and  $Y$  be drawn uniformly from the triangle below. Find the joint pdf  $f_{X,Y}(x,y)$ .



Let  $X$  and  $Y$  be drawn uniformly from the triangle below. Find the joint pdf  $f_{X,Y}(x,y)$ .

Since the joint density is constant, then

$$f(x,y) = \begin{cases} c & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$




Let  $X$  and  $Y$  be drawn uniformly from the triangle below. Find the joint pdf  $f_{X,Y}(x,y)$ .

Since the joint density is constant, then

$$f(x,y) = \begin{cases} c & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Because

$$1 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x,y) \, dx \, dy$$






Let  $X$  and  $Y$  be drawn uniformly from the triangle below. Find the joint pdf  $f_{X,Y}(x,y)$ .

Since the joint density is constant, then

$$f(x,y) = \begin{cases} c & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Because

$$\begin{aligned} 1 &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x,y) \, dx \, dy \\ &= \iint_{x \geq 0, y \geq 0, x+y \leq 3} c \, dx \, dy \end{aligned}$$



Let  $X$  and  $Y$  be drawn uniformly from the triangle below. Find the joint pdf  $f_{X,Y}(x,y)$ .


Since the joint density is constant, then

$$f(x,y) = \begin{cases} c & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Because

$$\begin{aligned} 1 &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x,y) \, dx \, dy \\ &= \iint_{x \geq 0, y \geq 0, x+y \leq 3} c \, dx \, dy \end{aligned}$$

$$= c \times \text{area of the triangle} = c \times \frac{3 \times 3}{2}$$



Let  $X$  and  $Y$  be drawn uniformly from the triangle below. Find the joint pdf  $f_{X,Y}(x,y)$ .

Since the joint density is constant, then


$$f(x,y) = \begin{cases} c & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Because

$$\begin{aligned} 1 &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x,y) \, dx \, dy \\ &= \iint_{x \geq 0, y \geq 0, x+y \leq 3} c \, dx \, dy \end{aligned}$$

$$= c \times \text{area of the triangle} = c \times \frac{3 \times 3}{2}$$

Therefore,  $c = \frac{2}{9}$ .



## Recap

Joint cdf of two random variables  $X$  and  $Y$ :

$$F_{X,Y}(x,y) = P[X \leq x, Y \leq y], -\infty < x, y < \infty$$

- Probability of  $(X, Y)$  in a rectangle

$$\begin{aligned} &P(x_1 < X \leq x_2, y_1 < Y \leq y_2) \\ &= F(x_2, y_2) + F(x_1, y_1) - F(x_1, y_2) - F(x_2, y_1) \end{aligned}$$

- Marginal cdfs

$$F_X(x) = F_{X,Y}(x, \infty), \quad F_Y(y) = F_{X,Y}(\infty, y)$$

## Joint distribution of two discrete random variables

- Joint pmf

$$p_{X,Y}(x,y) = P(X = x, Y = y)$$

- Marginal pmfs

$$p_X(x) = \sum_{y:p(x,y)>0} p_{X,Y}(x,y), \quad p_Y(y) = \sum_{x:p(x,y)>0} p_{X,Y}(x,y)$$

## Joint distribution of two discrete random variables

- Joint pmf

$$p_{X,Y}(x,y) = P(X = x, Y = y)$$

- Marginal pmfs

$$p_X(x) = \sum_{y:p(x,y)>0} p_{X,Y}(x,y), \quad p_Y(y) = \sum_{x:p(x,y)>0} p_{X,Y}(x,y)$$

## Joint distribution of two continuous random variables

- Joint pdf

- ▶ Non-negative  $f_{X,Y}(x,y) \geq 0$ , for any  $x, y \in \mathbb{R}$
- ▶  $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x,y) \, dx \, dy = 1$
- ▶ For any set  $C \subset \mathbb{R}^2$ ,

$$P[(X,Y) \in C] = \iint_{(x,y) \in C} f_{X,Y}(x,y) \, dx \, dy$$


- Marginal pdfs

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x,y) \, dy, \quad f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x,y) \, dx$$

Let  $X$  and  $Y$  have the following joint pdf

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Find the marginal pdf  $f_X(x)$ .




Let  $X$  and  $Y$  have the following joint pdf

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Find the marginal pdf  $f_X(x)$ .

For  $x \in [0, 3]$ ,






Let  $X$  and  $Y$  have the following joint pdf

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Find the marginal pdf  $f_X(x)$ .

For  $x \in [0, 3]$ ,

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy$$




Let  $X$  and  $Y$  have the following joint pdf

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Find the marginal pdf  $f_X(x)$ .

For  $x \in [0, 3]$ ,

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy = \int_0^{3-x} \frac{2}{9} dy$$



Let  $X$  and  $Y$  have the following joint pdf


$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Find the marginal pdf  $f_X(x)$ .

For  $x \in [0, 3]$ ,

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy = \int_0^{3-x} \frac{2}{9} dy$$

*Be careful about the range of  $Y$  given  $X = x$ .*



Let  $X$  and  $Y$  have the following joint pdf


$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Find the marginal pdf  $f_X(x)$ .

For  $x \in [0, 3]$ ,

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) \, dy = \int_0^{3-x} \frac{2}{9} \, dy = \frac{2}{9}(3 - x)$$

*Be careful about the range of  $Y$  given  $X = x$ .*



Let  $X$  and  $Y$  have the following joint pdf

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$


Find the marginal pdf  $f_X(x)$ .

For  $x \in [0, 3]$ ,

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy = \int_0^{3-x} \frac{2}{9} dy = \frac{2}{9}(3 - x)$$


*Be careful about the range of  $Y$  given  $X = x$ .*

$$f_X(x) = \begin{cases} \frac{2}{9}(3 - x) & \text{for } x \in [0, 3] \\ 0 & \text{otherwise} \end{cases}$$



In the previous example, find  $P(X < Y)$ .

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$




In the previous example, find  $P(X < Y)$ .

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Identify the region

$$C = \{(x, y) : x \geq 0, y \geq 0 \text{ and } x + y \leq 3, x < y\}.$$




In the previous example, find  $P(X < Y)$ .

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Identify the region

$$C = \{(x, y) : x \geq 0, y \geq 0 \text{ and } x + y \leq 3, x < y\}.$$



$$P(X < Y) = \iint_{(x,y) \in C} f(x, y) \, dx \, dy$$




In the previous example, find  $P(X < Y)$ .

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Identify the region

$$C = \{(x, y) : x \geq 0, y \geq 0 \text{ and } x + y \leq 3, x < y\}.$$




$$\begin{aligned} P(X < Y) &= \iint_{(x,y) \in C} f(x, y) \, dx \, dy \\ &= \int_0^{\frac{3}{2}} \left[ \int_x^{3-x} \frac{2}{9} \, dy \right] dx \end{aligned}$$

In the previous example, find  $P(X < Y)$ .

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Identify the region

$$C = \{(x, y) : x \geq 0, y \geq 0 \text{ and } x + y \leq 3, x < y\}.$$




$$\begin{aligned} P(X < Y) &= \iint_{(x,y) \in C} f(x, y) \, dx \, dy \\ &= \int_0^{\frac{3}{2}} \left[ \int_x^{3-x} \frac{2}{9} \, dy \right] dx \\ &= \int_0^{\frac{3}{2}} \frac{2}{9} (3 - 2x) \, dx \end{aligned}$$

In the previous example, find  $P(X < Y)$ .

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Identify the region

$$C = \{(x, y) : x \geq 0, y \geq 0 \text{ and } x + y \leq 3, x < y\}.$$




$$\begin{aligned} P(X < Y) &= \iint_{(x,y) \in C} f(x, y) \, dx \, dy \\ &= \int_0^{\frac{3}{2}} \left[ \int_x^{3-x} \frac{2}{9} \, dy \right] dx \\ &= \int_0^{\frac{3}{2}} \frac{2}{9} (3 - 2x) \, dx = \frac{2}{9} \times \left[ 3x - x^2 \right]_0^{\frac{3}{2}} \end{aligned}$$

In the previous example, find  $P(X < Y)$ .

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Identify the region

$$C = \{(x, y) : x \geq 0, y \geq 0 \text{ and } x + y \leq 3, x < y\}.$$




$$\begin{aligned} P(X < Y) &= \iint_{(x,y) \in C} f(x, y) \, dx \, dy \\ &= \int_0^{\frac{3}{2}} \left[ \int_x^{3-x} \frac{2}{9} \, dy \right] dx \\ &= \int_0^{\frac{3}{2}} \frac{2}{9} (3 - 2x) \, dx = \frac{2}{9} \times \left[ 3x - x^2 \right]_0^{\frac{3}{2}} \\ &= \frac{2}{9} \left( \frac{9}{2} - \frac{9}{4} \right) \end{aligned}$$

In the previous example, find  $P(X < Y)$ .

$$f(x, y) = \begin{cases} \frac{2}{9} & \text{for } x \geq 0, y \geq 0 \text{ and } x + y \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

Identify the region

$$C = \{(x, y) : x \geq 0, y \geq 0 \text{ and } x + y \leq 3, x < y\}.$$



$$\begin{aligned} P(X < Y) &= \iint_{(x,y) \in C} f(x, y) \, dx \, dy \\ &= \int_0^{\frac{3}{2}} \left[ \int_x^{3-x} \frac{2}{9} \, dy \right] dx \\ &= \int_0^{\frac{3}{2}} \frac{2}{9} (3 - 2x) \, dx = \frac{2}{9} \times \left[ 3x - x^2 \right]_0^{\frac{3}{2}} \\ &= \frac{2}{9} \left( \frac{9}{2} - \frac{9}{4} \right) = \frac{1}{2} \end{aligned}$$

## Joint distribution of two continuous random variables

### • Joint pdf

- ▶ Non-negative  $f_{X,Y}(x,y) \geq 0$ , for any  $x, y \in \mathbb{R}$
- ▶  $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x,y) \, dx \, dy = 1$
- ▶ For any set  $C \subset \mathbb{R}^2$ ,

$$P[(X,Y) \in C] = \iint_{(x,y) \in C} f_{X,Y}(x,y) \, dx \, dy$$

### • Between joint cdf and joint pdf

$$F_{X,Y}(a,b) = \int_{-\infty}^b \int_{-\infty}^a f_{X,Y}(x,y) \, dx \, dy$$

$$f_{X,Y}(x,y) = \frac{\partial^2}{\partial x \partial y} F_{X,Y}(x,y)$$

### • Marginal pdfs

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x,y) \, dy, \quad f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x,y) \, dx$$

# Outline

1 Joint distribution

2 Independent random variables

# Independent random variables

## Definition

Random variables  $X$  and  $Y$  are *independent* if any real sets  $A, B \subset \mathbb{R}$ ,

$$P(X \in A, Y \in B) = P(X \in A)P(Y \in B)$$



# Independent random variables

## Definition

Random variables  $X$  and  $Y$  are *independent* if any real sets  $A, B \subset \mathbb{R}$ ,

$$P(X \in A, Y \in B) = P(X \in A)P(Y \in B)$$

Random variables  $X$  and  $Y$  are independent **if and only if**

- Cdf: for any  $x, y \in \mathbb{R}$

$$F_{X,Y}(x, y) = F_X(x)F_Y(y)$$

# Independent random variables

## Definition

Random variables  $X$  and  $Y$  are *independent* if any real sets  $A, B \subset \mathbb{R}$ ,

$$P(X \in A, Y \in B) = P(X \in A)P(Y \in B)$$

Random variables  $X$  and  $Y$  are independent **if and only if**

- Cdf: for any  $x, y \in \mathbb{R}$

$$F_{X,Y}(x, y) = F_X(x)F_Y(y)$$

- If both are discrete, pmf: for any  $x, y \in \mathbb{R}$

$$p_{X,Y}(x, y) = p_X(x)p_Y(y)$$

# Independent random variables

## Definition

Random variables  $X$  and  $Y$  are *independent* if any real sets  $A, B \subset \mathbb{R}$ ,

$$P(X \in A, Y \in B) = P(X \in A)P(Y \in B)$$

Random variables  $X$  and  $Y$  are independent **if and only if**

- Cdf: for any  $x, y \in \mathbb{R}$

$$F_{X,Y}(x, y) = F_X(x)F_Y(y)$$

- If both are discrete, pmf: for any  $x, y \in \mathbb{R}$

$$p_{X,Y}(x, y) = p_X(x)p_Y(y)$$


- If both are continuous, pdf: for any  $x, y \in \mathbb{R}$

$$f_{X,Y}(x, y) = f_X(x)f_Y(y)$$

A man and a woman decide to meet at a certain location. If each of them independently arrives at a time uniformly distributed between 12 noon and 1 P.M., find the probability that the first to arrive has to wait longer than 10 minutes.

A man and a woman decide to meet at a certain location. If each of them independently arrives at a time uniformly distributed between 12 noon and 1 P.M., find the probability that the first to arrive has to wait longer than 10 minutes.


Let random variables  $X, Y$  be the time they arrive (uniform between 0 to 60 minutes).



A man and a woman decide to meet at a certain location. If each of them independently arrives at a time uniformly distributed between 12 noon and 1 P.M., find the probability that the first to arrive has to wait longer than 10 minutes.


Let random variables  $X, Y$  be the time they arrive (uniform between 0 to 60 minutes).

$$P(|X - Y| > 10)$$



A man and a woman decide to meet at a certain location. If each of them independently arrives at a time uniformly distributed between 12 noon and 1 P.M., find the probability that the first to arrive has to wait longer than 10 minutes.


Let random variables  $X, Y$  be the time they arrive (uniform between 0 to 60 minutes).



$$\begin{aligned}
 &P(|X - Y| > 10) \\
 &= P(Y > X + 10) + P(Y < X - 10)
 \end{aligned}$$

A man and a woman decide to meet at a certain location. If each of them independently arrives at a time uniformly distributed between 12 noon and 1 P.M., find the probability that the first to arrive has to wait longer than 10 minutes.

Let random variables  $X, Y$  be the time they arrive (uniform between 0 to 60 minutes).




$$\begin{aligned}
 &P(|X - Y| > 10) \\
 &= P(Y > X + 10) + P(Y < X - 10) \\
 &= 2P(Y > X + 10)
 \end{aligned}$$



A man and a woman decide to meet at a certain location. If each of them independently arrives at a time uniformly distributed between 12 noon and 1 P.M., find the probability that the first to arrive has to wait longer than 10 minutes.


Let random variables  $X, Y$  be the time they arrive (uniform between 0 to 60 minutes).



$$\begin{aligned}
 &P(|X - Y| > 10) \\
 &= P(Y > X + 10) + P(Y < X - 10) \\
 &= 2P(Y > X + 10) \\
 &= 2 \iint_{y > x+10} f_{X,Y}(x, y) \, dx \, dy
 \end{aligned}$$

A man and a woman decide to meet at a certain location. If each of them independently arrives at a time uniformly distributed between 12 noon and 1 P.M., find the probability that the first to arrive has to wait longer than 10 minutes.


Let random variables  $X, Y$  be the time they arrive (uniform between 0 to 60 minutes).



$$\begin{aligned}
 &P(|X - Y| > 10) \\
 &= P(Y > X + 10) + P(Y < X - 10) \\
 &= 2P(Y > X + 10) \\
 &= 2 \iint_{y > x + 10} f_{X,Y}(x, y) \, dx \, dy \\
 &= 2 \iint_{y > x + 10} f_X(x) f_Y(y) \, dx \, dy
 \end{aligned}$$

A man and a woman decide to meet at a certain location. If each of them independently arrives at a time uniformly distributed between 12 noon and 1 P.M., find the probability that the first to arrive has to wait longer than 10 minutes.


Let random variables  $X, Y$  be the time they arrive (uniform between 0 to 60 minutes).



$$\begin{aligned}
 &P(|X - Y| > 10) \\
 &= P(Y > X + 10) + P(Y < X - 10) \\
 &= 2P(Y > X + 10) \\
 &= 2 \iint_{y > x + 10} f_{X,Y}(x, y) \, dx \, dy \\
 &= 2 \iint_{y > x + 10} f_X(x) f_Y(y) \, dx \, dy \\
 &= 2 \int_{10}^{60} \int_0^{y-10} (1/60)^2 \, dx \, dy
 \end{aligned}$$

A man and a woman decide to meet at a certain location. If each of them independently arrives at a time uniformly distributed between 12 noon and 1 P.M., find the probability that the first to arrive has to wait longer than 10 minutes.

Let random variables  $X, Y$  be the time they arrive (uniform between 0 to 60 minutes).



$$\begin{aligned}
 &P(|X - Y| > 10) \\
 &= P(Y > X + 10) + P(Y < X - 10) \\
 &= 2P(Y > X + 10) \\
 &= 2 \iint_{y > x + 10} f_{X,Y}(x, y) \, dx \, dy \\
 &= 2 \iint_{y > x + 10} f_X(x) f_Y(y) \, dx \, dy \\
 &= 2 \int_{10}^{60} \int_0^{y-10} (1/60)^2 \, dx \, dy = 25/36
 \end{aligned}$$

# Independent random variables

The continuous (discrete) random variables  $X$  and  $Y$  are independent **if and only if** their joint probability density (mass) function can be expressed as

$$f_{X,Y}(x,y) = g(x)h(y), \quad -\infty < x, y < \infty$$

## More than two random variables

☞ Random variables  $X_1, X_2, \dots, X_n$  are *independent* if any real sets  $A_1, A_2, \dots, A_n \subset \mathbb{R}$ ,

$$P(X_1 \in A_1, \dots, X_n \in A_n) = P(X_1 \in A_1) \cdots P(X_n \in A_n)$$

## More than two random variables

☞ Random variables  $X_1, X_2, \dots, X_n$  are *independent* if any real sets  $A_1, A_2, \dots, A_n \subset \mathbb{R}$ ,

$$P(X_1 \in A_1, \dots, X_n \in A_n) = P(X_1 \in A_1) \cdots P(X_n \in A_n)$$

Random variables  $X_1, X_2, \dots, X_n$  are independent **if and only if**

- Cdf: for any  $x_1, x_2, \dots, x_n \in \mathbb{R}$

$$F(x_1, \dots, x_n) = F_{X_1}(x_1) \cdots F_{X_n}(x_n)$$

## More than two random variables

- ☞ Random variables  $X_1, X_2, \dots, X_n$  are *independent* if any real sets  $A_1, A_2, \dots, A_n \subset \mathbb{R}$ ,

$$P(X_1 \in A_1, \dots, X_n \in A_n) = P(X_1 \in A_1) \cdots P(X_n \in A_n)$$

Random variables  $X_1, X_2, \dots, X_n$  are independent **if and only if**

- Cdf: for any  $x_1, x_2, \dots, x_n \in \mathbb{R}$

$$F(x_1, \dots, x_n) = F_{X_1}(x_1) \cdots F_{X_n}(x_n)$$

- If both are discrete, pmf: for any  $x_1, x_2, \dots, x_n \in \mathbb{R}$

$$p(x_1, \dots, x_n) = p_{X_1}(x_1) \cdots p_{X_n}(x_n)$$



## More than two random variables

- ➡ Random variables  $X_1, X_2, \dots, X_n$  are *independent* if any real sets  $A_1, A_2, \dots, A_n \subset \mathbb{R}$ ,

$$P(X_1 \in A_1, \dots, X_n \in A_n) = P(X_1 \in A_1) \cdots P(X_n \in A_n)$$

Random variables  $X_1, X_2, \dots, X_n$  are independent **if and only if**

- Cdf: for any  $x_1, x_2, \dots, x_n \in \mathbb{R}$

$$F(x_1, \dots, x_n) = F_{X_1}(x_1) \cdots F_{X_n}(x_n)$$

- If both are discrete, pmf: for any  $x_1, x_2, \dots, x_n \in \mathbb{R}$

$$p(x_1, \dots, x_n) = p_{X_1}(x_1) \cdots p_{X_n}(x_n)$$

- If both are continuous, pdf: for any  $x_1, x_2, \dots, x_n \in \mathbb{R}$

$$f(x_1, \dots, x_n) = f_{X_1}(x_1) \cdots f_{X_n}(x_n)$$