

# Chapter 8

## Limit Theorems

Jingchen (Monika) Hu

Vassar College

MATH 241

# Outline

## 1 Central limit theorem

## The central limit theorem

Let  $X_1, X_2, \dots$  be a sequence of independent and identically distributed random variables, each having mean  $\mu$  and variance  $\sigma^2$ . Then the distribution of

$$\frac{X_1 + X_2 + \dots + X_n - n\mu}{\sigma\sqrt{n}}$$

tends to the standard normal as  $n \rightarrow \infty$ .

## The central limit theorem

Let  $X_1, X_2, \dots$  be a sequence of independent and identically distributed random variables, each having mean  $\mu$  and variance  $\sigma^2$ . Then the distribution of

$$\frac{X_1 + X_2 + \dots + X_n - n\mu}{\sigma\sqrt{n}}$$

tends to the standard normal as  $n \rightarrow \infty$ .

In words, if the random variables have a finite mean  $\mu$  and a finite variance  $\sigma^2$ , then the distribution of the sum of the first  $n$  of them is, for large  $n$ , approximately that of a normal random variable with mean  $n\mu$  and variance  $n\sigma^2$ .

# Applications of the CLT

A person has 100 light bulbs whose lifetimes are independent exponentials with mean 5 hours. If the bulbs are used one at a time, with a failed bulb being replaced immediately by a new one, approximate the probability that there is still a working bulb after 525 hours. Note that if  $X \sim \text{Exponential}(\lambda)$ , then  $E[X] = \frac{1}{\lambda}$  and  $\text{Var}(X) = \frac{1}{\lambda^2}$ .

# Applications of the CLT

A person has 100 light bulbs whose lifetimes are independent exponentials with mean 5 hours. If the bulbs are used one at a time, with a failed bulb being replaced immediately by a new one, approximate the probability that there is still a working bulb after 525 hours. Note that if  $X \sim \text{Exponential}(\lambda)$ , then  $E[X] = \frac{1}{\lambda}$  and  $\text{Var}(X) = \frac{1}{\lambda^2}$ .

- For each light bulb:  $\mu = 5, \sigma^2 = 25$  ( $\lambda = \frac{1}{5}$ )

# Applications of the CLT

A person has 100 light bulbs whose lifetimes are independent exponentials with mean 5 hours. If the bulbs are used one at a time, with a failed bulb being replaced immediately by a new one, approximate the probability that there is still a working bulb after 525 hours. Note that if  $X \sim \text{Exponential}(\lambda)$ , then  $E[X] = \frac{1}{\lambda}$  and  $\text{Var}(X) = \frac{1}{\lambda^2}$ .

- For each light bulb:  $\mu = 5, \sigma^2 = 25$  ( $\lambda = \frac{1}{5}$ )
- Using the CLT

$$P\left\{\sum_{i=1}^{100} X_i > 525\right\}$$

# Applications of the CLT

A person has 100 light bulbs whose lifetimes are independent exponentials with mean 5 hours. If the bulbs are used one at a time, with a failed bulb being replaced immediately by a new one, approximate the probability that there is still a working bulb after 525 hours. Note that if  $X \sim \text{Exponential}(\lambda)$ , then  $E[X] = \frac{1}{\lambda}$  and  $\text{Var}(X) = \frac{1}{\lambda^2}$ .

- For each light bulb:  $\mu = 5, \sigma^2 = 25$  ( $\lambda = \frac{1}{5}$ )
- Using the CLT

$$\begin{aligned} P\left\{\sum_{i=1}^{100} X_i > 525\right\} &\approx P\left\{N(0, 1) > \frac{525 - 500}{\sqrt{100 \times 25}}\right\} = P\{N(0, 1) > .5\} \\ &= .3085 \end{aligned}$$



# Applications of the CLT

A person has 100 light bulbs whose lifetimes are independent exponentials with mean 5 hours. If the bulbs are used one at a time, with a failed bulb being replaced immediately by a new one, approximate the probability that there is still a working bulb after 525 hours. Note that if  $X \sim \text{Exponential}(\lambda)$ , then  $E[X] = \frac{1}{\lambda}$  and  $\text{Var}(X) = \frac{1}{\lambda^2}$ .

- For each light bulb:  $\mu = 5, \sigma^2 = 25$  ( $\lambda = \frac{1}{5}$ )
- Using the CLT

$$\begin{aligned} P\left\{\sum_{i=1}^{100} X_i > 525\right\} &\approx P\left\{N(0, 1) > \frac{525 - 500}{\sqrt{100 \times 25}}\right\} = P\{N(0, 1) > .5\} \\ &= .3085 \end{aligned}$$

- Note that because exponential distribution is also continuous, we do not apply the continuity correction (.5); for discrete distributions (e.g. binomial, poisson), the continuity correction is necessary