

Chapter 4 part 2

Discrete Random Variables

Jingchen (Monika) Hu

Vassar College

MATH 241

Outline

1 Variance

2 Bernoulli distribution and Binomial distribution

Variance

- Expected value (or mean) $\mu = E[X]$ yields the weighted average of the possible values of X

Variance

- Expected value (or mean) $\mu = E[X]$ yields the weighted average of the possible values of X

Definition

Variance measures the variation (or spread) of these values

$$\sigma^2 = \text{Var}(X) = E[(X - E(X))^2] = E[(X - \mu)^2]$$

This holds for all random variable X (not necessary discrete random variable).

Variance

- Expected value (or mean) $\mu = E[X]$ yields the weighted average of the possible values of X

Definition

Variance measures the variation (or spread) of these values

$$\sigma^2 = \text{Var}(X) = E[(X - E(X))^2] = E[(X - \mu)^2]$$

This holds for all random variable X (not necessary discrete random variable).

- One common simplification:

$$\text{Var}(X) = E(X^2) - \mu^2$$

Standard deviation

Definition

Standard Deviation is the square root of the variance

$$\sigma = SD(X) = \sqrt{Var(X)}$$

Standard deviation

Definition

Standard Deviation is the square root of the variance

$$\sigma = SD(X) = \sqrt{Var(X)}$$

Example: compute the standard deviation of a fair 4-sided die toss.

Standard deviation

Definition

Standard Deviation is the square root of the variance

$$\sigma = SD(X) = \sqrt{Var(X)}$$

Example: compute the standard deviation of a fair 4-sided die toss.

$$Var(X) = E[X^2] - (E[X])^2$$

Standard deviation

Definition

Standard Deviation is the square root of the variance

$$\sigma = SD(X) = \sqrt{Var(X)}$$

Example: compute the standard deviation of a fair 4-sided die toss.

$$Var(X) = E[X^2] - (E[X])^2$$

$$E[X] = 1 \times \frac{1}{4} + 2 \times \frac{1}{4} + 3 \times \frac{1}{4} + 4 \times \frac{1}{4} = 2.5$$

Standard deviation

Definition

Standard Deviation is the square root of the variance

$$\sigma = SD(X) = \sqrt{Var(X)}$$

Example: compute the standard deviation of a fair 4-sided die toss.

$$Var(X) = E[X^2] - (E[X])^2$$

$$E[X] = 1 \times \frac{1}{4} + 2 \times \frac{1}{4} + 3 \times \frac{1}{4} + 4 \times \frac{1}{4} = 2.5$$

$$E[X^2] = 1 \times \frac{1}{4} + 4 \times \frac{1}{4} + 9 \times \frac{1}{4} + 16 \times \frac{1}{4} = 7.5$$

Standard deviation

Definition

Standard Deviation is the square root of the variance

$$\sigma = SD(X) = \sqrt{Var(X)}$$

Example: compute the standard deviation of a fair 4-sided die toss.

$$Var(X) = E[X^2] - (E[X])^2$$

$$E[X] = 1 \times \frac{1}{4} + 2 \times \frac{1}{4} + 3 \times \frac{1}{4} + 4 \times \frac{1}{4} = 2.5$$

$$E[X^2] = 1 \times \frac{1}{4} + 4 \times \frac{1}{4} + 9 \times \frac{1}{4} + 16 \times \frac{1}{4} = 7.5$$

$$Var(X) = 7.5 - 2.5^2 = 1.25$$

Standard deviation

Definition

Standard Deviation is the square root of the variance

$$\sigma = SD(X) = \sqrt{Var(X)}$$

Example: compute the standard deviation of a fair 4-sided die toss.

$$Var(X) = E[X^2] - (E[X])^2$$

$$E[X] = 1 \times \frac{1}{4} + 2 \times \frac{1}{4} + 3 \times \frac{1}{4} + 4 \times \frac{1}{4} = 2.5$$

$$E[X^2] = 1 \times \frac{1}{4} + 4 \times \frac{1}{4} + 9 \times \frac{1}{4} + 16 \times \frac{1}{4} = 7.5$$

$$Var(X) = 7.5 - 2.5^2 = 1.25$$

$$SD(X) = \sqrt{Var[X]} = 1.12$$

Variance measures the spread of X

X_1 is a discrete random variable with pmf

x	-1	1
$p_{X_1}(x)$	$1/2$	$1/2$

X_2 is a discrete random variable with pmf

x	-2	2
$p_{X_2}(x)$	$1/2$	$1/2$

Variance measures the spread of X

X_1 is a discrete random variable with pmf X_2 is a discrete random variable with pmf

x	-1	1
$p_{X_1}(x)$	$1/2$	$1/2$

x	-2	2
$p_{X_2}(x)$	$1/2$	$1/2$

$$\mu_1 = 0, \sigma_1^2 = E[X_1^2] - 0^2 = 1, \sigma_1 = 1$$

Variance measures the spread of X

X_1 is a discrete random variable with pmf X_2 is a discrete random variable with pmf

x	-1	1
$p_{X_1}(x)$	$1/2$	$1/2$

x	-2	2
$p_{X_2}(x)$	$1/2$	$1/2$

$$\mu_1 = 0, \sigma_1^2 = E[X_1^2] - 0^2 = 1, \sigma_1 = 1$$

$$\mu_2 = 0, \sigma_2^2 = E[X_2^2] - 0^2 = 4, \sigma_2 = 2$$

Variance measures the spread of X

X_1 is a discrete random variable with pmf X_2 is a discrete random variable with pmf


x	-1	1
$p_{X_1}(x)$	1/2	1/2

x	-2	2
$p_{X_2}(x)$	1/2	1/2


$$\mu_1 = 0, \sigma_1^2 = E[X_1^2] - 0^2 = 1, \sigma_1 = 1$$

$$\mu_2 = 0, \sigma_2^2 = E[X_2^2] - 0^2 = 4, \sigma_2 = 2$$

Increasing variance (or sd) reflects increasing spread.



Distributions with $SD = 1$



Property of variance

We know that $\text{Var}(X) = E[(X - \mu)^2]$, and for constants a, b ,

$$E[aX + b] = aE[X] + b.$$

Write $\text{Var}(aX + b)$ as a function of $\text{Var}(X)$.

Property of variance

We know that $Var(X) = E[(X - \mu)^2]$, and for constants a, b ,

$$E[aX + b] = aE[X] + b.$$

Write $Var(aX + b)$ as a function of $Var(X)$.

$$Var(aX + b) = E[(aX + b - E[aX + b])^2]$$

Property of variance

We know that $Var(X) = E[(X - \mu)^2]$, and for constants a, b ,

$$E[aX + b] = aE[X] + b.$$

Write $Var(aX + b)$ as a function of $Var(X)$.

$$\begin{aligned} Var(aX + b) &= E[(aX + b - E[aX + b])^2] \\ &= E[(aX + b - aE[X] - b)^2] \\ &= E[(aX - a\mu)^2] \\ &= E[a^2(X - \mu)^2] \\ &= a^2 Var(X) \end{aligned}$$

Properties of variance

$$\text{Var}(X) \geq 0$$

- $\text{Var}(X) = 0$ if and only if X is a constant.

Properties of variance

$$\text{Var}(X) \geq 0$$

- $\text{Var}(X) = 0$ if and only if X is a constant.

If a and b are constants, then

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Properties of variance

$$\text{Var}(X) \geq 0$$

- $\text{Var}(X) = 0$ if and only if X is a constant.

If a and b are constants, then

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

-

$$\text{Var}(aX) = a^2 \text{Var}(X)$$

-

$$\text{Var}(X + b) = \text{Var}(X)$$

-

$$\text{Var}(b) = 0$$

Outline

1 Variance

2 Bernoulli distribution and Binomial distribution

Bernoulli distribution

A trial has two outcomes: success (1) or failure (0).

Let random variable X be the number of success in a single trial.

Bernoulli distribution

A trial has two outcomes: success (1) or failure (0).

Let random variable X be the number of success in a single trial.

Definition

Random variable X has a *Bernoulli distribution*, if

$$P(X = 1) = p, \quad P(X = 0) = 1 - p$$

where $0 \leq p \leq 1$ is the probability of a success.

Bernoulli distribution

A trial has two outcomes: success (1) or failure (0).

Let random variable X be the number of success in a single trial.

Definition

Random variable X has a *Bernoulli distribution*, if

$$P(X = 1) = p, \quad P(X = 0) = 1 - p$$

where $0 \leq p \leq 1$ is the probability of a success.

- The pmf of Bernoulli distribution

$$X \sim \text{Ber}(p) \iff p(1) = p, \quad p(0) = 1 - p$$

Bernoulli distribution

A trial has two outcomes: success (1) or failure (0).

Let random variable X be the number of success in a single trial.

Definition

Random variable X has a *Bernoulli distribution*, if

$$P(X = 1) = p, \quad P(X = 0) = 1 - p$$

where $0 \leq p \leq 1$ is the probability of a success.

- The pmf of Bernoulli distribution

$$X \sim \text{Ber}(p) \iff p(1) = p, \quad p(0) = 1 - p$$

- Found by Swiss mathematician Jacob Bernoulli.



Examples of Bernoulli distributions

- Toss a fair coin and obtain a head. $p = 0.5$.
- Roll a fair 6-sided die and obtain a 6. $p = 1/6$.
- Earn an A for this class. $p \in [0, 1]$.

Examples of Bernoulli distributions

- Toss a fair coin and obtain a head. $p = 0.5$.
- Roll a fair 6-sided die and obtain a 6. $p = 1/6$.
- Earn an A for this class. $p \in [0, 1]$.

Find the mean and variance of $X \sim \text{Ber}(p)$.

Examples of Bernoulli distributions

- Toss a fair coin and obtain a head. $p = 0.5$.
- Roll a fair 6-sided die and obtain a 6. $p = 1/6$.
- Earn an A for this class. $p \in [0, 1]$.

Find the mean and variance of $X \sim \text{Ber}(p)$.

$$E(X) = 1 \times p + 0 \times (1 - p) = p$$

$$\text{Var}(X) = E(X^2) - (E[X])^2 = 1^2 \times p + 0^2 \times (1 - p) - p^2 = p - p^2 = p(1 - p)$$

Recap

Variance σ^2

- **For all random variable**

$$Var(X) = E[(X - \mu)^2] = E[X^2] - \mu^2$$

- **Linear function**

$$Var(aX + b) = a^2 Var(X)$$

- **Constants**

$$Var(c) = 0$$

Standard deviation

$$SD(X) = \sqrt{Var(X)}$$

Bernoulli distribution

$$p(1) = p, \quad p(0) = 1 - p$$

$$\mu = p, \quad \sigma^2 = p(1 - p)$$

Binomial distribution

Definition

Define X to be the number of successes in a fixed number n of independent trials with the same probability of success p as having a *Binomial distribution*.

Binomial distribution

Definition

Define X to be the number of successes in a fixed number n of independent trials with the same probability of success p as having a *Binomial distribution*.

Then the pmf of X is $P(X = k) = P(\text{getting } k \text{ successes in } n \text{ trials})$

Binomial distribution

Definition

Define X to be the number of successes in a fixed number n of independent trials with the same probability of success p as having a *Binomial distribution*.

Then the pmf of X is $P(X = k) = P(\text{getting } k \text{ successes in } n \text{ trials})$

$$X \sim \text{Bin}(n, p) \iff p(k) = \binom{n}{k} p^k (1-p)^{n-k}, \quad k = 0, 1, \dots, n$$

Binomial distribution

Definition

Define X to be the number of successes in a fixed number n of independent trials with the same probability of success p as having a *Binomial distribution*.

Then the pmf of X is $P(X = k) = P(\text{getting } k \text{ successes in } n \text{ trials})$

$$X \sim \text{Bin}(n, p) \iff p(k) = \binom{n}{k} p^k (1-p)^{n-k}, \quad k = 0, 1, \dots, n$$

Example of Binomial distributions

- The number of heads obtained by tossing a fair coin 10 times.
 $n = 10, p = 0.5$.

Binomial distribution

Definition

Define X to be the number of successes in a fixed number n of independent trials with the same probability of success p as having a *Binomial distribution*.

Then the pmf of X is $P(X = k) = P(\text{getting } k \text{ successes in } n \text{ trials})$

$$X \sim \text{Bin}(n, p) \iff p(k) = \binom{n}{k} p^k (1-p)^{n-k}, \quad k = 0, 1, \dots, n$$

Example of Binomial distributions

- The number of heads obtained by tossing a fair coin 10 times.
 $n = 10, p = 0.5$.
- The number of 6 obtained when roll four fair 6-sided dice simultaneously. $n = 4, p = 1/6$.

Binomial distribution

Definition

Define X to be the number of successes in a fixed number n of independent trials with the same probability of success p as having a *Binomial distribution*.

Then the pmf of X is $P(X = k) = P(\text{getting } k \text{ successes in } n \text{ trials})$

$$X \sim \text{Bin}(n, p) \iff p(k) = \binom{n}{k} p^k (1-p)^{n-k}, \quad k = 0, 1, \dots, n$$

Example of Binomial distributions

- The number of heads obtained by tossing a fair coin 10 times.
 $n = 10, p = 0.5$.
- The number of 6 obtained when roll four fair 6-sided dice simultaneously. $n = 4, p = 1/6$.
- The number of A's students will earn in this semester.
 $n = \text{number of classes you're taking}, p \text{ varies by classes...}$
 \implies not really a Binomial random variable!

Example: let X be the number of 6 obtained when roll four fair 6-sided dice simultaneously. Find its pmf.

Example: let X be the number of 6 obtained when roll four fair 6-sided dice simultaneously. Find its pmf.

$$n = 4, p = 1/6,$$

$$p(k) = \binom{n}{k} p^k (1 - p)^{n-k}$$

Example: let X be the number of 6 obtained when roll four fair 6-sided dice simultaneously. Find its pmf.

$$n = 4, p = 1/6,$$

$$p(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

$$p(0) = \binom{4}{0} (1/6)^0 (5/6)^4 = 0.4823$$

Example: let X be the number of 6 obtained when roll four fair 6-sided dice simultaneously. Find its pmf.

$$n = 4, p = 1/6,$$

$$p(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

$$p(0) = \binom{4}{0} (1/6)^0 (5/6)^4 = 0.4823$$

$$p(1) = \binom{4}{1} (1/6)^1 (5/6)^3 = 0.3858$$

Example: let X be the number of 6 obtained when roll four fair 6-sided dice simultaneously. Find its pmf.

$$n = 4, p = 1/6,$$

$$p(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

$$p(0) = \binom{4}{0} (1/6)^0 (5/6)^4 = 0.4823$$

$$p(1) = \binom{4}{1} (1/6)^1 (5/6)^3 = 0.3858$$

$$p(2) = \binom{4}{2} (1/6)^2 (5/6)^2 = 0.1157$$

Example: let X be the number of 6 obtained when roll four fair 6-sided dice simultaneously. Find its pmf.

$$n = 4, p = 1/6,$$

$$p(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

$$p(0) = \binom{4}{0} (1/6)^0 (5/6)^4 = 0.4823$$

$$p(1) = \binom{4}{1} (1/6)^1 (5/6)^3 = 0.3858$$

$$p(2) = \binom{4}{2} (1/6)^2 (5/6)^2 = 0.1157$$

$$p(3) = \binom{4}{3} (1/6)^3 (5/6)^1 = 0.0154$$

Example: let X be the number of 6 obtained when roll four fair 6-sided dice simultaneously. Find its pmf.

$$n = 4, p = 1/6,$$

$$p(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

$$p(0) = \binom{4}{0} (1/6)^0 (5/6)^4 = 0.4823$$

$$p(1) = \binom{4}{1} (1/6)^1 (5/6)^3 = 0.3858$$

$$p(2) = \binom{4}{2} (1/6)^2 (5/6)^2 = 0.1157$$


$$p(3) = \binom{4}{3} (1/6)^3 (5/6)^1 = 0.0154$$

$$p(4) = \binom{4}{4} (1/6)^4 (5/6)^0 = 0.0008$$

Example: let X be the number of 6 obtained when roll four fair 6-sided dice simultaneously. Find its pmf.

$$n = 4, p = 1/6,$$

$$p(k) = \binom{n}{k} p^k (1-p)^{n-k}$$



$$p(0) = \binom{4}{0} (1/6)^0 (5/6)^4 = 0.4823$$

$$p(1) = \binom{4}{1} (1/6)^1 (5/6)^3 = 0.3858$$

$$p(2) = \binom{4}{2} (1/6)^2 (5/6)^2 = 0.1157$$

$$p(3) = \binom{4}{3} (1/6)^3 (5/6)^1 = 0.0154$$

$$p(4) = \binom{4}{4} (1/6)^4 (5/6)^0 = 0.0008$$

Suppose 42% of the students in a class have part time jobs. Among a random sample of 10 students in that class, what is the probability that exactly 8 have part time jobs?

Suppose 42% of the students in a class have part time jobs. Among a random sample of 10 students in that class, what is the probability that exactly 8 have part time jobs?

Binomial distribution. $n = 10, p = 0.42$.

$$P(X = 8) = \binom{10}{8} \times 0.42^8 \times 0.58^2 = 0.0147$$

Suppose 42% of the students in a class have part time jobs. Among a random sample of 10 students in that class, what is the probability that exactly 8 have part time jobs?

Binomial distribution. $n = 10, p = 0.42$.

$$P(X = 8) = \binom{10}{8} \times 0.42^8 \times 0.58^2 = 0.0147$$

We can use the binomial distribution to calculate the probability of k successes in n trials, as long as

Suppose 42% of the students in a class have part time jobs. Among a random sample of 10 students in that class, what is the probability that exactly 8 have part time jobs?

Binomial distribution. $n = 10, p = 0.42$.

$$P(X = 8) = \binom{10}{8} \times 0.42^8 \times 0.58^2 = 0.0147$$

We can use the binomial distribution to calculate the probability of k successes in n trials, as long as

- 1 the trials are independent

Suppose 42% of the students in a class have part time jobs. Among a random sample of 10 students in that class, what is the probability that exactly 8 have part time jobs?

Binomial distribution. $n = 10, p = 0.42$.

$$P(X = 8) = \binom{10}{8} \times 0.42^8 \times 0.58^2 = 0.0147$$

We can use the binomial distribution to calculate the probability of k successes in n trials, as long as

- 1 the trials are independent
- 2 the number of trials, n , is fixed

Suppose 42% of the students in a class have part time jobs. Among a random sample of 10 students in that class, what is the probability that exactly 8 have part time jobs?

Binomial distribution. $n = 10, p = 0.42$.

$$P(X = 8) = \binom{10}{8} \times 0.42^8 \times 0.58^2 = 0.0147$$

We can use the binomial distribution to calculate the probability of k successes in n trials, as long as

- 1 the trials are independent
- 2 the number of trials, n , is fixed
- 3 each trial outcome can be classified as a *success* or a *failure*

Suppose 42% of the students in a class have part time jobs. Among a random sample of 10 students in that class, what is the probability that exactly 8 have part time jobs?

Binomial distribution. $n = 10, p = 0.42$.


$$P(X = 8) = \binom{10}{8} \times 0.42^8 \times 0.58^2 = 0.0147$$

We can use the binomial distribution to calculate the probability of k successes in n trials, as long as


- 1 the trials are independent
- 2 the number of trials, n , is fixed
- 3 each trial outcome can be classified as a *success* or a *failure*
- 4 the probability of success, p , is the same for each trial

Pmf of Binomial distribution: uni-modal


pmf: Bin(4, 1/6)




pmf: Bin(20, 1/6)



pmf: Bin(4, 1/2)



pmf: Bin(20, 1/2)



Binomial pmf is valid (or well-defined)

- Positive: for any $x \in \mathbb{R}$

$$p(x) \geq 0$$

- Total one (**required**):

$$\sum_{k=0}^n p(k) = 1$$

Binomial pmf is valid (or well-defined)

- Positive: for any $x \in \mathbb{R}$

$$p(x) \geq 0$$

- Total one (**required**):

$$\sum_{k=0}^n p(k) = 1$$

Recall the Binomial Theorem $(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}$

$$\sum_{k=0}^n p(k) = \sum_{k=0}^n \binom{n}{k} p^k (1-p)^{n-k} = [p + (1-p)]^n = 1^n = 1$$

Mean of Binomial random variable

Toss a coin n times, each toss has prob p being a head. On average, total number of heads equals np .

Mean of Binomial random variable

Toss a coin n times, each toss has prob p being a head. On average, total number of heads equals np .

$$E[X] = np$$

Check textbook page 131 for the derivation (**not required**).

Properties of Binomial distribution

- Variance

$$\text{Var}[X] = np(1 - p)$$

Check textbook page 132 for the derivation (**not required**).

Properties of Binomial distribution

- Variance

$$\text{Var}[X] = np(1 - p)$$

Check textbook page 132 for the derivation (**not required**).

- If we have independent Bernoulli random variable's X_1, X_2, \dots, X_n with the same probability of success p , then their sum has a Binomial distribution

$$X = X_1 + X_2 + \dots + X_n \sim \text{Bin}(n, p)$$

Roll 4 fair six-sided dice. Let X be the number of 6 obtained. Find its mean and variance.

Roll 4 fair six-sided dice. Let X be the number of 6 obtained. Find its mean and variance.

$$X \sim \text{Bin}(n = 4, p = 1/6)$$

Roll 4 fair six-sided dice. Let X be the number of 6 obtained. Find its mean and variance.

$$X \sim \text{Bin}(n = 4, p = 1/6)$$

$$E[X] = np = 4 \times (1/6) = 2/3$$

$$\text{Var}(X) = np(1 - p) = 4 \times (1/6) \times (5/6) = 5/9$$

Recap

Binomial distribution $X \sim \text{Bin}(n, p)$

$$p(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

- mean $\mu = np$
- variance $\sigma^2 = np(1-p)$