

# Chapter 3

## Conditional Probability and Independence

Jingchen (Monika) Hu

Vassar College

MATH 241

# Outline

- 1 Conditional probability
- 2 Bayes theorem
- 3 Independent events
- 4 Perceptions of probability and biases

## Conditional probability: motivation

The probability of getting a one when rolling a fair 6-sided die is  $1/6$

## Conditional probability: motivation

The probability of getting a one when rolling a fair 6-sided die is  $1/6$

Suppose you were given the extra information that the die roll was an odd number (hence 1, 3 or 5)

## Conditional probability: motivation

The probability of getting a one when rolling a fair 6-sided die is  $1/6$

Suppose you were given the extra information that the die roll was an odd number (hence 1, 3 or 5)

*Conditional on this new information*, the probability of a one is now  $1/3$

# Conditional probability

## Definition

Given two events  $A$  and  $B$  with  $P(B) > 0$ , the *conditional probability* of  $A$  given  $B$  has occurred is defined as

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)}$$

# Conditional probability

## Definition

Given two events  $A$  and  $B$  with  $P(B) > 0$ , the *conditional probability* of  $A$  given  $B$  has occurred is defined as

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)}$$

- When  $B$  is the sample space:  $P(A \mid B) = P(A)$
- Intuition: in a sample space with equally likely outcomes,

$$P(A \mid B) = \frac{\#(A \cap B)}{\#(B)}$$

## Example

Consider the die roll example:  $B = \{1 \text{ or } 3 \text{ or } 5\}$ ,  $A = \{1\}$

$$\begin{aligned} P(\text{get 1 given that roll is odd}) &= P(A \mid B) \\ &= \frac{P(A \cap B)}{P(B)} \\ &= \frac{P(\text{get 1})}{P(\text{get 1 or 3 or 5})} \\ &= \frac{1/6}{3/6} = \frac{1}{3} \end{aligned}$$



Example: roll two fair 6-sided dice. Find the probability of the sum of two rolls is 7, given the condition that the first roll is 4.

Example: roll two fair 6-sided dice. Find the probability of the sum of two rolls is 7, given the condition that the first roll is 4.

Denote two events  $A = \{\text{Sum is 7}\}$  and  $B = \{\text{First roll is 4}\}$ .

Find  $P(A \mid B)$ . Formulas to use?

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)}$$

Example: roll two fair 6-sided dice. Find the probability of the sum of two rolls is 7, given the condition that the first roll is 4.

Denote two events  $A = \{\text{Sum is 7}\}$  and  $B = \{\text{First roll is 4}\}$ .  
Find  $P(A \mid B)$ . Formulas to use?

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)}$$

In context,

$$P(A \cap B) = P(\{(4, 3)\}) = 1/36, \quad P(B) = 1/6$$

$$P(A \mid B) = \frac{1/36}{1/6} = \frac{1}{6}$$

# Propositions of conditional probability

①  $P(A \mid A) = 1$

# Propositions of conditional probability

$$\textcircled{1} \quad P(A \mid A) = 1$$

$$\textcircled{2} \quad P(A^c \mid A) = 0$$

# Propositions of conditional probability

$$\textcircled{1} \quad P(A \mid A) = 1$$

$$\textcircled{2} \quad P(A^c \mid A) = 0$$

$$\textcircled{3} \quad P(A^c \mid B) = 1 - P(A \mid B)$$

# Multiplication rule

- 👉 By the definition of conditional probability, the *joint probability* of  $A$  and  $B$  is

$$P(A \cap B) = P(A \mid B)P(B)$$

- Usually,  $P(A)$  and  $P(B)$  are called *marginal probabilities*.

# Multiplication rule

- 👉 By the definition of conditional probability, the *joint probability* of  $A$  and  $B$  is

$$P(A \cap B) = P(A \mid B)P(B)$$

- Usually,  $P(A)$  and  $P(B)$  are called *marginal probabilities*.

- 👉 Generalize to  $n$  events: chaining of probabilities

$$P\left(\bigcap_{i=1}^n A_i\right) = P(A_1)P(A_2 \mid A_1)P(A_3 \mid A_1, A_2) \cdots P(A_n \mid A_1, \dots, A_{n-1})$$



A box contains 8 white balls and 4 black ones. 3 balls are drawn at random without replacement.

- 1 What's the probability of getting  $W_1B_2B_3$ ?
- 2 What's the probability of getting black in draw 2?

A box contains 8 white balls and 4 black ones. 3 balls are drawn at random without replacement.

- 1 What's the probability of getting  $W_1B_2B_3$ ?
- 2 What's the probability of getting black in draw 2?

1 Multiplication rule

$$\begin{aligned}P(W_1B_2B_3) &= P(W_1)P(B_2 | W_1)P(B_3 | W_1B_2) \\&= \frac{8}{12} \cdot \frac{4}{11} \cdot \frac{3}{10}\end{aligned}$$

A box contains 8 white balls and 4 black ones. 3 balls are drawn at random without replacement.

- ➊ What's the probability of getting  $W_1B_2B_3$ ?
- ➋ What's the probability of getting black in draw 2?

- ➊ Multiplication rule

$$\begin{aligned} P(W_1B_2B_3) &= P(W_1)P(B_2 | W_1)P(B_3 | W_1B_2) \\ &= \frac{8}{12} \cdot \frac{4}{11} \cdot \frac{3}{10} \end{aligned}$$

- ➋ If we want to consider draw 1, which can be  $W_1$  or  $B_1$  (disjoint!)

$$\begin{aligned} P(B_2) &= P(W_1B_2) + P(B_1B_2) \\ &= \frac{8}{12} \cdot \frac{4}{11} + \frac{4}{12} \cdot \frac{3}{11} = \frac{44}{132} = \frac{4}{12} \end{aligned}$$

## Law of total probability

For events  $A_1, \dots, A_n$  are **disjoint**, and

$$\bigcup_{i=1}^n A_i = S,$$

then for any event  $B$ ,

## Law of total probability

For events  $A_1, \dots, A_n$  are **disjoint**, and

$$\bigcup_{i=1}^n A_i = S,$$

then for any event  $B$ ,

👉 law of total probability

$$P(B) = P(B \mid A_1)P(A_1) + \dots + P(B \mid A_n)P(A_n)$$

# Law of total probability

For events  $A_1, \dots, A_n$  are **disjoint**, and

$$\bigcup_{i=1}^n A_i = S,$$

then for any event  $B$ ,

👉 law of total probability

$$P(B) = P(B \mid A_1)P(A_1) + \dots + P(B \mid A_n)P(A_n)$$

- Such collection of sets  $A_1, \dots, A_n$  is also called a partition of sample space.

Example: a firm is considering a drug-testing program for its employees, but before it begins it wants to know the scope of the problem if any exists. Realizing the sensitivity of this issue, the personnel director decides to use a randomized response survey. It is believed that respondents are more likely to be honest when such forms are used. Each employee is asked to flip a fair coin. If the coin comes up heads, answer the question “Do you carpool to work?”. If the coin comes up tails, answer the question “Have you used illegal drugs within the last month?”. Assume that all employees answer the survey honestly. Out of 8000 responses, 1420 answered “YES”. Suppose the firm knows that 35% of its employees carpool to work. What is the probability that an employee chosen at random used illegal drugs within the last month?

Example: a firm is considering a drug-testing program for its employees, but before it begins it wants to know the scope of the problem if any exists. Realizing the sensitivity of this issue, the personnel director decides to use a randomized response survey. It is believed that respondents are more likely to be honest when such forms are used. Each employee is asked to flip a fair coin. If the coin comes up heads, answer the question “Do you carpool to work?”. If the coin comes up tails, answer the question “Have you used illegal drugs within the last month?”. Assume that all employees answer the survey honestly. Out of 8000 responses, 1420 answered “YES”. Suppose the firm knows that 35% of its employees carpool to work. What is the probability that an employee chosen at random used illegal drugs within the last month?

Let events  $A_1 = \{\text{head}\}$ ,  $A_2 = \{\text{tail}\}$ ,  $B = \{\text{answer YES}\}$ .

Since  $\{A_1, A_2\}$  is a partition, so we can use the law to total probability



Example: a firm is considering a drug-testing program for its employees, but before it begins it wants to know the scope of the problem if any exists. Realizing the sensitivity of this issue, the personnel director decides to use a randomized response survey. It is believed that respondents are more likely to be honest when such forms are used. Each employee is asked to flip a fair coin. If the coin comes up heads, answer the question “Do you carpool to work?”. If the coin comes up tails, answer the question “Have you used illegal drugs within the last month?”. Assume that all employees answer the survey honestly. Out of 8000 responses, 1420 answered “YES”. Suppose the firm knows that 35% of its employees carpool to work. What is the probability that an employee chosen at random used illegal drugs within the last month?

Let events  $A_1 = \{\text{head}\}$ ,  $A_2 = \{\text{tail}\}$ ,  $B = \{\text{answer YES}\}$ .

Since  $\{A_1, A_2\}$  is a partition, so we can use the law to total probability

$$P(B) = P(B | A_1)P(A_1) + P(B | A_2)P(A_2)$$

$$P(\text{YES}) = P(\text{carpool})P(\text{head}) + P(\text{drug})P(\text{tail})$$

$$\frac{1420}{8000} = 0.35 \times 0.5 + P(\text{drug}) \times 0.5$$

$$P(\text{drug}) = 0.005$$

# Recap

- Marginal probability:  $P(A), P(B)$
- Joint probability:  $P(A \cap B)$
- Conditional probability

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)}$$

# Recap

- Marginal probability:  $P(A), P(B)$
- Joint probability:  $P(A \cap B)$
- Conditional probability

$$P(A | B) = \frac{P(A \cap B)}{P(B)}$$

- Multiplication rule:

$$P(A \cap B) = P(A | B) \times P(B)$$

- Law of total probability: for a partition  $\{A_1, A_2, \dots, A_n\}$  of  $S$ ,

$$P(B) = \sum_{j=1}^n P(B | A_j)P(A_j)$$

# Outline

- 1 Conditional probability
- 2 Bayes theorem**
- 3 Independent events
- 4 Perceptions of probability and biases

## Bayes theorem (also called Bayes rule)

Suppose events  $A_1, \dots, A_n$  are disjoint, and  $\bigcup_{i=1}^n A_i = S$ , with  $P(A_i) > 0$ ,  $i = 1, 2, \dots, n$ . Then for any event  $B$  with  $P(B) > 0$ ,

## Bayes theorem (also called Bayes rule)

Suppose events  $A_1, \dots, A_n$  are disjoint, and  $\bigcup_{i=1}^n A_i = S$ , with  $P(A_i) > 0$ ,  $i = 1, 2, \dots, n$ . Then for any event  $B$  with  $P(B) > 0$ ,

$$\begin{aligned} P(A_i | B) &= \frac{P(B | A_i)P(A_i)}{\sum_{j=1}^n P(B | A_j)P(A_j)}, \quad i = 1, \dots, n \\ &= \frac{P(B | A_i)P(A_i)}{P(B | A_1)P(A_1) + \dots + P(B | A_n)P(A_n)} \end{aligned}$$

## Bayes theorem (also called Bayes rule)

Suppose events  $A_1, \dots, A_n$  are disjoint, and  $\bigcup_{i=1}^n A_i = S$ , with  $P(A_i) > 0$ ,  $i = 1, 2, \dots, n$ . Then for any event  $B$  with  $P(B) > 0$ ,

$$P(A_i | B) = \frac{P(B | A_i)P(A_i)}{\sum_{j=1}^n P(B | A_j)P(A_j)}, \quad i = 1, \dots, n$$

$$= \frac{P(B | A_i)P(A_i)}{P(B | A_1)P(A_1) + \dots + P(B | A_n)P(A_n)}$$

- $P(A_i)$  is often called *prior probability*
- $P(A_i | B)$  is called *posterior probability*.

## Example: Vassar students

- Suppose Vassar has only three possible majors: math, music, and econ. And 25% of the students are in math, 55% are in music, and the rest are in econ (with no double majors).



## Example: Vassar students

- Suppose Vassar has only three possible majors: math, music, and econ. And 25% of the students are in math, 55% are in music, and the rest are in econ (with no double majors).
- Campus folklore say that 90% of math majors want to go on dates, compared with 50% of music and 10% of econ majors.

## Example: Vassar students

- Suppose Vassar has only three possible majors: math, music, and econ. And 25% of the students are in math, 55% are in music, and the rest are in econ (with no double majors).
- Campus folklore say that 90% of math majors want to go on dates, compared with 50% of music and 10% of econ majors.
- Your roommate meets someone at an event and gets a date. What is the probability that she is dating a math major?

## Example: Vassar students

- Suppose Vassar has only three possible majors: math, music, and econ. And 25% of the students are in math, 55% are in music, and the rest are in econ (with no double majors).
- Campus folklore say that 90% of math majors want to go on dates, compared with 50% of music and 10% of econ majors.
- Your roommate meets someone at an event and gets a date. What is the probability that she is dating a math major?
- Note that the majors define a finite partition, and the campus folklore gives us the conditional probabilities  $\Pr(B \mid A_i)$ .

## Example: Vassar students

- Suppose Vassar has only three possible majors: math, music, and econ. And 25% of the students are in math, 55% are in music, and the rest are in econ (with no double majors).
- Campus folklore say that 90% of math majors want to go on dates, compared with 50% of music and 10% of econ majors.
- Your roommate meets someone at an event and gets a date. What is the probability that she is dating a math major?
- Note that the majors define a finite partition, and the campus folklore gives us the conditional probabilities  $\Pr(B \mid A_i)$ .
- The point of Bayes' rule is to reverse the conditioning to get  $\Pr(A_i \mid B)$ .

Example: Suppose Vassar has only three possible majors: math, music, and econ. And 25% of the students are in math, 55% are in music, and the rest are in econ (with no double majors). Campus folklore say that 90% of math majors want to go on dates, compared with 50% of music and 10% of econ majors. Find  $P(\text{math} \mid \text{date})$ .

Example: Suppose Vassar has only three possible majors: math, music, and econ. And 25% of the students are in math, 55% are in music, and the rest are in econ (with no double majors). Campus folklore say that 90% of math majors want to go on dates, compared with 50% of music and 10% of econ majors. Find  $P(\text{math} \mid \text{date})$ .

Events  $A_1 = \{\text{math}\}$ ,  $A_2 = \{\text{music}\}$ ,  $A_3 = \{\text{econ}\}$  define a partition. Let  $B = \{\text{date}\}$ . In order to calculate  $P(A_1 \mid B)$ , use the Bayes theorem:

Example: Suppose Vassar has only three possible majors: math, music, and econ. And 25% of the students are in math, 55% are in music, and the rest are in econ (with no double majors). Campus folklore say that 90% of math majors want to go on dates, compared with 50% of music and 10% of econ majors. Find  $P(\text{math} \mid \text{date})$ .

Events  $A_1 = \{\text{math}\}$ ,  $A_2 = \{\text{music}\}$ ,  $A_3 = \{\text{econ}\}$  define a partition. Let  $B = \{\text{date}\}$ . In order to calculate  $P(A_1 \mid B)$ , use the Bayes theorem:

$$\begin{aligned} P(A_1 \mid B) &= \frac{P(B \mid A_1)P(A_1)}{P(B \mid A_1)P(A_1) + P(B \mid A_2)P(A_2) + P(B \mid A_3)P(A_3)} \\ &= \frac{0.9 \times 0.25}{0.9 \times 0.25 + 0.5 \times 0.55 + 0.1 \times 0.20} = 0.43 \end{aligned}$$

# Recap

## Bayes theorem

$$P(A_i | B) = \frac{P(B | A_i)P(A_i)}{\sum_{j=1}^n P(B | A_j)P(A_j)}, \quad i = 1, \dots, n$$



# Outline

- 1 Conditional probability
- 2 Bayes theorem
- 3 Independent events**
- 4 Perceptions of probability and biases

# Product rule for independent events

## Definition

Events  $A$  and  $B$  are *independent* if

$$P(A \cap B) = P(A) \times P(B)$$

# Product rule for independent events

## Definition

Events  $A$  and  $B$  are *independent* if

$$P(A \cap B) = P(A) \times P(B)$$

- If  $A$  and  $B$  are independent, then

$$P(A \mid B) = \frac{P(A)P(B)}{P(B)} = P(A).$$

Knowing  $B$  doesn't affect the odds of  $A$ .

Example: you toss a coin twice, what is the probability of getting two tails in a row?

Example: you toss a coin twice, what is the probability of getting two tails in a row?

- Method 1: sample space  $S = \{HH, TT, HT, TH\}$ .

Since one out of four possible outcomes match this definition, the probability is  $\frac{1}{4}$ .

*Inefficient way if the number of trials was much higher.*

Example: you toss a coin twice, what is the probability of getting two tails in a row?

- Method 1: sample space  $S = \{HH, TT, HT, TH\}$ .

Since one out of four possible outcomes match this definition, the probability is  $\frac{1}{4}$ .

*Inefficient way if the number of trials was much higher.*

- Method 2: use independence.

$$P(\text{T on the first toss}) \times P(\text{T on the second toss}) = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$$

# Properties of independence

If  $A$  and  $B$  are independent, then

- $B$  and  $A$  are independent.

# Properties of independence

If  $A$  and  $B$  are independent, then

- $B$  and  $A$  are independent.
- $A$  and  $B^c$  are independent. And so are  $A^c$  and  $B^c$ .



# Properties of independence

If  $A$  and  $B$  are independent, then

- $B$  and  $A$  are independent.
- $A$  and  $B^c$  are independent. And so are  $A^c$  and  $B^c$ .
- Independent is not disjoint.

# Mutually independent

☞ Three events  $A, B, C$  are called *mutually independent* if

1

$$P(A \cap B) = P(A)P(B)$$

$$P(B \cap C) = P(B)P(C)$$

$$P(A \cap C) = P(A)P(C)$$

2

$$P(A \cap B \cap C) = P(A)P(B)P(C)$$

# Mutually independent

☞ Three events  $A, B, C$  are called *mutually independent* if

1

$$P(A \cap B) = P(A)P(B)$$

$$P(B \cap C) = P(B)P(C)$$

$$P(A \cap C) = P(A)P(C)$$

2

$$P(A \cap B \cap C) = P(A)P(B)P(C)$$

☞ If only (1) holds but not (2), then  $A, B, C$  are called *pairwise independent*.

## Example: pairwise indpt but not mutually indpt

Roll two fair 6-sided dice. Set

- $A = \{\text{Sum is 7}\}$
- $B = \{\text{First roll is 4}\}$
- $C = \{\text{Second roll is 3}\}$

## Example: pairwise indpt but not mutually indpt

Roll two fair 6-sided dice. Set

- $A = \{\text{Sum is 7}\}$
- $B = \{\text{First roll is 4}\}$
- $C = \{\text{Second roll is 3}\}$

$$P(A \cap B) \stackrel{?}{=} P(A) \times P(B)$$

## Example: pairwise indpt but not mutually indpt

Roll two fair 6-sided dice. Set

- $A = \{\text{Sum is 7}\}$
- $B = \{\text{First roll is 4}\}$
- $C = \{\text{Second roll is 3}\}$

$$P(A \cap B) \stackrel{?}{=} P(A) \times P(B)$$

$$\frac{1}{36} \qquad \frac{1}{6} \qquad \frac{1}{6}$$

## Example: pairwise indpt but not mutually indpt

Roll two fair 6-sided dice. Set

- $A = \{\text{Sum is 7}\}$
- $B = \{\text{First roll is 4}\}$
- $C = \{\text{Second roll is 3}\}$

$$P(A \cap B) \stackrel{?}{=} P(A) \times P(B)$$

$$\frac{1}{36} \qquad \frac{1}{6} \qquad \frac{1}{6}$$

$$P(B \cap C) \stackrel{?}{=} P(B) \times P(C)$$

## Example: pairwise indpt but not mutually indpt

Roll two fair 6-sided dice. Set

- $A = \{\text{Sum is 7}\}$
- $B = \{\text{First roll is 4}\}$
- $C = \{\text{Second roll is 3}\}$

$$P(A \cap B) \stackrel{?}{=} P(A) \times P(B)$$

$$\frac{1}{36} \qquad \frac{1}{6} \qquad \frac{1}{6}$$

$$P(B \cap C) \stackrel{?}{=} P(B) \times P(C)$$

$$\frac{1}{36} \qquad \frac{1}{6} \qquad \frac{1}{6}$$



## Example: pairwise indpt but not mutually indpt

Roll two fair 6-sided dice. Set

- $A = \{\text{Sum is 7}\}$
- $B = \{\text{First roll is 4}\}$
- $C = \{\text{Second roll is 3}\}$

$$P(A \cap B) \stackrel{?}{=} P(A) \times P(B)$$

$$\frac{1}{36} \qquad \frac{1}{6} \qquad \frac{1}{6}$$

$$P(B \cap C) \stackrel{?}{=} P(B) \times P(C)$$

$$\frac{1}{36} \qquad \frac{1}{6} \qquad \frac{1}{6}$$

$$P(A \cap C) \stackrel{?}{=} P(A) \times P(C)$$

## Example: pairwise indpt but not mutually indpt

Roll two fair 6-sided dice. Set

- $A = \{\text{Sum is 7}\}$
- $B = \{\text{First roll is 4}\}$
- $C = \{\text{Second roll is 3}\}$

$$\frac{P(A \cap B)}{1/36} \stackrel{?}{=} \frac{P(A)}{1/6} \times \frac{P(B)}{1/6}$$

$$\frac{P(B \cap C)}{1/36} \stackrel{?}{=} \frac{P(B)}{1/6} \times \frac{P(C)}{1/6}$$

$$\frac{P(A \cap C)}{1/36} \stackrel{?}{=} \frac{P(A)}{1/6} \times \frac{P(C)}{1/6}$$

# Example: pairwise indpt but not mutually indpt

Roll two fair 6-sided dice. Set

- $A = \{\text{Sum is 7}\}$
- $B = \{\text{First roll is 4}\}$
- $C = \{\text{Second roll is 3}\}$

$$\frac{P(A \cap B)}{1/36} \stackrel{?}{=} \frac{P(A)}{1/6} \times \frac{P(B)}{1/6}$$

$$\frac{P(B \cap C)}{1/36} \stackrel{?}{=} \frac{P(B)}{1/6} \times \frac{P(C)}{1/6}$$

$$\frac{P(A \cap C)}{1/36} \stackrel{?}{=} \frac{P(A)}{1/6} \times \frac{P(C)}{1/6}$$

$A, B, C$  are pairwise indpt

## Example: pairwise indpt but not mutually indpt

Roll two fair 6-sided dice. Set

- $A = \{\text{Sum is 7}\}$
- $B = \{\text{First roll is 4}\}$
- $C = \{\text{Second roll is 3}\}$

$$P(A \cap B) \stackrel{?}{=} P(A) \times P(B)$$

$$\frac{1}{36} \qquad \frac{1}{6} \qquad \frac{1}{6}$$

$$P(B \cap C) \stackrel{?}{=} P(B) \times P(C)$$

$$\frac{1}{36} \qquad \frac{1}{6} \qquad \frac{1}{6}$$

$$P(A \cap C) \stackrel{?}{=} P(A) \times P(C)$$

$$\frac{1}{36} \qquad \frac{1}{6} \qquad \frac{1}{6}$$

$$P(A \cap B \cap C) \stackrel{?}{=} P(A) \times P(B) \times P(C)$$

$A, B, C$  are pairwise indpt

# Example: pairwise indpt but not mutually indpt

Roll two fair 6-sided dice. Set

- $A = \{\text{Sum is 7}\}$
- $B = \{\text{First roll is 4}\}$
- $C = \{\text{Second roll is 3}\}$

$$\frac{P(A \cap B)}{1/36} \stackrel{?}{=} \frac{P(A)}{1/6} \times \frac{P(B)}{1/6}$$

$$\frac{P(B \cap C)}{1/36} \stackrel{?}{=} \frac{P(B)}{1/6} \times \frac{P(C)}{1/6}$$

$$\frac{P(A \cap C)}{1/36} \stackrel{?}{=} \frac{P(A)}{1/6} \times \frac{P(C)}{1/6}$$

$$\frac{P(A \cap B \cap C)}{1/36} \stackrel{?}{=} \frac{P(A)}{1/6} \times \frac{P(B)}{1/6} \times \frac{P(C)}{1/6}$$

$A, B, C$  are pairwise indpt

# Example: pairwise indpt but not mutually indpt

Roll two fair 6-sided dice. Set

- $A = \{\text{Sum is 7}\}$
- $B = \{\text{First roll is 4}\}$
- $C = \{\text{Second roll is 3}\}$

$$\frac{P(A \cap B)}{1/36} \stackrel{?}{=} \frac{P(A)}{1/6} \times \frac{P(B)}{1/6}$$

$$\frac{P(B \cap C)}{1/36} \stackrel{?}{=} \frac{P(B)}{1/6} \times \frac{P(C)}{1/6}$$

$$\frac{P(A \cap C)}{1/36} \stackrel{?}{=} \frac{P(A)}{1/6} \times \frac{P(C)}{1/6}$$

$A, B, C$  are pairwise indpt

$$\frac{P(A \cap B \cap C)}{1/36} \stackrel{?}{=} \frac{P(A)}{1/6} \times \frac{P(B)}{1/6} \times \frac{P(C)}{1/6} \quad \text{NOT mutually indpt}$$

👉 Events  $A_1, A_2, \dots, A_n$  are called *independent* if for any  $1 \leq r \leq n$  of them

$$P(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_r}) = P(A_{i_1})P(A_{i_2}) \cdots P(A_{i_r})$$

- 👉 Events  $A_1, A_2, \dots, A_n$  are called *independent* if for any  $1 \leq r \leq n$  of them

$$P(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_r}) = P(A_{i_1})P(A_{i_2}) \cdots P(A_{i_r})$$

- 👉 The key to compute the probability of independent events is to just multiply the probability of the individual events.



- 👉 Events  $A_1, A_2, \dots, A_n$  are called *independent* if for any  $1 \leq r \leq n$  of them

$$P(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_r}) = P(A_{i_1})P(A_{i_2}) \cdots P(A_{i_r})$$

- 👉 The key to compute the probability of independent events is to just multiply the probability of the individual events.

A fair coin is tossed 5 times. Compute  $P(5 \text{ tails})$ .

- 👉 Events  $A_1, A_2, \dots, A_n$  are called *independent* if for any  $1 \leq r \leq n$  of them

$$P(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_r}) = P(A_{i_1})P(A_{i_2}) \cdots P(A_{i_r})$$

- 👉 The key to compute the probability of independent events is to just multiply the probability of the individual events.

A fair coin is tossed 5 times. Compute  $P(5 \text{ tails})$ .

$$P(T_1 T_2 T_3 T_4 T_5) = P(T_1)P(T_2)P(T_3)P(T_4)P(T_5) = \frac{1}{2^5}$$

# Outline

- 1 Conditional probability
- 2 Bayes theorem
- 3 Independent events
- 4 Perceptions of probability and biases**

# The Linda problem

Linda is 31 years old, single, outspoken, and very bright. She majored in philosophy. As a student she was deeply concerned with issues of discrimination and social justice, and she also participated in anti-nuclear demonstrations. Please rank the following statements by their probability, using 1 for the most probable and 8 for the least probable.

- 1 Linda is a teacher in an elementary school.
- 2 Linda works in a bookstore and takes yoga classes.
- 3 Linda is active in the feminist movement.
- 4 Linda is a psychiatric social worker.
- 5 Linda is a member of the League of Women Voters.
- 6 Linda is a bank teller.
- 7 Linda sells insurance.
- 8 Linda is a bank teller and is active in the feminist movement.

## The Linda problem cont'd

Linda is 31 years old, single, outspoken, and very bright. She majored in philosophy. As a student she was deeply concerned with issues of discrimination and social justice, and she also participated in anti-nuclear demonstrations. Please rank the following statements by their probability, using 1 for the most probable and 8 for the least probable.

- 1 Linda is a teacher in an elementary school.
- 2 Linda works in a bookstore and takes yoga classes.
- 3 Linda is active in the feminist movement.
- 4 Linda is a psychiatric social worker.
- 5 Linda is a member of the League of Women Voters.
- 6 Linda is a bank teller.
- 7 Linda sells insurance.
- 8 Linda is a bank teller and is active in the feminist movement.

## Set theory - intersection

- Event  $A = \{\text{Linda is active in the feminist movement}\}$
- Event  $B = \{\text{Linda is a bank teller}\}$
- Event  $A \cap B = \{\text{Linda is a bank teller and is active in the feminist movement}\}$ .

## Set theory - intersection

- Event  $A = \{\text{Linda is active in the feminist movement}\}$
- Event  $B = \{\text{Linda is a bank teller}\}$
- Event  $A \cap B = \{\text{Linda is a bank teller and is active in the feminist movement}\}$ .
- $P(A \cap B) \leq P(A)$  and  $P(A \cap B) \leq P(B)$ !
- What did our class do? 21 of you responded.
  - ▶ Ranking: 1 is the most probable choice, and 8 is the least probable choice
  - ▶  $\text{Avg}(\text{Ranking of event } A) = 3.38$
  - ▶  $\text{Avg}(\text{Ranking of event } B) = 4.76$
  - ▶  $\text{Avg}(\text{Ranking of event } A \cap B) = 4.62$
  - ▶ Percentage of people ranking event  $A \cap B$  higher than event  $B \approx 62\%$  (13 out of 21 responses)

## The Linda problem - historical results

The Linda problem was proposed by Amos Tversky and Daniel Kahneman in 1983.

- Ranking: 1 is the most probable choice, and 8 is the least probable choice
- $\text{Avg}(\text{Ranking of event } A) = 2.1$
- $\text{Avg}(\text{Ranking of event } B) = 6.2$
- $\text{Avg}(\text{Ranking of event } A \cap B) = 4.1$
- Percentage of people ranking event  $A \cap B$  higher than event  $B = 90\%$  (even among those who have had several advanced courses in probability and statistics)



# Representativeness and Conjunction Bias

- Representativeness heuristic:
  - ▶ people determine the probability of an event by the degree to which the event is similar in essential characteristics to its parent population.
  - ▶ it leads people to inflate the believed probability of events that resembles the data available.

# Representativeness and Conjunction Bias

- Representativeness heuristic:
  - ▶ people determine the probability of an event by the degree to which the event is similar in essential characteristics to its parent population.
  - ▶ it leads people to inflate the believed probability of events that resembles the data available.
- Conjunction bias:
  - ▶ intersecting relatively unrepresentative events (bank teller) with very representative event (feminist).
  - ▶ the description of the conjunction of the two (feminist bank teller) is considered more probable than the unrepresentative event (bank teller).

# Representativeness and Conjunction Bias

- Representativeness heuristic:
  - ▶ people determine the probability of an event by the degree to which the event is similar in essential characteristics to its parent population.
  - ▶ it leads people to inflate the believed probability of events that resembles the data available.
- Conjunction bias:
  - ▶ intersecting relatively unrepresentative events (bank teller) with very representative event (feminist).
  - ▶ the description of the conjunction of the two (feminist bank teller) is considered more probable than the unrepresentative event (bank teller).
- Interested in these types of topics? ECON 333 Behavioral Economics!