

Chapter 2

Axioms of Probability

Jingchen (Monika) Hu

Vassar College

MATH 241

Outline

- 1 Sample space and events
- 2 Axioms of Probability
- 3 Some simple propositions
- 4 Sample spaces with equally likely outcomes

Sample space

Definition

A sample space S is the set of all possible outcomes of an experiment.

Examples: 3 coin tosses

$$S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$$

One die roll

$$S = \{1, 2, 3, 4, 5, 6\}$$

Sum of two rolls

$$S = \{2, 3, \dots, 11, 12\}$$

Seconds waiting for bus

$$S = [0, \infty)$$

Examples of sample spaces

- Experiment is playing five rounds of Russian roulette
Sample space is $\{D, LD, LLD, LLLD, LLLLD, LLLLL\}$.

Why is this different than coin flipping?

Examples of sample spaces

- Experiment is playing five rounds of Russian roulette
Sample space is $\{D, LD, LLD, LLLD, LLLLD, LLLLL\}$.

Why is this different than coin flipping?

- Experiment is sequencing three nucleotides
Sample space is $\{AAA, CCC, GGG, TTT, AAC, AAT, AAG, \dots\}$.

Examples of sample spaces

- Experiment is playing five rounds of Russian roulette
Sample space is $\{D, LD, LLD, LLLD, LLLLD, LLLLL\}$.

Why is this different than coin flipping?

- Experiment is sequencing three nucleotides
Sample space is $\{AAA, CCC, GGG, TTT, AAC, AAT, AAG, \dots\}$.

How big is this sample space? (Hint: there are four types of nucleotides.)

Examples of sample spaces

- Experiment is playing five rounds of Russian roulette
Sample space is $\{D, LD, LLD, LLLD, LLLLD, LLLLL\}$.

Why is this different than coin flipping?

- Experiment is sequencing three nucleotides
Sample space is $\{AAA, CCC, GGG, TTT, AAC, AAT, AAG, \dots\}$.

How big is this sample space? (Hint: there are four types of nucleotides.)

$$4 \times 4 \times 4 = 64$$

An event

Definition

An event E is any subset of the sample space S .

$$E \subset S$$

An event

Definition

An event E is any subset of the sample space S .

$$E \subset S$$

Examples: 2 heads $E = \{\text{HHT, HTH, THH}\}$

Even number $E = \{2, 4, 6\}$

< 2 minutes $E = [0, 120)$

An event

Definition

An event E is any subset of the sample space S .

$$E \subset S$$

Examples: 2 heads $E = \{\text{HHT, HTH, THH}\}$

Even number $E = \{2, 4, 6\}$

< 2 minutes $E = [0, 120)$

- Impossible event: empty set $\emptyset \subset S$
- $S \subset S$

Set theory

Let A, B be two events.

Definition

- ➊ **Intersection** $A \cap B$: *implies the event that both A and B occur*
- ➋ **Union** $A \cup B$: *implies the event that at least one of A or B occur*
- ➌ The **complement** of an event A denoted A^c (also notated A' or \bar{A}):
 $A^c = S \setminus A$ - *the event that A does not occur*
- ➍ $A \subset B$ *implies that the occurrence of A implies the occurrence of B*

Venn diagram

More set theory

Definition

Two events A and B are **mutually exclusive** or **disjoint** if they have no outcomes in common, i.e. $A \cap B = \emptyset$.

More set theory

Definition

Two events A and B are **mutually exclusive** or **disjoint** if they have no outcomes in common, i.e. $A \cap B = \emptyset$.

Examples of exclusive events?

Some rules

① Commutative laws

$$A \cup B = B \cup A, \quad A \cap B = B \cap A$$

Some rules

1 Commutative laws

$$A \cup B = B \cup A, \quad A \cap B = B \cap A$$

2 Associative laws

$$(A \cup B) \cup C = A \cup (B \cup C)$$

$$(A \cap B) \cap C = A \cap (B \cap C)$$

Some rules

1 Commutative laws

$$A \cup B = B \cup A, \quad A \cap B = B \cap A$$

2 Associative laws

$$(A \cup B) \cup C = A \cup (B \cup C)$$

$$(A \cap B) \cap C = A \cap (B \cap C)$$

3 Distributive laws

$$(A \cup B) \cap C = (A \cap C) \cup (B \cap C)$$

$$(A \cap B) \cup C = (A \cup C) \cap (B \cup C)$$

DeMorgan's Laws

$$(A \cup B)^c = A^c \cap B^c, \quad (A \cap B)^c = A^c \cup B^c$$

DeMorgan's Laws

$$(A \cup B)^c = A^c \cap B^c, \quad (A \cap B)^c = A^c \cup B^c$$

- Outline of proof (to the first equation): two steps
Left \subset Right \iff For any $x \in (A \cup B)^c$, then $x \in A^c \cap B^c$.
Right \subset Left \iff For any $x \in A^c \cap B^c$, then $x \in (A \cup B)^c$.

DeMorgan's Laws

$$(A \cup B)^c = A^c \cap B^c, \quad (A \cap B)^c = A^c \cup B^c$$

- Outline of proof (to the first equation): two steps
 Left \subset Right \iff For any $x \in (A \cup B)^c$, then $x \in A^c \cap B^c$.
 Right \subset Left \iff For any $x \in A^c \cap B^c$, then $x \in (A \cup B)^c$.

- DeMorgan's laws can be generalized to n events A_1, \dots, A_n :

$$\left(\bigcup_{i=1}^n A_i \right)^c = \bigcap_{i=1}^n A_i^c, \quad \left(\bigcap_{i=1}^n A_i \right)^c = \bigcup_{i=1}^n A_i^c$$

DeMorgan's Laws

$$(A \cup B)^c = A^c \cap B^c, \quad (A \cap B)^c = A^c \cup B^c$$

- Outline of proof (to the first equation): two steps
 Left \subset Right \iff For any $x \in (A \cup B)^c$, then $x \in A^c \cap B^c$.
 Right \subset Left \iff For any $x \in A^c \cap B^c$, then $x \in (A \cup B)^c$.

- DeMorgan's laws can be generalized to n events A_1, \dots, A_n :

$$\left(\bigcup_{i=1}^n A_i \right)^c = \bigcap_{i=1}^n A_i^c, \quad \left(\bigcap_{i=1}^n A_i \right)^c = \bigcup_{i=1}^n A_i^c$$

<https://www.youtube.com/watch?v=oOx7FSzSav4>

Recap

- Sample space S , event E
- Set operations: intersection, union, complement, subset, disjoint/mutually exclusive
- Rules: commutative, associative, distributive, DeMorgan's laws
- Hint: Venn Diagram is helpful

Outline

- 1 Sample space and events
- 2 Axioms of Probability**
- 3 Some simple propositions
- 4 Sample spaces with equally likely outcomes

What is probability?

Given an experiment and a sample space S , the objective of probability is to assign to each event A a number $P(A)$, called the probability of the event A , which will give a precise measure of the chance that A will occur.

What is probability?

Given an experiment and a sample space S , the objective of probability is to assign to each event A a number $P(A)$, called the probability of the event A , which will give a precise measure of the chance that A will occur.

What is $P(A)$ of the following events (A 's)?

- ① $A =$ Someone in this class room wins the MegaMillion.
- ② $A =$ You spin a quarter and it comes up heads.
- ③ $A =$ You spin a quarter and it stands up.
- ④ $A =$ At least two people in this class room have the same birthday.

Probability: Frequentist interpretation

- There are several possible interpretations of probability.
- There is no agreement, at all, in how probabilities should be interpreted.
- There is (nearly) complete agreement on the mathematical rules probability must follow (axioms).

Probability: Frequentist interpretation

- There are several possible interpretations of probability.
- There is no agreement, at all, in how probabilities should be interpreted.
- There is (nearly) complete agreement on the mathematical rules probability must follow (axioms).
- *Frequentist* interpretation: The probability of event A is the proportion of times (frequency) that A occurs in an infinite sequence (or very long run) of separate tries of the experiment.

$$P(A) = \lim_{n \rightarrow \infty} \frac{\# \text{ times } A \text{ happens}}{n}.$$

Probability: Frequentist interpretation

- There are several possible interpretations of probability.
- There is no agreement, at all, in how probabilities should be interpreted.
- There is (nearly) complete agreement on the mathematical rules probability must follow (axioms).
- *Frequentist* interpretation: The probability of event A is the proportion of times (frequency) that A occurs in an infinite sequence (or very long run) of separate tries of the experiment.

$$P(A) = \lim_{n \rightarrow \infty} \frac{\# \text{ times } A \text{ happens}}{n}.$$

- Often associated with Jerzy Neyman and Egon Pearson who described the logic of statistical hypothesis testing.

Probability: Bayesian interpretation

- A *Bayesian* can pick whatever number they prefer for $P(A)$, based on their own personal experience and intuition, provided that number is consistent with all of the other probabilities they choose in life.

Probability: Bayesian interpretation

- A *Bayesian* can pick whatever number they prefer for $P(A)$, based on their own personal experience and intuition, provided that number is consistent with all of the other probabilities they choose in life.
- A Bayesian interprets probability as a subjective degree of belief: *For the same event, two separate people could have differing probabilities.*

Probability: Bayesian interpretation

- A *Bayesian* can pick whatever number they prefer for $P(A)$, based on their own personal experience and intuition, provided that number is consistent with all of the other probabilities they choose in life.
- A Bayesian interprets probability as a subjective degree of belief: *For the same event, two separate people could have differing probabilities.*
- The Bayesian's view must: (1) conform to all other personal opinions; (2) change as new data arise according to *Bayes' Rule*.

Probability: Bayesian interpretation

- A *Bayesian* can pick whatever number they prefer for $P(A)$, based on their own personal experience and intuition, provided that number is consistent with all of the other probabilities they choose in life.
- A Bayesian interprets probability as a subjective degree of belief: *For the same event, two separate people could have differing probabilities.*
- The Bayesian's view must: (1) conform to all other personal opinions; (2) change as new data arise according to *Bayes' Rule*.
- Bayesian interpretations of probabilities avoid some of the philosophical difficulties of frequency interpretations.

Probability: Bayesian interpretation

- A *Bayesian* can pick whatever number they prefer for $P(A)$, based on their own personal experience and intuition, provided that number is consistent with all of the other probabilities they choose in life.
- A Bayesian interprets probability as a subjective degree of belief: *For the same event, two separate people could have differing probabilities.*
- The Bayesian's view must: (1) conform to all other personal opinions; (2) change as new data arise according to *Bayes' Rule*.
- Bayesian interpretations of probabilities avoid some of the philosophical difficulties of frequency interpretations.
- Named after the 18th century Presbyterian minister and mathematician Thomas Bayes.
- Largely popularized by revolutionary advance in computational technology and methods during the last twenty years.

Axiomatic Foundations of Probability

- The foundations of modern probability theory, the axiomatic basis, are laid by Andrey Kolmogorov in 1933.



Axiomatic Foundations of Probability

- The foundations of modern probability theory, the axiomatic basis, are laid by Andrey Kolmogorov in 1933.



- The axiomatic approach is not concerned with the interpretation of probabilities.
- Concerned only that probabilities are defined by a function satisfying the axioms.

Axiomatic Foundations of Probability

- The foundations of modern probability theory, the axiomatic basis, are laid by Andrey Kolmogorov in 1933.



- The axiomatic approach is not concerned with the interpretation of probabilities.
- Concerned only that probabilities are defined by a function satisfying the axioms.
- Kolmogorov (1903-1987) was one of the greatest mathematicians of the 20th century. This axiomatization was one of his “trivial” accomplishment.

Axioms of probability

Definition

*Let P be a function that assigns a nonnegative real number to each event E of a sample space S . We call P a **probability** if*

Axioms of probability

Definition

Let P be a function that assigns a nonnegative real number to each event E of a sample space S . We call P a *probability* if

① Axiom 1: non-negative

$$0 \leq P(E) \leq 1$$

Axioms of probability

Definition

Let P be a function that assigns a nonnegative real number to each event E of a sample space S . We call P a *probability* if

① Axiom 1: non-negative

$$0 \leq P(E) \leq 1$$

② Axiom 2: total one

$$P(S) = 1$$

Axioms of probability

Definition

Let P be a function that assigns a nonnegative real number to each event E of a sample space S . We call P a *probability* if

① Axiom 1: non-negative

$$0 \leq P(E) \leq 1$$

② Axiom 2: total one

$$P(S) = 1$$

③ Axiom 3: countable addition

$$P\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} P(E_i), \text{ if } E_i \cap E_j = \emptyset \text{ for } i \neq j$$

Axioms of probability

Definition

Let P be a function that assigns a nonnegative real number to each event E of a sample space S . We call P a **probability** if

- ① Axiom 1: non-negative

$$0 \leq P(E) \leq 1$$

- ② Axiom 2: total one

$$P(S) = 1$$

- ③ Axiom 3: countable addition

$$P\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} P(E_i), \text{ if } E_i \cap E_j = \emptyset \text{ for } i \neq j$$

In particular, for k **disjoint** events E_1, \dots, E_k ,

$$P\left(\bigcup_{i=1}^k E_i\right) = \sum_{i=1}^k P(E_i)$$

Outline

- 1 Sample space and events
- 2 Axioms of Probability
- 3 Some simple propositions**
- 4 Sample spaces with equally likely outcomes

Propositions

👉 Complement Rule:

$$P(A^c) = 1 - P(A)$$

Proof (hint: use Axiom 2 and Axiom 3)

Propositions

👉 Complement Rule:

$$P(A^c) = 1 - P(A)$$

Proof (hint: use Axiom 2 and Axiom 3)

- $P(\emptyset) = 0$

👉 Difference Rule:

$$P(B \cap A^c) = P(B) - P(A), \text{ if } A \subseteq B$$

Proof (hint: use Axiom 3)

👉 Difference Rule:

$$P(B \cap A^c) = P(B) - P(A), \text{ if } A \subseteq B$$

Proof (hint: use Axiom 3)

- $P(B) \geq P(A), \text{ if } A \subseteq B$

Proof (hint: use Axiom 1)

👉 Inclusion-Exclusion: two events A, B (not necessarily disjoint)

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

Proof (hint: use Axiom 3 and Venn-diagram)

Inclusion-Exclusion example

Suppose that for a randomly selected student in a probability class,

- $P(\text{live Eastern Time Zone at home}) = 63\%$.
- $P(\text{senior}) = 41\%$.
- $P(\text{live Eastern Time Zone at home and senior}) = 31\%$.

Find the probability that a student is either live Eastern Time Zone at home or a senior.

Inclusion-Exclusion example

Suppose that for a randomly selected student in a probability class,

- $P(\text{live Eastern Time Zone at home}) = 63\%$.
- $P(\text{senior}) = 41\%$.
- $P(\text{live Eastern Time Zone at home and senior}) = 31\%$.

Find the probability that a student is either live Eastern Time Zone at home or a senior.

- 1 Conditions: event $A = \{\text{live Eastern Time Zone at home}\}$,
 $B = \{\text{senior}\}$,

$$P(A) = 0.63, P(B) = 0.41, P(A \cap B) = 0.31$$

Inclusion-Exclusion example

Suppose that for a randomly selected student in a probability class,

- $P(\text{live Eastern Time Zone at home}) = 63\%$.
- $P(\text{senior}) = 41\%$.
- $P(\text{live Eastern Time Zone at home and senior}) = 31\%$.

Find the probability that a student is either live Eastern Time Zone at home or a senior.

- ① Conditions: event $A = \{\text{live Eastern Time Zone at home}\}$,
 $B = \{\text{senior}\}$,

$$P(A) = 0.63, P(B) = 0.41, P(A \cap B) = 0.31$$

- ② Question: find $P(A \cup B)$.

Inclusion-Exclusion example

Suppose that for a randomly selected student in a probability class,

- $P(\text{live Eastern Time Zone at home}) = 63\%$.
- $P(\text{senior}) = 41\%$.
- $P(\text{live Eastern Time Zone at home and senior}) = 31\%$.

Find the probability that a student is either live Eastern Time Zone at home or a senior.

- ① Conditions: event $A = \{\text{live Eastern Time Zone at home}\}$,
 $B = \{\text{senior}\}$,

$$P(A) = 0.63, P(B) = 0.41, P(A \cap B) = 0.31$$

- ② Question: find $P(A \cup B)$.


- ③ Formula: inclusion-exclusion

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.63 + 0.41 - 0.31 = 0.73$$

Propositions

👉 Inclusion-Exclusion: three events A, B, C (not necessarily disjoint)

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$$



Recap

Three axioms of probability P

① $0 \leq P(E) \leq 1$

② $P(S) = 1$

③ $P\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} P(E_i)$, if $E_i \cap E_j = \emptyset$ for $i \neq j$

Recap

Three axioms of probability P

- 1 $0 \leq P(E) \leq 1$
- 2 $P(S) = 1$
- 3 $P\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} P(E_i)$, if $E_i \cap E_j = \emptyset$ for $i \neq j$

Propositions of probability

- $P(A^c) = 1 - P(A)$
- $P(B \cap A^c) = P(B) - P(A)$, if $A \subseteq B$
- $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
- $P(A \cup B \cup C) =$
 $P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$

Outline

- 1 Sample space and events
- 2 Axioms of Probability
- 3 Some simple propositions
- 4 Sample spaces with equally likely outcomes**

Sample spaces with equally likely outcomes

- ☞ Suppose a sample space has N equally likely outcomes $\{1\}, \dots, \{N\}$, then

$$S = \{1\} \cup \dots \cup \{N\}$$

Disjointness gives

$$1 = P(S) = P(\{1\}) + \dots + P(\{N\}) = NP(\{i\}),$$

so for each $1 \leq i \leq N$,

$$P(\{i\}) = \frac{1}{N}$$

Sample spaces with equally likely outcomes

- ☞ Suppose a sample space has N equally likely outcomes $\{1\}, \dots, \{N\}$, then

$$S = \{1\} \cup \dots \cup \{N\}$$

Disjointness gives

$$1 = P(S) = P(\{1\}) + \dots + P(\{N\}) = NP(\{i\}),$$

so for each $1 \leq i \leq N$,

$$P(\{i\}) = \frac{1}{N}$$

- ☞ For event E in a sample space S with equally likely outcomes,

$$P(E) = \frac{\#(E)}{\#(S)}$$

Notation:

Cardinality - $\#(E)$ = number of elements in set E

Probability of rolling an even number with a six sided die?

Probability of rolling an even number with a six sided die?

$$E = \{2, 4, 6\} \text{ and } S = \{1, 2, 3, 4, 5, 6\}$$

$$P(E) = 3/6 = 1/2$$

Probability of rolling an even number with a six sided die?

$$E = \{2, 4, 6\} \text{ and } S = \{1, 2, 3, 4, 5, 6\}$$

$$P(E) = 3/6 = 1/2$$

A couple has two kids, what is the probability that they are not both girls?

Probability of rolling an even number with a six sided die?

$$E = \{2, 4, 6\} \text{ and } S = \{1, 2, 3, 4, 5, 6\}$$

$$P(E) = 3/6 = 1/2$$

A couple has two kids, what is the probability that they are not both girls?

$$E = \{BB, GB, BG\} \text{ and } S = \{BB, GG, GB, BG\}$$

$$P(E) = 3/4$$

Example: birthday problem

Ignoring leap years, and assuming birthdays are equally likely to be any day of the year, what is the chance of no tie in birthdays among n students?

Example: birthday problem

Ignoring leap years, and assuming birthdays are equally likely to be any day of the year, what is the chance of no tie in birthdays among n students?

$$\#(\text{birthdays of } n \text{ people}) = 365^n$$

$$\#(\text{no match}) = 365 \times 364 \times \cdots \times (365 - n + 1)$$

$$P(\text{no match}) = \frac{365!}{(365 - n)! 365^n}$$

Example: birthday problem

Ignoring leap years, and assuming birthdays are equally likely to be any day of the year, what is the chance of no tie in birthdays among n students?

$$\#(\text{birthdays of } n \text{ people}) = 365^n$$

$$\#(\text{no match}) = 365 \times 364 \times \cdots \times (365 - n + 1)$$

$$P(\text{no match}) = \frac{365!}{(365 - n)! 365^n}$$

