

Chapter 4 part 1

Discrete Random Variables

Jingchen (Monika) Hu

Vassar College

MATH 241

Outline

- 1 Discrete random variables
- 2 Expectation

Outline

1 Discrete random variables

2 Expectation

Random Variables

Definition

Random Variable X is a real-valued function on the sample space S .

Random Variables

Definition

Random Variable X is a real-valued function on the sample space S .

Example: If $S = \{a = (a_1, a_2) : 1 \leq a_1, a_2 \leq 6\}$ is the 36 element space resulting from rolling two fair six-sided dice, then the following are all random variables

$$X(a) = a_1$$

$$Y(a) = |a_1 - a_2|$$

$$Z(a) = a_1 + a_2$$

Random Variables

Definition

Random Variable X is a real-valued function on the sample space S .

Example: If $S = \{a = (a_1, a_2) : 1 \leq a_1, a_2 \leq 6\}$ is the 36 element space resulting from rolling two fair six-sided dice, then the following are all random variables

$$X(a) = a_1$$

$$Y(a) = |a_1 - a_2|$$

$$Z(a) = a_1 + a_2$$

- Random variable is a number associated with a random experiment.
- Random variables are in essence a fancy way of describing an event, e.g.

$$P(X = 1) = 1/6$$

Example: a coin is flipped until a head is obtained. The flips are independent and each one has probability p of heads. Let the random variable X denote the number of flips until a head is obtained.

Example: a coin is flipped until a head is obtained. The flips are independent and each one has probability p of heads. Let the random variable X denote the number of flips until a head is obtained.

$$P(X = 1) = P(\text{H}) = p$$

$$P(X = 2) = P(\text{TH}) = (1 - p)p$$

$$P(X = 3) = P(\text{TTH}) = (1 - p)^2 p$$

.....

$$P(X = n) = P(\text{TT} \cdots \text{TH}) = (1 - p)^{n-1} p$$

.....

Cumulative distribution function (cdf)

Definition

For a random variable X , the function F defined by

$$F(x) = P(X \leq x), \quad -\infty < x < \infty$$

is called the *cumulative distribution function* of X .

Cumulative distribution function (cdf)

Definition

For a random variable X , the function F defined by

$$F(x) = P(X \leq x), \quad -\infty < x < \infty$$

is called the *cumulative distribution function* of X .

Note that

- Capital X : random variable
- Little x : a real-valued number
- \leq : smaller than or equal to

Cumulative distribution function (cdf)

Definition

For a random variable X , the function F defined by

$$F(x) = P(X \leq x), \quad -\infty < x < \infty$$

is called the *cumulative distribution function* of X .

Note that

- Capital X : random variable
- Little x : a real-valued number
- \leq : smaller than or equal to

In the previous coin flipping example,

$$F(n) = P(X \leq n) = \sum_{i=1}^n (1-p)^{i-1} p = \frac{[1 - (1-p)^n]p}{1 - (1-p)} = 1 - (1-p)^n$$

Properties of the cdf (Ch 4.10)

- $P(a < X \leq b) = F(b) - F(a)$, for all $a < b$

Properties of the cdf (Ch 4.10)

- $P(a < X \leq b) = F(b) - F(a)$, for all $a < b$
- $P(X < b)$ does not necessary equal to $P(X \leq b)$.

Properties of the cdf (Ch 4.10)

- $P(a < X \leq b) = F(b) - F(a)$, for all $a < b$
- $P(X < b)$ does not necessary equal to $P(X \leq b)$.
- ① $F(x)$ is a non-decreasing function; i.e., if $x_1 < x_2$, then

$$F(x_1) \leq F(x_2)$$

Properties of the cdf (Ch 4.10)

- $P(a < X \leq b) = F(b) - F(a)$, for all $a < b$
- $P(X < b)$ does not necessary equal to $P(X \leq b)$.
- ① $F(x)$ is a non-decreasing function; i.e., if $x_1 < x_2$, then

$$F(x_1) \leq F(x_2)$$

②

$$\lim_{x \rightarrow \infty} F(x) = 1 \iff P(X \leq \infty) = 1$$

Properties of the cdf (Ch 4.10)

- $P(a < X \leq b) = F(b) - F(a)$, for all $a < b$
- $P(X < b)$ does not necessary equal to $P(X \leq b)$.
- ① $F(x)$ is a non-decreasing function; i.e., if $x_1 < x_2$, then

$$F(x_1) \leq F(x_2)$$

②

$$\lim_{x \rightarrow \infty} F(x) = 1 \iff P(X \leq \infty) = 1$$

③

$$\lim_{x \rightarrow -\infty} F(x) = 0 \iff P(X > -\infty) = 1$$

Properties of the cdf (Ch 4.10)

- $P(a < X \leq b) = F(b) - F(a)$, for all $a < b$
- $P(X < b)$ does not necessary equal to $P(X \leq b)$.
- ① $F(x)$ is a non-decreasing function; i.e., if $x_1 < x_2$, then

$$F(x_1) \leq F(x_2)$$

- ②
- $$\lim_{x \rightarrow \infty} F(x) = 1 \iff P(X \leq \infty) = 1$$

- ③
- $$\lim_{x \rightarrow -\infty} F(x) = 0 \iff P(X > -\infty) = 1$$

- ④ $F(x)$ is right continuous; i.e., for any decreasing sequence $\{x_n : n = 1, 2, \dots\}$ that converges to x ,

$$\lim_{n \rightarrow \infty} F(x_n) = F(x) \implies \lim_{n \rightarrow \infty} P(X \leq x + \frac{1}{n}) = P(X \leq x)$$

The cdf tells us everything about a random variable X




FIGURE 4.8: Graph of $F(x)$.

The cdf tells us everything about a random variable X




FIGURE 4.8: Graph of $F(x)$.

- $P(X \leq 0.5) = 0.5 \times \frac{1/2}{1} = 0.25$

The cdf tells us everything about a random variable X




FIGURE 4.8: Graph of $F(x)$.

- $P(X \leq 0.5) = 0.5 \times \frac{1/2}{1} = 0.25$
- $P(X > 2) = 1 - P(X \leq 2) = \frac{1}{12}$

The cdf tells us everything about a random variable X




FIGURE 4.8: Graph of $F(x)$.

- $P(X \leq 0.5) = 0.5 \times \frac{1}{2} = 0.25$
- $P(X < 1) = 0.5, P(X \leq 1) = \frac{2}{3}$
- $P(X > 2) = 1 - P(X \leq 2) = \frac{1}{12}$

The cdf tells us everything about a random variable X




FIGURE 4.8: Graph of $F(x)$.

- $P(X \leq 0.5) = 0.5 \times \frac{1}{1} = 0.25$
- $P(X > 2) = 1 - P(X \leq 2) = \frac{1}{12}$
- $P(X < 1) = 0.5, P(X \leq 1) = \frac{2}{3}$
- $P(X = 1) = P(X \leq 1) - P(X < 1) = \frac{2}{3} - \frac{1}{2} = \frac{1}{6}$

Discrete random variables

Definition

☞ A random variable X that can take on at most a countable number of possible values is a *discrete random variable*.

Discrete random variables

Definition

- ➡ A random variable X that can take on at most a countable number of possible values is a *discrete random variable*.
- ➡ For a discrete random variable X , we define the *probability mass function* (pmf) by

$$p(x) = P(X = x)$$

Note: pmf is also written as $f(x)$.

Discrete random variables

Definition

- ➡ A random variable X that can take on at most a countable number of possible values is a *discrete random variable*.
- ➡ For a discrete random variable X , we define the *probability mass function* (pmf) by

$$p(x) = P(X = x)$$

Note: pmf is also written as $f(x)$.

Examples of discrete random variable

- Flip a coin 10 times, X = number of heads.
 X can be $0, 1, 2, \dots, 10$, finite. \implies discrete random variable.

Discrete random variables

Definition

- ☞ A random variable X that can take on at most a countable number of possible values is a *discrete random variable*.
- ☞ For a discrete random variable X , we define the *probability mass function* (pmf) by

$$p(x) = P(X = x)$$

Note: pmf is also written as $f(x)$.

Examples of discrete random variable

- Flip a coin 10 times, X = number of heads.
 X can be $0, 1, 2, \dots, 10$, finite. \implies discrete random variable.
- X = number of your ex boyfriends / girlfriends.
 X can be $0, 1, 2, \dots, \infty$, countable. \implies discrete random variable.

Discrete random variables

Definition

- 👉 A random variable X that can take on at most a countable number of possible values is a *discrete random variable*.
- 👉 For a discrete random variable X , we define the *probability mass function* (pmf) by

$$p(x) = P(X = x)$$

Note: pmf is also written as $f(x)$.

Examples of discrete random variable

- Flip a coin 10 times, X = number of heads.
 X can be $0, 1, 2, \dots, 10$, finite. \implies discrete random variable.
- X = number of your ex boyfriends / girlfriends.
 X can be $0, 1, 2, \dots, \infty$, countable. \implies discrete random variable.
- X = a random number in $[0, 1]$ generated by computer
 X can anything in $[0, 1]$, uncountable. \implies not discrete random variable.

pmf and cdf

- For a discrete random variable X , there exists a countable sequence x_1, x_2, \dots , such that

$$p(x_i) > 0 \quad \text{for } i = 1, 2, \dots$$

$$p(x) = 0 \quad \text{for all other values of } x$$

pmf and cdf

- For a discrete random variable X , there exists a countable sequence x_1, x_2, \dots , such that

$$p(x_i) > 0 \quad \text{for } i = 1, 2, \dots$$

$$p(x) = 0 \quad \text{for all other values of } x$$

and

$$\sum_{i=1}^{\infty} p(x_i) = 1$$

pmf and cdf

- For a discrete random variable X , there exists a countable sequence x_1, x_2, \dots , such that

$$p(x_i) > 0 \quad \text{for } i = 1, 2, \dots$$

$$p(x) = 0 \quad \text{for all other values of } x$$

and

$$\sum_{i=1}^{\infty} p(x_i) = 1$$

- Relationship between pmf and cdf (for discrete random variable)

$$F(a) = \sum_{\text{all } x \leq a} p(x)$$

- If we know pmf, we can compute cdf. And vice versa.

X has a pmf given by

$$p(1) = \frac{1}{4}, p(2) = \frac{1}{2}, p(3) = \frac{1}{8}, p(4) = \frac{1}{8}$$

X has a pmf given by


$$p(1) = \frac{1}{4}, \quad p(2) = \frac{1}{2}, \quad p(3) = \frac{1}{8}, \quad p(4) = \frac{1}{8}$$

$$F(a) = \begin{cases} 0 & a < 1 \\ \frac{1}{4} & 1 \leq a < 2 \\ \frac{3}{4} & 2 \leq a < 3 \\ \frac{7}{8} & 3 \leq a < 4 \\ 1 & 4 \leq a \end{cases}$$

X has a pmf given by

$$p(1) = \frac{1}{4}, \quad p(2) = \frac{1}{2}, \quad p(3) = \frac{1}{8}, \quad p(4) = \frac{1}{8}$$

$$F(a) = \begin{cases} 0 & a < 1 \\ \frac{1}{4} & 1 \leq a < 2 \\ \frac{3}{4} & 2 \leq a < 3 \\ \frac{7}{8} & 3 \leq a < 4 \\ 1 & 4 \leq a \end{cases}$$



Recap

Random Variable

- Random Variable X is a real-valued function on the sample space S .

$$X : S \longrightarrow \mathbb{R}$$

- Cumulative distribution function (cdf)

$$F_X(x) = P(X \leq x), \quad \text{for any } x \in \mathbb{R}$$

Recap

Random Variable

- Random Variable X is a real-valued function on the sample space S .

$$X : S \longrightarrow \mathbb{R}$$

- Cumulative distribution function (cdf)

$$F_X(x) = P(X \leq x), \quad \text{for any } x \in \mathbb{R}$$

Discrete random variable

- can only take at most a countable number of possible values.
- probability mass function (pmf)

$$p_X(x) = P(X = x)$$

Recap

Random Variable

- Random Variable X is a real-valued function on the sample space S .

$$X : S \longrightarrow \mathbb{R}$$

- Cumulative distribution function (cdf)

$$F_X(x) = P(X \leq x), \quad \text{for any } x \in \mathbb{R}$$

Discrete random variable

- can only take at most a countable number of possible values.
- probability mass function (pmf)

$$p_X(x) = P(X = x)$$

- $\sum_{i=1}^{\infty} p(x_i) = 1$

- $F(a) = \sum_{\text{all } x \leq a} p(x)$

Outline

1 Discrete random variables

2 **Expectation**

Expected value

Definition

The *expected value* (or *mean*) of a discrete random variable is defined as

$$E[X] = \sum_{x:p(x)>0} x \cdot P(X = x) = \sum_{x:p(x)>0} xp(x)$$

Expected value

Definition

The *expected value* (or *mean*) of a discrete random variable is defined as

$$E[X] = \sum_{x:p(x)>0} x \cdot P(X = x) = \sum_{x:p(x)>0} xp(x)$$

- $E[X]$ is a weighted average of the possible values x that X can take on, each value being weighted by the probability $p(x)$.

Expected value

Definition

The *expected value* (or *mean*) of a discrete random variable is defined as

$$E[X] = \sum_{x:p(x)>0} x \cdot P(X = x) = \sum_{x:p(x)>0} xp(x)$$

- $E[X]$ is a weighted average of the possible values x that X can take on, each value being weighted by the probability $p(x)$.



When she told me I was average,
she was just being mean.

Example: X is the outcome when we roll a 4-sided fair die. Find $E[X]$.

Example: X is the outcome when we roll a 4-sided fair die. Find $E[X]$.

$$p(1) = p(2) = p(3) = p(4) = \frac{1}{4}$$

$$E[X] = 1 \times \frac{1}{4} + 2 \times \frac{1}{4} + 3 \times \frac{1}{4} + 4 \times \frac{1}{4} = 2.5$$

Expectation of a function of a random variable

- If X is a discrete random variable, and g is a real-valued function then the expectation (or expected value) of $Y = g(X)$ is

$$E[g(X)] = \sum_{x:p_X(x)>0} g(x) \cdot p_X(x)$$

X is a discrete random variable with pmf

x	-1	0	1
$p_X(x)$	$1/4$	$1/2$	$1/4$

$Y = X^2$. Compute $E[Y]$.

X is a discrete random variable with pmf

x	-1	0	1
$p_X(x)$	$1/4$	$1/2$	$1/4$

$Y = X^2$. Compute $E[Y]$.

$$\begin{aligned}
 E[Y] &= \sum_{\text{all } x} x^2 \cdot p_X(x) \\
 &= (-1)^2 \times \frac{1}{4} + 0^2 \times \frac{1}{2} + 1^2 \times \frac{1}{4} \\
 &= \frac{1}{2}
 \end{aligned}$$

Properties of expected values

If a and b are constants, then

$$E[aX + b] = aE[X] + b$$

- Holds for all random variable X (not necessary discrete random variable).

Properties of expected values

If a and b are constants, then

$$E[aX + b] = aE[X] + b$$

- Holds for all random variable X (not necessary discrete random variable).
- Proof for discrete random variable: let $g(X) = aX + b$

Special cases of linear transformation $E[aX + b] = aE[X] + b$

- constant factor

$$E[aX] = aE[X]$$

- constant

$$E[b] = b$$

Recap

Expectation μ

- **For discrete random variable:** $E[X] = \sum_{\text{all } x} x \cdot p(x)$
- **Functions:** $E[g(X)] = \sum_{\text{all } x} g(x) p(x)$
- **Indicators:** $E[\delta_A] = P(A)$ where δ_A is an indicator function
- **Linear function:** $E[aX + b] = aE[X] + b$
- **Constants:** $E[c] = c$ if c is constant

Recap

Expectation μ

- **For discrete random variable:** $E[X] = \sum_{\text{all } x} x \cdot p(x)$
- **Functions:** $E[g(X)] = \sum_{\text{all } x} g(x) p(x)$
- **Indicators:** $E[\delta_A] = P(A)$ where δ_A is an indicator function
- **Linear function:** $E[aX + b] = aE[X] + b$
- **Constants:** $E[c] = c$ if c is constant

For two random variable's X and Y

$$E[X + Y] = E[X] + E[Y]$$