

# Chapter 7 part 2

## Properties of Expectations

Jingchen (Monika) Hu

Vassar College

MATH 241

# Outline

## 1 Moment generating functions

# Moment generating functions

## Definition

The moment generating function  $M(t)$  of the random variable  $X$  is defined for all real values of  $t$  by

$$M(t) = E[e^{tX}] = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ is discrete with pmf } p(x) \\ \int_{-\infty}^{\infty} e^{tx} f(x) dx & \text{if } X \text{ is continuous with pdf } f(x) \end{cases}$$

# Moment generating functions

## Definition

The moment generating function  $M(t)$  of the random variable  $X$  is defined for all real values of  $t$  by

$$M(t) = E[e^{tX}] = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ is discrete with pmf } p(x) \\ \int_{-\infty}^{\infty} e^{tx} f(x) dx & \text{if } X \text{ is continuous with pdf } f(x) \end{cases}$$

All the moments of  $X$  can be obtained by successively differentiating  $M(t)$  and then evaluating the result at  $t = 0$

# Moment generating functions

## Definition

The moment generating function  $M(t)$  of the random variable  $X$  is defined for all real values of  $t$  by

$$M(t) = E[e^{tX}] = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ is discrete with pmf } p(x) \\ \int_{-\infty}^{\infty} e^{tx} f(x) dx & \text{if } X \text{ is continuous with pdf } f(x) \end{cases}$$

All the moments of  $X$  can be obtained by successively differentiating  $M(t)$  and then evaluating the result at  $t = 0$

$$M'(t) = \frac{d}{dt} E[e^{tX}]$$

# Moment generating functions

## Definition

The moment generating function  $M(t)$  of the random variable  $X$  is defined for all real values of  $t$  by

$$M(t) = E[e^{tX}] = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ is discrete with pmf } p(x) \\ \int_{-\infty}^{\infty} e^{tx} f(x) dx & \text{if } X \text{ is continuous with pdf } f(x) \end{cases}$$

All the moments of  $X$  can be obtained by successively differentiating  $M(t)$  and then evaluating the result at  $t = 0$

$$M'(t) = \frac{d}{dt} E[e^{tX}] = E\left[\frac{d}{dt}(e^{tX})\right]$$

# Moment generating functions

## Definition

The moment generating function  $M(t)$  of the random variable  $X$  is defined for all real values of  $t$  by

$$M(t) = E[e^{tX}] = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ is discrete with pmf } p(x) \\ \int_{-\infty}^{\infty} e^{tx} f(x) dx & \text{if } X \text{ is continuous with pdf } f(x) \end{cases}$$

All the moments of  $X$  can be obtained by successively differentiating  $M(t)$  and then evaluating the result at  $t = 0$

$$M'(t) = \frac{d}{dt} E[e^{tX}] = E\left[\frac{d}{dt}(e^{tX})\right] = E[Xe^{tX}],$$

# Moment generating functions

## Definition

The moment generating function  $M(t)$  of the random variable  $X$  is defined for all real values of  $t$  by

$$M(t) = E[e^{tX}] = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ is discrete with pmf } p(x) \\ \int_{-\infty}^{\infty} e^{tx} f(x) dx & \text{if } X \text{ is continuous with pdf } f(x) \end{cases}$$

All the moments of  $X$  can be obtained by successively differentiating  $M(t)$  and then evaluating the result at  $t = 0$

$$M'(t) = \frac{d}{dt} E[e^{tX}] = E\left[\frac{d}{dt}(e^{tX})\right] = E[Xe^{tX}], M'(0) = E[X]$$



# Moment generating functions

## Definition

The moment generating function  $M(t)$  of the random variable  $X$  is defined for all real values of  $t$  by

$$M(t) = E[e^{tX}] = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ is discrete with pmf } p(x) \\ \int_{-\infty}^{\infty} e^{tx} f(x) dx & \text{if } X \text{ is continuous with pdf } f(x) \end{cases}$$

All the moments of  $X$  can be obtained by successively differentiating  $M(t)$  and then evaluating the result at  $t = 0$

$$M'(t) = \frac{d}{dt} E[e^{tX}] = E\left[\frac{d}{dt}(e^{tX})\right] = E[Xe^{tX}], M'(0) = E[X]$$

$$M''(t) = \frac{d}{dt} E[Xe^{tX}]$$

# Moment generating functions

## Definition

The moment generating function  $M(t)$  of the random variable  $X$  is defined for all real values of  $t$  by

$$M(t) = E[e^{tX}] = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ is discrete with pmf } p(x) \\ \int_{-\infty}^{\infty} e^{tx} f(x) dx & \text{if } X \text{ is continuous with pdf } f(x) \end{cases}$$

All the moments of  $X$  can be obtained by successively differentiating  $M(t)$  and then evaluating the result at  $t = 0$

$$M'(t) = \frac{d}{dt} E[e^{tX}] = E\left[\frac{d}{dt}(e^{tX})\right] = E[Xe^{tX}], M'(0) = E[X]$$

$$M''(t) = \frac{d}{dt} E[Xe^{tX}] = E\left[\frac{d}{dt}(Xe^{tX})\right]$$

# Moment generating functions

## Definition

The moment generating function  $M(t)$  of the random variable  $X$  is defined for all real values of  $t$  by

$$M(t) = E[e^{tX}] = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ is discrete with pmf } p(x) \\ \int_{-\infty}^{\infty} e^{tx} f(x) dx & \text{if } X \text{ is continuous with pdf } f(x) \end{cases}$$

All the moments of  $X$  can be obtained by successively differentiating  $M(t)$  and then evaluating the result at  $t = 0$

$$M'(t) = \frac{d}{dt} E[e^{tX}] = E\left[\frac{d}{dt}(e^{tX})\right] = E[Xe^{tX}], M'(0) = E[X]$$

$$M''(t) = \frac{d}{dt} E[Xe^{tX}] = E\left[\frac{d}{dt}(Xe^{tX})\right] = E[X^2 e^{tX}],$$

# Moment generating functions

## Definition

The moment generating function  $M(t)$  of the random variable  $X$  is defined for all real values of  $t$  by

$$M(t) = E[e^{tX}] = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ is discrete with pmf } p(x) \\ \int_{-\infty}^{\infty} e^{tx} f(x) dx & \text{if } X \text{ is continuous with pdf } f(x) \end{cases}$$

All the moments of  $X$  can be obtained by successively differentiating  $M(t)$  and then evaluating the result at  $t = 0$

$$M'(t) = \frac{d}{dt} E[e^{tX}] = E\left[\frac{d}{dt}(e^{tX})\right] = E[Xe^{tX}], M'(0) = E[X]$$

$$M''(t) = \frac{d}{dt} E[Xe^{tX}] = E\left[\frac{d}{dt}(Xe^{tX})\right] = E[X^2 e^{tX}], M''(0) = E[X^2]$$

# Moment generating functions

## Definition

The moment generating function  $M(t)$  of the random variable  $X$  is defined for all real values of  $t$  by

$$M(t) = E[e^{tX}] = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ is discrete with pmf } p(x) \\ \int_{-\infty}^{\infty} e^{tx} f(x) dx & \text{if } X \text{ is continuous with pdf } f(x) \end{cases}$$

All the moments of  $X$  can be obtained by successively differentiating  $M(t)$  and then evaluating the result at  $t = 0$

$$M'(t) = \frac{d}{dt} E[e^{tX}] = E\left[\frac{d}{dt}(e^{tX})\right] = E[Xe^{tX}], M'(0) = E[X]$$

$$M''(t) = \frac{d}{dt} E[Xe^{tX}] = E\left[\frac{d}{dt}(Xe^{tX})\right] = E[X^2 e^{tX}], M''(0) = E[X^2]$$

In general

$$M^n(t) = E[X^n e^{tX}], M^n(0) = E[X^n], n \geq 1$$

If  $X$  is a binomial random variable with parameters  $n$  and  $p$ . Use the moment generating functions to obtain its mean and variance.

If  $X$  is a binomial random variable with parameters  $n$  and  $p$ . Use the moment generating functions to obtain its mean and variance.

- The moment generating function is

$$\begin{aligned}M(t) = E[e^{tX}] &= \sum_{k=0}^n e^{tk} \binom{n}{k} p^k (1-p)^{n-k} \\&= \sum_{k=0}^n \binom{n}{k} (pe^t)^k (1-p)^{n-k} \\&= (pe^t + 1 - p)^n\end{aligned}$$

If  $X$  is a binomial random variable with parameters  $n$  and  $p$ . Use the moment generating functions to obtain its mean and variance.

- The moment generating function is

$$\begin{aligned}
 M(t) = E[e^{tX}] &= \sum_{k=0}^n e^{tk} \binom{n}{k} p^k (1-p)^{n-k} \\
 &= \sum_{k=0}^n \binom{n}{k} (pe^t)^k (1-p)^{n-k} \\
 &= (pe^t + 1 - p)^n
 \end{aligned}$$

- Taking differentiations

$$M'(t) = n(pe^t + 1 - p)^{n-1} pe^t$$



If  $X$  is a binomial random variable with parameters  $n$  and  $p$ . Use the moment generating functions to obtain its mean and variance.

- The moment generating function is

$$\begin{aligned}
 M(t) = E[e^{tX}] &= \sum_{k=0}^n e^{tk} \binom{n}{k} p^k (1-p)^{n-k} \\
 &= \sum_{k=0}^n \binom{n}{k} (pe^t)^k (1-p)^{n-k} \\
 &= (pe^t + 1 - p)^n
 \end{aligned}$$

- Taking differentiations

$$M'(t) = n(pe^t + 1 - p)^{n-1} pe^t$$

$$M''(t) = n(n-1)(pe^t + 1 - p)^{n-2} (pe^t)^2 + n(pe^t + 1 - p)^{n-1} pe^t$$

$$M'(t) = n(pe^t + 1 - p)^{n-1}pe^t$$

$$M''(t) = n(n-1)(pe^t + 1 - p)^{n-2}(pe^t)^2 + n(pe^t + 1 - p)^{n-1}pe^t$$

Therefore

$$E[X] = M'(0) = np$$

$$M'(t) = n(pe^t + 1 - p)^{n-1}pe^t$$

$$M''(t) = n(n-1)(pe^t + 1 - p)^{n-2}(pe^t)^2 + n(pe^t + 1 - p)^{n-1}pe^t$$

Therefore

$$E[X] = M'(0) = np$$

$$\begin{aligned} \text{Var}(X) = E[X^2] - (E[X])^2 &= M''(0) - (M'(0))^2 \\ &= n(n-1)p^2 + np - (np)^2 \\ &= np(1-p) \end{aligned}$$

# Properties of MGFs

- Property 1: The MGF of the sum of independent random variables equals to the product of the individual MGFs.

$$M_{X+Y}(t) = E[e^{t(X+Y)}] = E[e^{tX}e^{tY}] = E[e^{tX}]E[e^{tY}] = M_X(t)M_Y(t)$$

# Properties of MGFs

- Property 1: The MGF of the sum of independent random variables equals to the product of the individual MGFs.

$$M_{X+Y}(t) = E[e^{t(X+Y)}] = E[e^{tX}e^{tY}] = E[e^{tX}]E[e^{tY}] = M_X(t)M_Y(t)$$

- Property 2: The MGF uniquely determines the distribution. Refer to the tables of MGF of some discrete and continuous distributions.

Textbook pages 339 and 340 for lists of MGFs of distributions.

# Joint moment generating functions

## Definition

For any  $n$  random variables  $X_1, \dots, X_n$ , the joint MGF  $M(t_1, \dots, t_n)$  is defined for all real values of  $t_1, \dots, t_n$  by

$$M(t_1, \dots, t_n) = E[e^{t_1 X_1 + \dots + t_n X_n}]$$

# Joint moment generating functions

## Definition

For any  $n$  random variables  $X_1, \dots, X_n$ , the joint MGF  $M(t_1, \dots, t_n)$  is defined for all real values of  $t_1, \dots, t_n$  by

$$M(t_1, \dots, t_n) = E[e^{t_1 X_1 + \dots + t_n X_n}]$$

- The individual MGF can be obtained from  $M(t_1, \dots, t_n)$  by letting all but one of the  $t_j$ 's be 0

$$M_{X_i}(t) = E[e^{tX_i}] = M(0, \dots, t, 0, \dots, 0)$$

# Joint moment generating functions

## Definition

For any  $n$  random variables  $X_1, \dots, X_n$ , the joint MGF  $M(t_1, \dots, t_n)$  is defined for all real values of  $t_1, \dots, t_n$  by

$$M(t_1, \dots, t_n) = E[e^{t_1 X_1 + \dots + t_n X_n}]$$

- The individual MGF can be obtained from  $M(t_1, \dots, t_n)$  by letting all but one of the  $t_j$ 's be 0

$$M_{X_i}(t) = E[e^{tX_i}] = M(0, \dots, t, 0, \dots, 0)$$

- The joint MGF  $M(t_1, \dots, t_n)$  uniquely determines the joint distribution of  $X_1, \dots, X_n$



# Joint moment generating functions

## Definition

For any  $n$  random variables  $X_1, \dots, X_n$ , the joint MGF  $M(t_1, \dots, t_n)$  is defined for all real values of  $t_1, \dots, t_n$  by

$$M(t_1, \dots, t_n) = E[e^{t_1 X_1 + \dots + t_n X_n}]$$

- The individual MGF can be obtained from  $M(t_1, \dots, t_n)$  by letting all but one of the  $t_j$ 's be 0

$$M_{X_i}(t) = E[e^{tX_i}] = M(0, \dots, t, 0, \dots, 0)$$

- The joint MGF  $M(t_1, \dots, t_n)$  uniquely determines the joint distribution of  $X_1, \dots, X_n$
- Then we have the  $n$  random variables  $X_1, \dots, X_n$  are independent **if and only if**

$$M(t_1, \dots, t_n) = M_{X_1}(t_1) \dots M_{X_n}(t_n)$$