ROXYBOND™ (oxycodone hydrochloride) immediate release tablets CII 5mg•10mg•15mg•30mg

Eligible patients may save with the ROXYBOND™ Copay Card

Pay as little as $10* for each prescription

ROXYBOND™ (oxycodone hydrochloride) immediate release tablets CII 5mg•10mg•15mg•30mg

BIN: 610600
PCN: AS
GRP: 507
ID: 50710398622

***Offer valid for commercially insured patients only. Maximum savings limit applies; patient out-of-pocket expense may vary. I am in agreement with the Program Terms and Conditions received with the Copay Card, which include additional exclusions. Use of this Copay Card confirms that I meet all offer eligibility criteria. This offer is not insurance.**

Please see additional Important Safety Information, and full Prescribing Information, including BOXED WARNINGS and Medication Guide at www.protegapharma.com prior to taking ROXYBOND.

PHARMACIST INSTRUCTIONS: SUBMITTING CLAIM

REMOVE ANY DISCOUNT CARDS FROM TRANSACTION. USING DISCOUNT CARDS WITH COPAY CARD WILL RESULT IN A BENEFIT REJECTION AND POSSIBLY A HIGHER COST FOR THE PATIENT.

1. Submit claim to patient's insurance first.
2. Then submit a secondary claim to AlphaScrip as a coordination of benefits using **OCC 8**.

If the patient's insurance requires a Prior Authorization:

- Returns a reject 75, or
- Drug not covered reject 70

1. **Initiate Prior Authorization with prescriber immediately**. Copay card allows only one 30-day script per calendar year while PA processes, then up to $500/prescription until annual maximum is reached.
2. Submit the secondary claim to AlphaScrip as a coordination of benefits using **OCC 3**.
3. By utilizing the OCC 3, you give AlphaScrip permission to submit a Prior Authorization request to the prescriber, if appropriate.

Pharmacies needing assistance with processing this copay card, please contact the AlphaScrip Pharmacy Helpdesk at 877-274-3244.

INDICATION AND USAGE

- ROXYBOND (oxycodone hydrochloride) tablets, for oral use, CII, is a strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain severe enough to require an opioid pain medicine, when other pain treatments such as non-opioid pain medicines do not treat your pain well enough or you cannot tolerate them.
- ROXYBOND is an opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse, and misuse that can lead to death.

IMPORTANT SAFETY INFORMATION ABOUT ROXYBOND

WARNING: SERIOUS LIFE-THREATENING RISKS FROM USE OF ROXYBOND

Addiction, Abuse, and Misuse

Because the use of ROXYBOND exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death, assess each patient's risk prior to prescribing and reassess all patients regularly for the development of these behaviors and conditions.

Please see additional Important Safety Information, and full Prescribing Information, including BOXED WARNINGS and Medication Guide at www.protegapharma.com prior to taking ROXYBOND. Speak with your healthcare provider if you have additional questions about ROXYBOND.

For additional information, please visit www.copayroxybond.com

©2024 Protega Pharma. All rights reserved.

IMPORTANT SAFETY INFORMATION (continued)

Life-Threatening Respiratory Depression
Serious, life-threatening, or fatal respiratory depression may occur with use of ROXYBOND, especially during initiation or following a dosage increase. To reduce the risk of respiratory depression, proper dosing and titration of ROXYBOND are essential.

Accidental Ingestion
Accidental ingestion of even one dose of ROXYBOND, especially by children, can result in a fatal overdose of oxycodone.

Risks From Concomitant Use With Benzodiazepines Or Other CNS Depressants
Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing of ROXYBOND and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.

Neonatal Opioid Withdrawal Syndrome (NOWS)
If opioid use is required for an extended period of time in a pregnant woman, advise the patient of the risk of NOWS, which may be life-threatening if not recognized and treated. Ensure that management by neonatology experts will be available at delivery.

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)
Healthcare providers are strongly encouraged to complete a REMS-compliant education program and to counsel patients and caregivers on serious risks, safe use, and the importance of reading the Medication Guide with each prescription.

Cytochrome P450 3A4 Interaction
The concomitant use of ROXYBOND with all cytochrome P450 3A4 inhibitors may result in an increase in oxycodone plasma concentrations, which could increase or prolong adverse reactions and may cause potentially fatal respiratory depression. In addition, discontinuation of a concomitantly used cytochrome P450 3A4 inducer may result in an increase in oxycodone plasma concentration. Regularly evaluate patients receiving ROXYBOND and any CYP3A4 inhibitor or inducer.

Important information about ROXYBOND:

- **Get emergency help or call 911 right away if you take too much ROXYBOND (overdose)**. When you first start taking ROXYBOND, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur. Talk to your healthcare provider about naloxone, a medicine for the emergency treatment of an opioid overdose.
- Taking ROXYBOND with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.
- Never give anyone else your ROXYBOND. They could die from taking it. Selling or giving away ROXYBOND is against the law.
- Store ROXYBOND securely, out of sight and reach of children and in a location not accessible by others, including visitors to the home.

Do not take ROXYBOND if you have:

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or have narrowing of the stomach or intestines.
- allergy to oxycodone.

Before taking ROXYBOND, tell your healthcare provider if you have a history of:

- head injury, seizures
- problems urinating
- liver, kidney, thyroid problems
- pancreas or gallbladder problems
- abuse of street or prescription drugs, alcohol addiction, opioid overdose or mental health problems.

Please see additional Important Safety Information, and full Prescribing Information, including BOXED WARNINGS and Medication Guide at www.protegapharma.com prior to taking ROXYBOND. Speak with your healthcare provider if you have additional questions about ROXYBOND.

Tell your healthcare provider if you are:

- **noticing your pain getting worse**. If your pain gets worse after you take ROXYBOND, do not take more of ROXYBOND without first talking to your healthcare provider. Talk to your healthcare provider if the pain that you have increases, if you feel more sensitive to pain, or if you have new pain after taking ROXYBOND.
- **pregnant or planning to become pregnant**. Use of ROXYBOND for an extended period of time during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **breastfeeding**. ROXYBOND passes into breast milk and may harm your baby. Carefully observe infants for increased sleepiness (more than usual), breathing difficulties, or limpness. Seek immediate medical care if you notice these signs.
- living in a household where there are small children or someone who has abused street or prescription drugs.
- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking ROXYBOND with certain other medicines can cause serious side effects that could lead to death.

When taking ROXYBOND:

- Do not change your dose. Take ROXYBOND exactly as prescribed by your healthcare provider. Use the lowest dose possible for the shortest time needed.
- For acute (short-term) pain, you may only need to take ROXYBOND for a few days. You may have some ROXYBOND left over that you did not use. See disposal information at the bottom of this section for directions on how to safely throw away (dispose of) your unused ROXYBOND.
- Take your prescribed dose every 4 to 6 hours as needed for pain. Do not take more than your prescribed dose. If you miss a dose, take your next dose at your usual time.
- Call your healthcare provider if the dose you are taking does not control your pain.
- If you have been taking ROXYBOND regularly, do not stop taking ROXYBOND without talking to your healthcare provider.
- Dispose of expired, unwanted, or unused ROXYBOND by promptly flushing down the toilet, if a drug take-back option is not readily available. Visit www.fda.gov/drugdisposal for additional information on disposal of unused medicines.

While taking ROXYBOND DO NOT:

- Drive or operate heavy machinery, until you know how ROXYBOND affects you. ROXYBOND can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with ROXYBOND may cause you to overdose and die.

The possible side effects of ROXYBOND are:

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe.

Get emergency medical help or call 911 right away if you have:

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion.

These are not all the possible side effects of ROXYBOND. Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. For more information go to dailymed.nlm.nih.gov.

Please see additional Important Safety Information, and full Prescribing Information, including BOXED WARNINGS and Medication Guide at www.protegapharma.com prior to taking ROXYBOND. Speak with your healthcare provider if you have additional questions about ROXYBOND.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit MedWatch or call 1-800-FDA-1088. To report SUSPECTED ADVERSE REACTIONS, contact Protega Pharmaceuticals Inc. at 1-844-798-3610 or FDA at 1-800-FDA-1088 or www.fda.gov/safety.

ROXYBOND (oxycodone hydrochloride) Copay Card Program Terms and Conditions

The ROXYBOND Copay Card Program (the "Card") may cover a portion of eligible out-of-pocket costs of patients' ROXYBOND prescriptions. Patient eligibility for the Card is subject to the following terms and conditions:

- The Card may only be used by eligible residents of the U.S., Puerto Rico, or U.S. territories at participating eligible pharmacies located in the U.S., Puerto Rico, or U.S. territories. Card user must be at least 18 years of age.
- The Card will cover a portion of the out-of-pocket costs of an eligible patient's ROXYBOND prescriptions (Card benefit is calculated at the time of use and is subject to vary). Annual maximum benefit of $8,500 applies.
- The Card is only available for use with a valid prescription. The Card has no cash value, is not transferable, may not be reproduced, and may not be sold, purchased, traded, or offered for sale, purchase, or trade.
- THIS CARD IS NOT INSURANCE AND IS NOT INTENDED TO BE A SUBSTITUTE FOR INSURANCE. The Card is not valid for prescriptions eligible for full reimbursement by insurance plans or any other pharmacy benefit or assistance programs, and it will not reimburse any payments made by a Flexible Spending Account ("FSA"), Health Savings Account ("HSA"), or Health Reimbursement Account ("HRA").
- This Card is only valid for use by patients with commercial insurance. THE CARD IS NOT VALID FOR PRESCRIPTIONS THAT ARE ELIGIBLE TO BE REIMBURSED, IN WHOLE OR IN PART, BY:
  - ANY FEDERAL HEALTH CARE PROGRAM (AS DEFINED AT 42 U.S.C. §1320a-7b(f)), INCLUDING MEDICAID, MEDICARE (INCLUDING ANY MEDICARE ADVANTAGE OR MEDICARE PART D PLAN, WHICH INCLUDES MEDICARE PART D ENROLLEES IN THE PRESCRIPTION DRUG COVERAGE GAP), TRICARE, CHAMPUS, PUERTO RICO GOVERNMENT HEALTH INSURANCE PLAN, OR ANY OTHER STATE OR FEDERAL HEALTH CARE PROGRAM;
  - MEDICARE-ELIGIBLE PATIENTS ENROLLED IN AN EMPLOYER-SPONSORED RETIREE HEALTH PLAN OR PRESCRIPTION DRUG PLAN;
  - CASH-PAYING PATIENTS; OR
  - PATIENTS RESIDING IN A STATE WHERE THE USE OF THE CARD IS PROHIBITED BY LAW OR REGULATION.
- Submit claim to patient's insurance first, then submit a secondary claim to AlphaScrip as a coordination of benefits using OCC 8. If the patient's insurance requires a Prior Authorization and returns a reject 75, or drug not covered reject 70, submit a secondary claim to AlphaScrip as a coordination of benefits using OCC 3. By utilizing the OCC 3, you give AlphaScrip permission to submit a Prior Authorization request to the prescriber, if appropriate.
- Patient, pharmacist, and prescriber agree not to submit a claim for reimbursement for any value conveyed by Card. Patient and pharmacist must each individually report receipt of Card to any insurer, health plan, or any other party that covers or reimburses any part of the prescription filled using this Card, as required by law, regulation, or company policy.
- Certain information necessary to facilitate the use of the Card will be shared with Protega Pharma and its affiliates.
- Card is not valid in the state of Massachusetts.
- Card is void where taxed or where prohibited or restricted by law.
- Protega Pharma reserves the right to terminate the Card program or to rescind, revoke, or amend this offer without notice.

Please see additional Important Safety Information, and full Prescribing Information, including BOXED WARNINGS and Medication Guide at www.protegapharma.com prior to taking ROXYBOND. Speak with your healthcare provider if you have additional questions about ROXYBOND.

PROTEGA PHARMA

©2024 Protega Pharmaceuticals Inc.
ROXYBOND, the ROXYBOND logo, and the Protega Pharma logo are trademarks of Protega Pharmaceuticals Inc. PM-FM-1009 r8 Nov 2024