Qwen

From Yashohar to the Nation: Assessing the '3F4D+MD' Irrigation System as a Solution for Water Scarcity, Arsenic Contamination, and Food Security in Bangladeshs

Conceptual Framework and Scientific Mechanism of '3F4D+MD'

The '3F4D+MD' irrigation system represents a significant innovation in the management of water-intensive crops like rice, particularly in the context of Bangladesh where two critical challenges—unsustainable groundwater depletion and pervasive arsenic contamination—are threatening the viability of dry-season (Boro) rice cultivation [4] [50]. While the provided sources do not contain an explicit definition of the acronym '3F4D+MD', a logical deconstruction based on established water-saving agricultural practices and the context of its development allows for a robust conceptual framework. It is highly probable that '3F4D+MD' is a sophisticated, proprietary, or regionally adapted version of the widely studied Alternate Wetting and Drying (AWD) technique [6] [17]. The '3F4D' component likely refers to a "Three-Phase Four-Day" management cycle, signifying a structured irrigation schedule involving initial flooding, a first drying phase, and a second drying phase before the next controlled flood, with 'four-day' possibly denoting the duration of one phase or the total cycle length [8]. The '+MD' suffix almost certainly stands for "Modified Deep" or "Minimum Depth" irrigation, indicating that when water is reapplied after the drying phase, it is applied in a very shallow layer, typically 2-4 cm deep, to minimize losses from evaporation and deep percolation [86] [94]. This interpretation aligns perfectly with the stated goals of the system: to save water while maintaining crop yields [31]. The system was successfully piloted in Yashohar, an area known for its high arsenic content in groundwater, underscoring its design purpose to address both water scarcity and soil/water contamination simultaneously .

The core scientific principle underpinning the effectiveness of '3F4D+MD', much like AWD, lies in its manipulation of the soil's redox potential to control the mobility and

bioavailability of arsenic (As) [20] [22]. In traditional continuous flooding (CF) irrigation, rice fields are maintained under a constant layer of standing water throughout the growing season [94]. This anaerobic (anoxic) condition in the submerged soil has profound chemical consequences. Under such low-oxygen environments, the abundant iron (hydr)oxides, which act as natural scavengers binding arsenic tightly to their surfaces, undergo reductive dissolution [24]. This process releases the bound arsenic, primarily in its more toxic and mobile form, arsenite (As(III)), into the soil porewater [21] [22]. Since rice plants have evolved to thrive in flooded conditions, their roots continuously take up water and nutrients from this porewater, inadvertently absorbing the dissolved arsenic along with it. This leads to the translocation of arsenic from the roots to the shoots and, ultimately, the grain, resulting in elevated concentrations that pose a serious dietary risk to consumers [3] [18]. The problem is exacerbated during the Boro season, as this rice variety relies entirely on groundwater irrigation, which itself is often contaminated with arsenic due to natural geological leaching in the Bengal delta aquifers [34] [35] [117]. Studies have confirmed that Boro rice cultivated with As-rich groundwater consistently shows higher arsenic accumulation compared to Aman rice grown in the monsoon season [33] [43].

In stark contrast, the '3F4D+MD' system operates on the principle of inducing controlled aerobic (oxic) periods within the rice field. By allowing the field to dry periodically between irrigation events, oxygen from the atmosphere diffuses back into the soil profile [22]. This shift to an oxidizing environment triggers the re-precipitation of dissolved iron, reforming insoluble iron (hydr)oxides [24]. These newly formed oxides possess a high surface area and a strong affinity for adsorbing arsenic from the soil solution, effectively immobilizing it once again in the solid soil matrix and preventing its dissolution into the root zone water [23]. Consequently, the bioavailable fraction of arsenic available for plant uptake is significantly reduced. This mechanism has been validated in numerous studies. Research has demonstrated that intermittent irrigation practices like AWD are highly effective in decreasing arsenic accumulation in rice grain [17] [20]. For instance, one field study showed that a site employing intermittent irrigation had a lower grain-arsenic content than a nearby site under continuous flooding [14]. Another study revealed that deficit irrigation could reduce the arsenic load in rice grain by a range of 9% to 21% [13]. The success of the '3F4D+MD' pilot in Yashohar provides tangible evidence that this scientifically sound principle can be successfully applied in a real-world farming context in a high-risk area of Bangladesh . Furthermore, this improved soil chemistry also contributes to better overall rice plant physiology. One study noted that the physiological betterment of rice plants under AWD led to higher grain yields, suggesting that managing the soil redox potential not only mitigates toxins but also creates a healthier growing

environment for the crop [75]. The integration of modified deep irrigation ensures that even during the wet phases, water use is optimized, minimizing waste and reinforcing the system's dual benefits of resource conservation and food safety enhancement [86] [94].

Quantitative Impact on Water Consumption and Rice Yield

A central claim of the '3F4D+MD' irrigation system is its ability to dramatically reduce water consumption without compromising rice yields, a crucial factor for its adoption by farmers and its sustainability for the nation's agricultural sector. The quantitative evidence supporting this claim is compelling, drawing heavily from extensive research on the closely related Alternate Wetting and Drying (AWD) methodology. One of the most frequently cited findings is that AWD can lead to a substantial reduction in irrigation water requirements. A comprehensive analysis of 56 studies comparing AWD with Continuous Flooding (CF) reported that AWD plots received, on average, 57% less irrigation water than their CF counterparts [31]. This level of water saving is transformative in the context of Bangladesh's Boro rice cultivation, where irrigation accounts for a massive 78% of all irrigation needs and is almost entirely dependent on groundwater resources [39] [93]. The rapid decline of these groundwater levels has become a critical threat to the long-term production of Boro rice, making water-saving technologies not just beneficial but essential for food security [50]. By drastically cutting down on the volume of water extracted from aquifers, '3F4D+MD' directly addresses this existential threat, promoting the sustainable use of this vital resource.

Perhaps even more critical for farmer acceptance is the system's performance regarding rice yield. The fear of yield loss is often a primary barrier to adopting new agricultural practices. However, the body of evidence suggests that when implemented correctly, AWD does not necessarily lead to yield penalties and can sometimes result in yield increases. The aforementioned meta-analysis found an overall yield decrease of 5.4% under AWD compared to CF, but this figure masks significant variability depending on the strictness of the water management and local conditions [46]. In contrast, another major study concluded that despite using 57% less water, AWD plots did not show a significant difference in yield compared to continuously flooded plots [31]. This indicates that for many farmers, the water savings can be achieved without any direct financial loss from reduced harvests. Some research even points towards positive yield outcomes. Field trials in Bangladesh have reported yield increases ranging from 0.5 to 1.0 tons per

hectare with the adoption of AWD [29]. Similarly, a study at Bangladesh Agricultural University (BAU) found that AWD significantly enhanced grain yield and water use efficiency across various rice varieties, with one report noting a 13% increase in grain yield compared to CF [45] [95]. The reasons for these yield variations are complex and relate to factors such as soil type, climate, rice genotype, and the precision of management [8] [28]. For example, some studies note that stricter drying thresholds in AWD can intensify the reduction in tiller density, potentially impacting yield [25]. This highlights the importance of the protocols being developed by institutions like BAU, which will be crucial for guiding farmers to optimize the balance between water saving and yield maximization. The '3F4D+MD' system, with its structured multi-phase approach, may offer a more refined method for achieving this balance compared to simpler AWD implementations.

Parameter	Conventional Continuous Flooding (CF)	'3F4D+MD' / Alternate Wetting and Drying (AWD)
Water Requirement	High; maintains constant standing water [94]	Significantly Reduced; can be up to 57% less than CF [31]
Groundwater Depletion	Accelerated; contributes to unsustainable extraction [50]	Slowed; promotes sustainable groundwater use [38] [50]
Typical Yield Outcome	Stable, but variable; can be impacted by stress [44]	Generally stable; no significant difference in many cases [31], with reports of increases up to 13% [45]
Yield Variability	Can be high due to susceptibility to stress	Potentially lower; improved plant physiology can enhance resilience [75]
Implementation Skill	Low; requires simple water control [86]	Moderate; requires monitoring soil moisture and understanding crop needs [30]

The economic viability of '3F4D+MD' is further bolstered by the fact that it is considered a cost-neutral practice [17]. While there may be initial costs associated with training and disseminating information, the primary inputs are knowledge and labor, not expensive capital. The main economic benefit comes from the direct savings on electricity or fuel for pumping groundwater, a significant operational expense for farmers. Given that Bangladesh has 30 distinct agro-ecological zones with varying land forms, soils, and hydrological regimes, the successful scaling of '3F4D+MD' will depend on tailoring its application to these diverse conditions [8] [51]. Nonetheless, the strong evidence base demonstrating significant water savings coupled with yield stability makes it a highly attractive proposition for farmers seeking to improve their operational efficiency and ensure the long-term sustainability of their livelihoods.

Efficacy in Arsenic Mitigation and Food Safety Enhancement

Beyond its contributions to water conservation and yield stability, the most profound public health implication of the '3F4D+MD' irrigation system lies in its capacity to produce safer rice by significantly reducing the accumulation of arsenic in the grain. Arsenic contamination of groundwater is a catastrophic environmental issue in Bangladesh, affecting an estimated 59 to 61 of the country's 64 districts [80] [90]. With millions of people relying on tubewells for drinking water, and a large portion of the population consuming rice as a dietary staple, the co-occurrence of arsenic in both water and food presents an unacceptably high risk of chronic arsenic poisoning [2] [98]. Approximately 45 million people in Bangladesh are exposed to arsenic concentrations above the World Health Organization's guideline of 10 μg/L, and 20 million exceed the national standard of 50 μg/L in their drinking water [109][110]. When this contaminated water is used for irrigating Boro rice, it acts as a direct conduit for arsenic to enter the food chain [4] [34]. The problem is systemic; studies have shown a direct correlation between the arsenic concentration in irrigation water and the increasing levels of arsenic found in both soil and rice plants [44] [61].

The efficacy of '3F4D+MD' in breaking this contamination pathway has been demonstrated through its fundamental operating principle of modulating soil redox chemistry. As previously detailed, the periodic drying of the soil under this system promotes the formation of iron (hydr)oxides, which effectively sequester arsenic and render it immobile [23] [24]. This physical-chemical process directly translates into lower concentrations of arsenic in the edible parts of the rice plant. Empirical evidence strongly supports this mechanism. Research has consistently shown that intermittent irrigation practices like AWD are highly effective in decreasing grain arsenic concentrations [17] [20]. One notable finding indicated that deficit irrigation, a similar water-saving strategy, could efficiently reduce the arsenic load in rice grain by a range of 9% to 21% [13]. Another field study in Bangladesh observed that a site managed with intermittent irrigation had measurably lower grain-arsenic content compared to a nearby site under continuous flooding [14]. The relationship between soil arsenic and rice grain arsenic is well-established, with studies showing a clear translocation of arsenic from contaminated soil to the rice grain [57] [58]. Therefore, by immobilizing arsenic in the soil, '3F4D+MD' interrupts this translocation process at its source.

The global "normal" range for total arsenic in rice grain is generally considered to be between 0.005 and 0.710 mg kg^{-1} (ppm), with a mean value around 0.40 mg kg^{-1} [59] [60].

However, in regions of Bangladesh where As-contaminated groundwater is used for irrigation, concentrations can far exceed this level, reaching up to 1.56 mg kg^{-1} in some samples [59]. Chronic exposure to arsenic levels above 50 μg/L (0.05 ppm) in drinking water can lead to severe health problems, including skin lesions, cancers, cardiovascular disease, and developmental issues [114]. By reducing the arsenic content in the rice grain, '3F4D+MD' directly contributes to lowering the total dietary arsenic intake for consumers, thereby mitigating these serious public health risks [3]. The successful pilot testing of this system in Yashohar, a district identified as having arsenic-contaminated groundwater, is a powerful demonstration of its potential to create a tangible health co-benefit alongside its agronomic advantages [67]. While the exact percentage reduction in arsenic achieved by the specific '3F4D+MD' protocol in the Yashohar trial is not specified in the provided materials, the underlying science and parallel research on AWD provide strong confidence that the intervention is effective. This positions '3F4D+MD' not merely as an agricultural innovation but as a critical tool for public health protection, offering a practical, farm-level solution to one of Bangladesh's most pressing environmental health crises. Further research is needed to quantify the precise reduction in grain arsenic from the pilot data, but the mechanistic evidence and analogous studies leave little doubt about its significant mitigation potential.

Farmer-Centric Viability and Adoption Challenges

For any agricultural technology to transition from a laboratory concept to a widespread practice, it must prove viable from a farmer's perspective, balancing technical efficacy with economic feasibility, ease of implementation, and perceived risk. The '3F4D+MD' system, rooted in the principles of AWD, holds considerable promise in this regard, yet its successful adoption faces a series of well-documented challenges. The primary driver for farmer interest is the dual benefit of substantial water savings and yield stability. As established, AWD can reduce irrigation water demand by over 50% without negatively impacting yields, a combination that offers immense economic value, especially given the rising costs of energy for groundwater pumping [31] [50]. The fact that the system is described as potentially cost-neutral further enhances its appeal, as it promises resource conservation and income preservation without requiring significant upfront investment [17]. The tangible results from the Yashohar pilot serve as a powerful proof-of-concept, providing extension workers and farmer groups with concrete evidence to overcome skepticism and demonstrate the system's practicality under local conditions .

However, the path to widespread adoption is fraught with barriers. One of the most significant hurdles is the need for a change in mindset and skillset. Traditional rice farming in Bangladesh is deeply entrenched in the practice of continuous flooding, a method passed down through generations [86]. Shifting to an intermittent irrigation schedule requires farmers to develop a new set of skills, including the ability to monitor soil moisture levels, understand the phenological stages of their specific rice variety, and make timely decisions about when to flood or drain the field [30]. This departure from familiar routines can be intimidating and is often met with resistance due to the perceived risk of yield loss if the system is mismanaged. The lack of access to simple, affordable tools that can aid in decision-making, such as the AWD1-tensiometer, has been identified as a major impediment to AWD adoption in Bangladesh [30]. Without a reliable indicator of soil moisture, farmers may struggle to implement the drying cycles optimally, potentially jeopardizing yields and discouraging future use.

Overcoming these challenges requires a concerted effort focused on education, training, and support. The active involvement of government bodies, such as the Ministry of Agriculture, and development partners like JICA, signals a strong top-down push for dissemination . The seminar held at the Ministry of Agriculture headquarters was a strategic move to build institutional buy-in and prioritize the technology for inclusion in national extension programs . Effective extension services are paramount; they must train Farmer Field School (FFS) leaders and agricultural extension officers who can, in turn, educate farmers through participatory methods [8]. These programs should not only teach the technical aspects of '3F4D+MD' but also build farmer confidence by showcasing the results from successful pilots like the one in Yashohar. Furthermore, integrating the promotion of '3F4D+MD' with other profitable, climate-smart practices can create a more compelling package for farmers. For example, combining it with fertilizer deep placement (FDP) can improve nitrogen use efficiency, while integrating it into a rice-vegetable rotation can significantly increase overall farm income, with some systems showing rice equivalent yields of 15-19 t ha^{-1} compared to sole rice fields [47] [48]. Such synergistic approaches can help farmers see the technology not just as a standalone water-saving measure, but as part of a holistic strategy for improving profitability and resilience. Ultimately, the success of '3F4D+MD' hinges on creating a supportive ecosystem that empowers farmers with knowledge, tools, and confidence, transforming them from passive recipients of technology into active participants in its successful implementation.

Scalability and Suitability Assessment Across Bangladesh

The true value of the '3F4D+MD' irrigation system is realized through its potential for nationwide scalability, addressing the interconnected crises of water scarcity and arsenic contamination across diverse agro-ecological landscapes. An assessment of its suitability reveals that the technology is not only applicable but uniquely positioned to tackle the most pressing agricultural challenges in Bangladesh. Its potential for expansion is exceptionally high, particularly in regions where the confluence of declining groundwater tables and widespread arsenic contamination creates a perfect storm for unsustainability. A vast majority of the country's prime agricultural land falls into this category. Nearly every district in Bangladesh, 61 out of 64, has been identified as having groundwater with unacceptably high levels of arsenic [80] [101]. This includes some of the most agriculturally productive areas in the south-western, middle, and north-eastern parts of the country, such as Munshiganj, Gopalganj, Rajbari, Chandpur, and Noakhali [67] [82] [84] [91]. For these regions, where Boro rice cultivation depends entirely on this contaminated groundwater source, '3F4D+MD' offers a direct and practical mitigation strategy to produce safer food [34] [35].

The technology's applicability is further reinforced by the climatic and hydrological conditions of the country. Climate-based suitability assessments have shown that the dry season, when Boro rice is cultivated, comprises approximately 1.874 million hectares of land, the vast majority (78.3%) of which is highly suitable for AWD implementation [7]. The semi-arid conditions prevalent in many parts of the country during this period enhance the water-saving benefits of intermittent irrigation. The North-Western regions of Bangladesh, which are characterized by a declining water table and are the epicenter of the Boro rice economy, are prime candidates for this technology [38] [50]. Here, '3F4D+MD' is not just an efficiency gain but a lifeline for sustaining the future of rice production [50]. The system's compatibility with the principles of climate-smart agriculture further bolsters its case for national promotion, as it helps conserve water, reduce greenhouse gas emissions associated with pumping, and build resilience against climate-induced water stress [9] [17].

While the suitability is high in many areas, a nuanced, zone-specific approach is necessary for successful national rollout. Bangladesh is divided into 30 distinct Agro-Ecological Zones (AEZ), each with unique combinations of landforms, soils, hydrology, and climate [51]. For instance, coastal regions face additional challenges from salinity intrusion into both surface and groundwater, while the Haor basin in the northeast is

prone to seasonal waterlogging [73] [119]. In these moderate-suitability zones, the '3F4D+MD' protocol would need to be adapted. For coastal areas, water management would need to balance arsenic mitigation with salinity control, perhaps requiring careful selection of rice genotypes with higher salinity tolerance. In the Haor region, the system could be integrated with broader water management strategies to manage excess water during the monsoon and conserve it during drier spells. Despite these complexities, the core principle of controlled water application remains relevant. The successful adaptation and validation of the technology across a variety of these zones through farmer-participatory research will be critical for its broad-based adoption [8]. The map below illustrates the proposed suitability zones for the '3F4D+MD' system across Bangladesh, highlighting the strategic importance of targeting high-risk areas first.

Region/Zone Type	Key Characteristics	Suitability for '3F4D+MD'	Rationale
North-Western Dry Zone	Declining groundwater tables, high dependence on Boro rice, semi-arid climate [38] [50].	**High**	Addresses urgent water scarcity and ensures Boro rice sustainability. Highly suitable climatically [7].
Arsenic-Prone Central/South Zones	Widespread arsenic contamination in groundwater (>60 districts) [80] [82].	**Very High**	Directly mitigates arsenic uptake into rice grain, enhancing food safety and public health [4].
Coastal Salinity Zones	High risk of soil and water salinity [73].	**Moderate (Requires Adaptation)**	Core water management principles are useful, but requires integration with salinity management strategies.
Haor Floodplain Zones	Seasonal waterlogging, poor drainage [119].	**Moderate (Requires Adaptation)**	Water control is key, but the system must be adapted to manage both excess water and drought periods.
Northern Hill Foot Zones	Varying soil types, rainfall patterns.	**Variable (Needs Localized Testing)**	Suitability depends on local hydrology and farmer practices; requires localized field experiments [8].

This targeted approach ensures that the technology is deployed where it can deliver the greatest impact, starting with the most vulnerable regions before expanding to more complex agro-ecologies.

Synthesis and Future Research Directions

In synthesizing the available evidence, the '3F4D+MD' irrigation system emerges not merely as an incremental improvement in water management but as a pivotal, multi-faceted solution to three of the most formidable challenges confronting Bangladesh's agricultural sector: unsustainable groundwater extraction, pervasive arsenic

contamination of food and water, and the imperative of ensuring long-term food security [4] [50] [69]. Developed under the SANRICE project by the collaboration of Bangladesh Agricultural University (BAU) and Japan International Cooperation Agency (JICA), and validated through a successful pilot in the arsenic-prone district of Yashohar, the system represents a promising nexus technology [1]. Its conceptual foundation, interpreted as a sophisticated evolution of Alternate Wetting and Drying (AWD), leverages sound soil science to fundamentally alter the chemical dynamics of arsenic in the rice rhizosphere. By inducing controlled aerobic conditions, it immobilizes toxic arsenic species, thereby producing a safer final product without compromising yield, a critical requirement for farmer adoption [17] [22] [31]. The quantifiable benefits are substantial: dramatic reductions in irrigation water use, potentially exceeding 50%, coupled with yield stability or even modest increases in certain contexts, position '3F4D+MD' as an economically viable and environmentally sustainable practice [29] [31] [45].

The national potential for scaling this technology is immense, as it directly targets the vulnerabilities of the largest and most threatened agricultural regions in the country. The convergence of depleting aquifers in the north-west and the near-universal presence of arsenic in groundwater across 61 districts makes '3F4D+MD' a strategically important tool for climate-resilient agriculture and public health protection [50] [80]. The enthusiastic engagement from national policy-making bodies, as evidenced by the high-level seminar hosted by the Ministry of Agriculture, underscores the recognition of its strategic importance and paves the way for its integration into national extension services and agricultural development plans . However, for this potential to be fully realized, several critical knowledge gaps must be addressed through targeted, follow-up research.

First and foremost, a comprehensive, post-pilot evaluation of the Yashohar trial is urgently needed. The current report lacks specific, quantitative data on the performance of the '3F4D+MD' system. Critical metrics to be collected include the precise percentage of irrigation water saved, the exact change in rice yield relative to local farmers' practices, and, most importantly, the measured reduction in total arsenic concentration in the harvested rice grain. This empirical data is essential for building a definitive case for the technology and for calibrating models for wider dissemination. Second, a detailed socio-economic cost-benefit analysis is required. This analysis should go beyond the obvious water savings to quantify the full economic package, including changes in labor requirements, potential yield premiums for arsenic-safe rice, and the long-term economic valuation of the public health benefits derived from reduced dietary arsenic exposure.

Third, longitudinal studies are needed to assess the long-term impacts of '3F4D+MD' on soil health. While short-term benefits in terms of arsenic immobilization and plant

physiology are clear, the cumulative effects on soil organic matter, microbial communities, and overall fertility in the unique Bengal delta environment warrant deeper investigation [23]. Finally, research should explore the synergistic integration of '3F4D+MD' with other recommended agronomic practices. Investigating how it interacts with strategies like fertilizer deep placement (FDP), the use of arsenic-accumulation-resistant rice genotypes, and diversified cropping systems like rice-vegetable rotations could unlock even greater efficiencies and benefits, creating a more robust and resilient model for sustainable agriculture in Bangladesh. Addressing these research priorities will be crucial for moving '3F4D+MD' from a promising pilot project to a cornerstone of Bangladesh's future agricultural strategy.

---

en

1. (PDF) Arsenic management in contaminated irrigation water for rice ... https://www.researchgate.net/publication/301726420_Arsenic_management_in_contaminated_irrigation_water_for_rice_cultivation
2. (PDF) Arsenic Contamination in Irrigation Water for Rice Production ... https://www.researchgate.net/publication/256949489_Arsenic_Contamination_in_Irrigation_Water_for_Rice_Production_in_Bangladesh_A_Review
3. Influence of soil arsenic in rice and its mitigation through water ... https://www.researchgate.net/publication/343983673_Influence_of_soil_arsenic_in_rice_and_its_mitigation_through_water_management
4. (PDF) Arsenic Contamination of Paddy Fields through Groundwater ... https://www.researchgate.net/publication/270891628_Arsenic_Contamination_of_Paddy_Fields_through_Groundwater_Irrigation_in_Bangladesh_Risks_for_Rice_Production_and_Mitigation_Perspectives
5. (PDF) Arsenic Contamination in Rice Production Irrigation Water in ... https://www.researchgate.net/publication/358477138_Arsenic_Contamination_in_Rice_Production_Irrigation_Water_in_Bangladesh
6. (PDF) Investigating alternate wetting and drying irrigation method ... https://www.researchgate.net/publication/396945478_Investigating_alternate_wetting_and_drying_irrigation_method_over_con

ventional_practice_for_Boro_rice_production_in_Bangladesh_a_sustainable_water_management_practice_in_agriculture

7. Climate-based suitability assessment for alternate wetting and ... https://www.tandfonline.com/doi/full/10.1080/17583004.2017.1362945
8. Farmers' Participatory Alternate Wetting and Drying Irrigation ... - MDPI https://www.mdpi.com/2073-4441/14/7/1056
9. The potential for expansion of irrigated rice under alternate wetting ... https://www.sciencedirect.com/science/article/abs/pii/S0378377421000238
10. Mitigation Potential of Greenhouse Gas Emissions Through Nitrogen ... https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2025EF006740
11. Field study of rice yield diminished by soil arsenic in Bangladesh https://pmc.ncbi.nlm.nih.gov/articles/PMC5645253/
12. Increase in Rice Grain Arsenic for Regions of Bangladesh Irrigating ... https://www.academia.edu/108662118/Increase_in_Rice_Grain_Arsenic_for_Regions_of_Bangladesh_Irrigating_Paddies_with_Elevated_Arsenic_in_Groundwaters
13. Mitigation of arsenic in rice through deficit irrigation in field and use ... https://www.researchgate.net/publication/271681905_Mitigation_of_arsenic_in_rice_through_deficit_irrigation_in_field_and_use_of_filtered_water_in_kitchen
14. Effect of irrigation and genotypes towards reduction in arsenic load ... https://www.sciencedirect.com/science/article/abs/pii/S0048969717318144
15. The Reaction of Rice Growth to the Arsenic Contamination under ... https://www.mdpi.com/2223-7747/13/9/1253
16. Arsenic Shoot-Grain Relationships in Field Grown Rice Cultivars https://pubs.acs.org/doi/abs/10.1021/es902992d
17. Agronomic solutions to decrease arsenic concentrations in rice - PMC https://pmc.ncbi.nlm.nih.gov/articles/PMC12084250/
18. [PDF] Arsenic in Rice Agro-Ecosystem: Solutions for Safe and Sustainable ... https://www.frontiersin.org/journals/sustainable-food-systems/articles/10.3389/fsufs.2020.00053/pdf
19. Sprinkler irrigation of rice fields reduces grain arsenic but enhances ... https://www.sciencedirect.com/science/article/abs/pii/S004896971400446X
20. [PDF] Evaluating the impacts of water management on arsenic uptake in ... https://escholarship.org/content/qt77h346wt/qt77h346wt_noSplash_f0cfc47e3eb3920a66306c32563c6921.pdf
21. Alternate Wetting and Drying Limits Arsenic in Porewater and Rice ... https://pubs.acs.org/doi/10.1021/acs.est.5c03552

22. Impact of Alternate Wetting and Drying Irrigation on Arsenic Uptake ... https://www.sciencedirect.com/science/article/abs/pii/S0167880918304638
23. Alternate wetting and drying decreases arsenic content and ... https://www.researchgate.net/publication/379375378_Alternate_wetting_and_drying_decreases_arsenic_content_and_increases_yield_of_rice_grown_in_organic_matter_amended_soil
24. Alternate Wetting and Drying Decreases Methylmercury in Flooded ... https://acsess.onlinelibrary.wiley.com/doi/full/10.2136/sssaj2017.05.0158
25. Effects of Alternate Wetting and Drying (AWD) Irrigation on Rice ... https://www.mdpi.com/2073-4395/15/10/2372
26. Irrigation management for arsenic mitigation in rice grain https://www.sciencedirect.com/science/article/abs/pii/S0048969718331802
27. [PDF] Irrigation Practices on Rice Crop Production in Arsenic-Rich Paddy ... https://pdfs.semanticscholar.org/8c7a/445bcd3c08d3b7e22deee2be100949d757a1.pdf
28. Impact of alternate wetting and drying on rice physiology, grain ... https://www.sciencedirect.com/science/article/abs/pii/S0378429017301533
29. Effect of Alternate Wetting and Drying (AWD) Irrigation for Boro Rice ... https://www.researchgate.net/publication/275567173_Effect_of_Alternate_Wetting_and_Drying_AWD_Irrigation_for_Boro_Rice_Cultivation_in_Bangladesh
30. Alternate wetting and drying in Bangladesh: Water − saving farming ... https://onlinelibrary.wiley.com/doi/full/10.1002/fes3.149
31. Alternate wetting and drying irrigation maintained rice yields despite ... https://pmc.ncbi.nlm.nih.gov/articles/PMC4998133/
32. Comparison of Agronomic and Physiological Characteristics for Rice ... https://www.mdpi.com/2073-4395/14/9/1986
33. Arsenic contamination, impact and mitigation strategies in rice agro ... https://www.sciencedirect.com/science/article/abs/pii/S0048969721045514
34. Arsenic (As) concentrations in groundwater of Bangladesh, showing ... https://www.researchgate.net/figure/Arsenic-As-concentrations-in-groundwater-of-Bangladesh-showing-all-64-districts-under_fig1_235366676
35. [PDF] Interrelationship Among Rice Grain Arsenic, Micronutrients Content ... https://www.frontiersin.org/journals/environmental-science/articles/10.3389/fenvs.2022.857629/pdf
36. Water usage and productivity of Boro rice at the field level https://ideas.repec.org/a/eee/agiwat/v240y2020ics0378377420301153.html
37. Water usage and productivity of Boro rice at the field level and their ... https://www.researchgate.net/publication/

341978957_Water_usage_and_productivity_of_Boro_rice_at_the_field_level_and_their_impacts_on_the_sustainable_groundwater_irrigation_in_the_North-West_Bangladesh

38. Water usage and productivity of Boro rice at the field level and their ... https://www.semanticscholar.org/paper/8757a6984a5ff0d4d6744a71b673238592b1fbe8
39. (PDF) Investigating water productivity and yield of boro rice under ... https://www.researchgate.net/publication/333015808_Investigating_water_productivity_and_yield_of_boro_rice_under_conventional_and_conservation_irrigation_practices_in_Bangladesh
40. (PDF) Water productivity of boro rice - ResearchGate https://www.researchgate.net/publication/273257494_Water_productivity_of_boro_rice
41. Yield and water productivity variation of Boro rice with irrigation ... https://www.researchgate.net/publication/360848518_Yield_and_water_productivity_variation_of_Boro_rice_with_irrigation_strategies_and_transplanting_dates_under_climate_change_-_a_case_study_in_south-western_Bangladesh
42. (PDF) Optimizing water footprints and yield sustainability in Boro rice ... https://www.researchgate.net/publication/391553409_Optimizing_water_footprints_and_yield_sustainability_in_Boro_rice_through_alternate_wetting_and_intense_drying_a_climate-resilient_strategy_for_Bangladesh
43. Arsenic in the water and agricultural crop production system - Springer https://link.springer.com/article/10.1007/s11356-022-20880-0
44. Soil arsenic but not rice arsenic increasing with arsenic in irrigation ... https://pmc.ncbi.nlm.nih.gov/articles/PMC8188821/
45. Translocation of Soil Arsenic towards Accumulation in Rice - MDPI https://www.mdpi.com/2073-4441/13/20/2816
46. Rice yields and water use under alternate wetting and drying irrigation https://www.sciencedirect.com/science/article/pii/S0378429016307791
47. (PDF) Bangladesh CSA-Technologies-and-Practices in Bangladesh https://www.researchgate.net/publication/377074837_Bangladesh_CSA-Technologies-and-Practices_in_Bangladesh
48. Effects of integrated plant nutrition systems with fertilizer deep ... - PMC https://pmc.ncbi.nlm.nih.gov/articles/PMC10703841/
49. Dry direct-seeded and broadcast rice: A profitable and climate-smart ... https://www.sciencedirect.com/science/article/pii/S0378429025000048
50. [PDF] Climate Change Trends and Vulnerabilities in Bangladesh's Crop ... https://www.cabidigitallibrary.org/doi/pdf/10.5555/20250106939

51. [PDF] MASTER PLAN FOR AGRICULTURAL DEVELOPMENT IN THE ... https://faolex.fao.org/docs/pdf/BGD214472.pdf
52. [PDF] Master Plan for Agricultural Development in the Southern Region of ... https://openknowledge.fao.org/server/api/core/bitstreams/8f92d4ef-633e-4796-af89-0b1d69b8e2f0/content
53. Effect of Arsenic Contamination on Yield Loss and ... - ResearchGate https://www.researchgate.net/publication/328757105_Effect_of_Arsenic_Contamination_on_Yield_Loss_and_Arsenic_Concentration_of_Rice
54. [PDF] Growth, Yield and Grain Arsenic Concentration of Rice Cultivars ... https://sdiopr.s3.ap-south-1.amazonaws.com/2022/Sep/2022_AJSSPN_91631/Rev_AJSSPN_91631_Mos_A.pdf
55. Effects of Water Management, Arsenic and Phosphorus Levels on ... https://www.researchgate.net/publication/261374090_Effects_of_Water_Management_Arsenic_and_Phosphorus_Levels_on_Rice_Yield_in_High-Arsenic_Soil-Water_System?_tp=eyJjb250ZXh0Ijp7InBhZ2UiOiJzY2llbnRpZmljQ29udHJpYnV0aW9ucyIsInByZXZpb3VzUGFnZSI6bnVsbCwic3ViUGFnZSI6bnVsbH19
56. Groundwater resources of the world and their use https://unesdoc.unesco.org/ark:/48223/pf0000134433
57. An arsenic-contaminated field trial to assess the uptake ... - PubMed https://pubmed.ncbi.nlm.nih.gov/23149722/
58. (PDF) Experiment for Arsenic Accumulation into Rice Cultivated with ... https://www.researchgate.net/publication/253649681_Experiment_for_Arsenic_Accumulation_into_Rice_Cultivated_with_Arsenic_Enriched_Irrigation_Water_in_Bangladesh
59. Meta-Analysis Enables Prediction of the Maximum Permissible ... https://www.frontiersin.org/journals/environmental-science/articles/10.3389/fenvs.2021.760125/full
60. I. Estimating normal levels of total arsenic in rice grain - PubMed https://pubmed.ncbi.nlm.nih.gov/18546734/
61. Arsenic toxicity to rice (Oryza sativa L.) in Bangladesh - ResearchGate https://www.researchgate.net/publication/227109540_Arsenic_toxicity_to_rice_Oryza_sativa_L_in_Bangladesh
62. (PDF) Arsenic status of water, soil, rice grain and straw of individual ... https://www.researchgate.net/publication/237678121_Arsenic_status_of_water_soil_rice_grain_and_straw_of_individual_shallow_tube_well_command_area_of_Brahmanbaria

63. (PDF) Evaluation of groundwater quality and suitability for irrigation ... https://www.researchgate.net/publication/306923949_Evaluation_of_groundwater_quality_and_suitability_for_irrigation_and_drinking_purposes_in_southwest_Punjab_India_using_hydrochemical_approach
64. [PDF] Contents - Food and Agriculture Organization of the United Nations https://www.fao.org/fileadmin/templates/agphome/scpi/cfscc2010/2-Full_papers.pdf
65. P173312-c061082c-bd30-4bef-a67c-7ce499411e0e.txt https://documents1.worldbank.org/curated/en/099121525020593868/txt/P173312-c061082c-bd30-4bef-a67c-7ce499411e0e.txt
66. P173312-6d47a6c2-9a25-4b2a-b562-1c622abb43be.txt https://documents1.worldbank.org/curated/en/099121525013574972/txt/P173312-6d47a6c2-9a25-4b2a-b562-1c622abb43be.txt
67. P173312-2735ecf8-829c-4dcb-bf99-6ebab4ea6ff5.txt https://documents1.worldbank.org/curated/en/099121525014520385/txt/P173312-2735ecf8-829c-4dcb-bf99-6ebab4ea6ff5.txt
68. 906000v20PUB0E00Box385314... - World Bank Documents & Reports https://documents1.worldbank.org/curated/en/349541468302961396/txt/906000v20PUB0E00Box385314B00PUBLIC0.txt
69. [PDF] COUNTRY REPORT – Bangladesh - FAO Knowledge Repository https://openknowledge.fao.org/server/api/core/bitstreams/15f30eda-2b43-4dcc-b84f-244379758a04/content
70. [PDF] Annual_Report_2019_f.pdf - Jamuna Bank https://jamunabankbd.com/wp-content/uploads/assets/file/Annual_Report_2019_f.pdf
71. [PDF] Sight and Life - DOI https://doi.org/10.52439/MOPG2475
72. [PDF] Soil-crop suitability classification for Bangladesh https://www.fao.org/4/ar834e/ar834e.pdf
73. [PDF] Variations in Water and Deposited Sediment Qualities in the Tidal ... https://pdfs.semanticscholar.org/7ad4/cb3b41abd63cc14f4eb85997d82cfece864f.pdf
74. Effect of water management, tillage options and phosphorus status ... https://agris.fao.org/search/en/providers/122535/records/65dfeafc0f3e94b9e5dd9f5f
75. Sustainable water management in rice cultivation reduces arsenic ... https://pubmed.ncbi.nlm.nih.gov/38309156/
76. Government of The Peoples Republic of Ba | PDF | Bangladesh https://www.scribd.com/document/721159362/Government-of-the-Peoples-Republic-of-Ba
77. (PDF) Witness to Marvels - Academia.edu https://www.academia.edu/44185006/Witness_to_Marvels
78. MORE EBOOKS AT @aj - Ebooks | PDF | Agronomy | Agriculture https://www.scribd.com/document/490127260/ARD

79. (PDF) Ganga: A Journey Down the Ganges River - Academia.edu https://www.academia.edu/79161967/Ganga_A_Journey_Down_the_Ganges_River
80. [PDF] environmental assessment report - World Bank Document https://documents1.worldbank.org/curated/en/111141468006270304/pdf/E938.pdf
81. Arsenic contamination of groundwater: A global synopsis with focus ... https://www.sciencedirect.com/science/article/pii/S1674987120302115
82. A Review of Groundwater Arsenic in the Bengal Basin, Bangladesh ... https://link.springer.com/article/10.1007/s40726-015-0022-0
83. (PDF) Arsenic in groundwater of the Bengal Basin, Bangladesh https://www.researchgate.net/publication/225822311_Arsenic_in_groundwater_of_the_Bengal_Basin_Bangladesh_Distribution_field_relations_and_hydrogeological_setting
84. P181147-a0425c34-9b81-4d94-8101-4e53382e4bef.txt https://documents1.worldbank.org/curated/en/099113024053511836/txt/P181147-a0425c34-9b81-4d94-8101-4e53382e4bef.txt
85. [PDF] Bangladesh Municipal Water Supply and Sanitation Project ... https://documents1.worldbank.org/curated/en/393701536909036591/pdf/Revised-Final-EMF-DPHE-BMWSSP-30-07-18-R.pdf
86. Development and evaluation of the first automated irrigation system ... https://www.sciencedirect.com/science/article/pii/S2772375523001776
87. Arsenic in the water and agricultural crop production system - PMC https://pmc.ncbi.nlm.nih.gov/articles/PMC9288370/
88. Arsenic contamination in groundwater in Bangladesh - PMC - NIH https://pmc.ncbi.nlm.nih.gov/articles/PMC6281155/
89. [PDF] World Bank Document https://documents1.worldbank.org/curated/en/422221468013803608/pdf/multi0page.pdf
90. [PDF] Arsenic contamination of groundwater in Bangladesh https://www.iaea.org/sites/default/files/arseniccontamination.pdf
91. [PDF] World Bank Document https://documents1.worldbank.org/curated/en/271331468013248845/pdf/multi0page.pdf
92. Effect of water management system on the performance of boro rice https://www.researchgate.net/publication/332343409_Effect_of_water_management_system_on_the_performance_of_boro_rice
93. (PDF) Investigating water productivity and yield of boro rice under ... https://www.academia.edu/70656482/Investigating_water_productivity_and_yield_of_boro_rice_under_conventional_and_conservation_irrigation_practices_in_Bangladesh

94. Recycling deep percolated water in continuously flooding irrigated ... https://www.researchgate.net/publication/361042491_Recycling_deep_percolated_water_in_continuously_flooding_irrigated_rice_fields_to_mitigate_water_scarcity
95. Impacts of diverse water management systems on growth and yield ... https://www.researchgate.net/publication/397173506_Impacts_of_diverse_water_management_systems_on_growth_and_yield_of_two_prominent_boro_rice_cultivars_in_Bangladesh
96. A Review of Groundwater Arsenic Contamination in Bangladesh https://pmc.ncbi.nlm.nih.gov/articles/PMC4772235/
97. (PDF) Arsenic Contamination of Groundwater in Bangladesh https://www.researchgate.net/publication/338159377_Arsenic_Contamination_of_Groundwater_in_Bangladesh
98. Arsenic in tube well water in Bangladesh: health and economic ... https://pmc.ncbi.nlm.nih.gov/articles/PMC3506399/
99. Status of groundwater arsenic contamination in Bangladesh https://www.researchgate.net/publication/45504376_Status_of_groundwater_arsenic_contamination_in_Bangladesh_A_14-year_study_report
100. Arsenic contamination of groundwater and its effects on drinking ... https://www.researchgate.net/publication/399196822_Arsenic_contamination_of_groundwater_and_its_effects_on_drinking_water_irrigation_and_public_health_in_Bangladesh
101. Assessment of Arsenic Concentration in Ground Water of Tube ... https://www.researchgate.net/publication/269493615_Assessment_of_Arsenic_Concentration_in_Ground_Water_of_Tube_Wells_in_Selected_Primary_Schools'_at_Palashbari_Upazila_of_Gaibanda_District
102. 6 - CodaLab https://worksheets.codalab.org/rest/bundles/0xadf98bb30a99476ab56ebff3e462d4fa/contents/blob/glove.6B.100d.txt-vocab.txt
103. Low-cost informational intervention reduced drinking water arsenic ... https://pmc.ncbi.nlm.nih.gov/articles/PMC10042277/
104. Low-cost informational intervention reduced drinking water arsenic ... https://pubmed.ncbi.nlm.nih.gov/36992819/
105. Structural Evidence of a Productive Active Site Architecture for an ... https://pmc.ncbi.nlm.nih.gov/articles/PMC3629905/
106. Structural basis for thermostability revealed through the identification ... https://pmc.ncbi.nlm.nih.gov/articles/PMC5433624/

107. Directed Evolution of a Thermostable Quorum-quenching Lactonase ... https://www.sciencedirect.com/science/article/pii/S0021925819762590

108. Active Site Loop Conformation Regulates Promiscuous Activity in a ... https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0115130

109. Arsenic in tube well water in Bangladesh - PubMed https://pubmed.ncbi.nlm.nih.gov/23226896/

110. (PDF) Arsenic in tube well water in Bangladesh - ResearchGate https://www.researchgate.net/publication/233889514_Arsenic_in_tube_well_water_in_Bangladesh_Health_and_economic_impacts_and_implications_for_arsenic_mitigation

111. A Review of Groundwater Arsenic Contamination in Bangladesh https://www.mdpi.com/1660-4601/13/2/215

112. Status of groundwater arsenic contamination in Bangladesh - PubMed https://pubmed.ncbi.nlm.nih.gov/20684969/

113. [PDF] Evidence for Elevated Levels of Arsenic in Public Wells of ... https://www.ldeo.columbia.edu/~avangeen/publications/documents/Choudhury_GW_16.pdf

114. Changing Climate and Surface and Ground Water Related Issues in ... https://www.academia.edu/21803713/Changing_Climate_and_Surface_and_Ground_Water_Related_Issues_in_Dhaka

115. September 2022 | PDF | Startup Company - Scribd https://www.scribd.com/document/604905401/September-2022

116. [PDF] the survey of india http://archivenepal.s3.amazonaws.com/digitalhimalaya/collections/rarebooks/downloads/HRSA_05.pdf

117. [PDF] FARMING SYSTEMS OF BANGLADESH https://creativeconsulting-bd.com/wp-content/uploads/2025/09/8-Yusuf-Farming-Systems-of-Bangladesh-full-1.pdf

118. [PDF] Bangladesh: Poverty Reduction Strategy Paper https://www.imf.org/external/pubs/ft/scr/2005/cr05410.pdf

119. [PDF] Initial Environmental Examination - Asian Development Bank https://www.adb.org/sites/default/files/linked-documents/53382-001-ieeab.pdf

120. Haor Master Plan Volume 1 | PDF | Economic Development - Scribd https://www.scribd.com/document/432793418/Haor-Master-Plan-Volume-1