

First Name: Zhihong

Last Name: Wang

Student ID: 1002095207

First Name: Aofei

Last Name: Liu

Student ID: 1002291334

First Name: Heng

Last Name: Ye

Student ID: 1001846363

We declare that this assignment is solely our own work, and is in accordance with the University of Toronto Code of Behaviour on Academic Matters.

This submission has been prepared using L^AT_EX.

Problem 1.

(6 MARKS)

Define the sequence $\{a_k\}$ as follows: $a_0 = 0, a_1 = 0, a_2 = 2, a_k = 3a_{\lfloor \frac{k}{2} \rfloor} + 2$ for all $k \geq 3$.

1. (2 MARKS) Find the first 8 terms of the sequence.

2. (4 MARKS) Prove that

$$\forall n \in \mathbb{N} : E(a_n)$$

where $E(n)$ is the usual predicate "n is even".

Solution

1. $a_0 = 0$

$$a_1 = 0$$

$$a_2 = 2$$

$$\begin{aligned} a_3 &= 3a_{\lfloor \frac{3}{2} \rfloor} + 2 \\ &= 3a_{\lfloor 1.5 \rfloor} + 2 \\ &= 3a_1 + 2 \\ &= 3 \times 0 + 2 \\ &= 2 \end{aligned}$$

$$\begin{aligned} a_4 &= 3a_{\lfloor \frac{4}{2} \rfloor} + 2 \\ &= 3a_{\lfloor 2 \rfloor} + 2 \\ &= 3a_2 + 2 \\ &= 3 \times 2 + 2 \\ &= 8 \end{aligned}$$

$$\begin{aligned} a_5 &= 3a_{\lfloor \frac{5}{2} \rfloor} + 2 \\ &= 3a_{\lfloor 2.5 \rfloor} + 2 \\ &= 3a_2 + 2 \\ &= 3 \times 2 + 2 \\ &= 8 \end{aligned}$$

$$\begin{aligned} a_6 &= 3a_{\lfloor \frac{6}{2} \rfloor} + 2 \\ &= 3a_{\lfloor 3 \rfloor} + 2 \\ &= 3a_3 + 2 \\ &= 3 \times 2 + 2 \\ &= 8 \end{aligned}$$

$$\begin{aligned} a_7 &= 3a_{\lfloor \frac{7}{2} \rfloor} + 2 \\ &= 3a_{\lfloor 3.5 \rfloor} + 2 \end{aligned}$$

$$\begin{aligned}
&= 3a_3 + 2 \\
&= 3 \times 2 + 2 \\
&= 8
\end{aligned}$$

2. Let $n \in \mathbb{N}$

Let $P(n)$ represent $E(a_n)$

‡ Prove by strong induction

Basis step: prove $P(0)$, $P(1)$, $P(2)$, $P(3)$

$$a_0 = 0$$

Then $P(0)$

$$a_1 = 0$$

Then $P(1)$

$$a_2 = 2$$

Then $P(2)$

$$a_3 = 3a_{\lfloor \frac{3}{2} \rfloor} + 2 = 3a_1 + 2 = 3 \times 0 + 2 = 2$$

Then $P(3)$

Inductive step: prove $\forall k \in \mathbb{N} : [k \geq 3 \rightarrow ((\forall i \in \{0, \dots, k\} : P(i)) \rightarrow P(k+1))]$

Let $k \in \mathbb{N}$

Assume $k \geq 3$

Assume $\forall i \in \{0, \dots, k\} : P(i)$

$$\text{Then } a_{k+1} = 3a_{\lfloor \frac{k+1}{2} \rfloor} + 2$$

$$\text{Then } 0 < \lfloor \frac{k+1}{2} \rfloor \leq \frac{k+1}{2} \text{ and } \frac{k+1}{2} < k+1 \text{ ‡ } k \geq 3$$

$$\text{Then } \lfloor \frac{k+1}{2} \rfloor \leq \frac{k+1}{2} < k+1$$

$$\text{Then } \lfloor \frac{k+1}{2} \rfloor < k+1$$

$$\text{Then } \lfloor \frac{k+1}{2} \rfloor \leq k \text{ ‡ } \lfloor \frac{k+1}{2} \rfloor \text{ and } k \text{ are integers}$$

$$\text{Then } \lfloor \frac{k+1}{2} \rfloor \in \{0, \dots, k\}$$

$$\text{Then } a_{\lfloor \frac{k+1}{2} \rfloor} \text{ is even ‡ By assumption}$$

$$\text{Then } \exists j \in \mathbb{Z} : a_{\lfloor \frac{k+1}{2} \rfloor} = 2j$$

$$\text{Let } j_0 \in \mathbb{Z} \text{ such that } a_{\lfloor \frac{k+1}{2} \rfloor} = 2j_0$$

$$\text{Then } 3 \times a_{\lfloor \frac{k+1}{2} \rfloor} = 6j_0$$

$$\begin{aligned}
\text{Then } 3 \times a_{\lfloor \frac{k+1}{2} \rfloor} + 2 &= 6j_0 + 2 \\
&= 2(3j_0 + 1)
\end{aligned}$$

$$\text{Let } j_1 = 3j_0 + 1$$

$$\text{Then } j_1 \in \mathbb{Z}$$

$$\text{Then } 3 \times a_{\lfloor \frac{k+1}{2} \rfloor} + 2 = 2j_1$$

Then $\exists j \in \mathbb{Z} : 3 \times a_{\lfloor \frac{k+1}{2} \rfloor} + 2 = 2j$

Then $3a_{\lfloor \frac{k+1}{2} \rfloor} + 2$ is even

Then a_{k+1} is even

Then $P(k+1)$

Therefore $(\forall i \in \{0, \dots, k\} : P(i)) \rightarrow P(k+1)$

Therefore $k \geq 3 \rightarrow ((\forall i \in \{0, \dots, k\} : P(i)) \rightarrow P(k+1))$

Therefore $\forall k \in \mathbb{N} : [k \geq 3 \rightarrow ((\forall i \in \{0, \dots, k\} : P(i)) \rightarrow P(k+1))]$

Therefore $\forall n \in \mathbb{N} : E(a_n)$ \sharp By strong induction

Problem 2.

(6 MARKS)

Prove or disprove:

1. (3 MARKS) $\forall x \in \mathbb{R} : \forall n \in \mathbb{N} : \lfloor x + n \rfloor = \lfloor x \rfloor + n$
2. (3 MARKS) $\forall x \in \mathbb{R} : \forall n \in \mathbb{N} : \lfloor nx \rfloor = n \lfloor x \rfloor$

Solution

1. True statement. Prove.

Let $x \in \mathbb{R}$

Let $n \in \mathbb{N}$

‡ Definition of floor: $\forall x \in \mathbb{R} : (\lfloor x \rfloor \in \mathbb{Z}) \wedge (\lfloor x \rfloor \leq x) \wedge (\forall z \in \mathbb{Z} : (z \leq x) \rightarrow (z \leq \lfloor x \rfloor))$

‡ Prove $\lfloor x + n \rfloor \leq n + \lfloor x \rfloor$

Then $\lfloor x + n \rfloor \leq x + n$ ‡ By definition of $\lfloor x + n \rfloor$

Then $\lfloor x + n \rfloor - n \leq x$ and $\lfloor x + n \rfloor - n \in \mathbb{Z}$ ‡ $\lfloor x + n \rfloor \in \mathbb{Z}$ and $n \in \mathbb{N}$

Then $\lfloor x + n \rfloor - n \leq \lfloor x \rfloor$ ‡ By definition of $\lfloor x \rfloor$

Then $\lfloor x + n \rfloor \leq n + \lfloor x \rfloor$

‡ Prove $\lfloor x \rfloor + n \leq \lfloor x + n \rfloor$

Then $\lfloor x \rfloor \leq x$ ‡ By definition of $\lfloor x \rfloor$

Then $\lfloor x \rfloor + n \leq x + n$ and $\lfloor x \rfloor + n \in \mathbb{Z}$ ‡ $\lfloor x \rfloor \in \mathbb{Z}$ and $n \in \mathbb{N}$

Then $\lfloor x \rfloor + n \leq \lfloor x + n \rfloor$ ‡ By definition of $\lfloor x + n \rfloor$

Then $(\lfloor x + n \rfloor \leq n + \lfloor x \rfloor) \wedge (\lfloor x + n \rfloor \geq \lfloor x \rfloor + n)$

Then $\lfloor x + n \rfloor = \lfloor x \rfloor + n$

Therefore $\forall n \in \mathbb{N} : \lfloor x + n \rfloor = \lfloor x \rfloor + n$

Therefore $\forall x \in \mathbb{R} : \forall n \in \mathbb{N} : \lfloor x + n \rfloor = \lfloor x \rfloor + n$

2. False statement. Disprove by proving negation is true.

Prove $\neg(\forall x \in \mathbb{R} : \forall n \in \mathbb{N} : \lfloor nx \rfloor = n \lfloor x \rfloor)$

Prove $\exists x \in \mathbb{R} : [\exists n \in \mathbb{N} : [\lfloor nx \rfloor \neq n \lfloor x \rfloor]]$

Let $x = 1.5, n = 2$

Then $x \in \mathbb{R}, n \in \mathbb{N}$

Then $nx = 3$

Then $\lfloor nx \rfloor = 3$ # By definition of $\lfloor nx \rfloor$

Then $\lfloor x \rfloor = 1$ # By definition of $\lfloor x \rfloor$

Then $n\lfloor x \rfloor = 2$

Then $\lfloor nx \rfloor \neq n\lfloor x \rfloor$ # $3 \neq 2$

Therefore $\exists n \in \mathbb{N} : [\lfloor nx \rfloor \neq n\lfloor x \rfloor]$

Therefore $\exists x \in \mathbb{R} : [\exists n \in \mathbb{N} : [\lfloor nx \rfloor \neq n\lfloor x \rfloor]]$

Therefore the negation is true.

Therefore the original statement " $(\forall x \in \mathbb{R} : \forall n \in \mathbb{N} : \lfloor nx \rfloor = n\lfloor x \rfloor)$ " is false.

Problem 3.

(5 MARKS)

Prove the following claims:

1. (3 MARKS) $\forall x \in [0, \pi/2] : \sin x + \cos x \geq 1$.
2. (2 MARKS) Prove that $\log_2 3$ is irrational.

Solution

1. Prove by contradiction

Assume $\neg(\forall x \in [0, \frac{\pi}{2}] : \sin x + \cos x \geq 1)$

Then $\exists x \in [0, \frac{\pi}{2}] : \sin x + \cos x < 1$

Let $x_0 \in [0, \frac{\pi}{2}]$ such that $\sin x_0 + \cos x_0 < 1$

Then $(1 \geq \sin x_0 \geq 0) \wedge (1 \geq \cos x_0 \geq 0) \nmid x \in [0, \frac{\pi}{2}]$

Then $\sin x_0 \cos x_0 \geq 0$

Then $\sin x_0 + \cos x_0 \geq 0$

Then $0 \leq \sin x_0 + \cos x_0 < 1 \nmid$ By assumption

Then $(\sin x_0 + \cos x_0)^2 < 1$

Then $\sin^2 x_0 + \cos^2 x_0 + 2\sin x_0 \cos x_0 < 1$

Then $1 + 2\sin x_0 \cos x_0 < 1 \nmid \sin^2 x_0 + \cos^2 x_0 = 1$

Then $2\sin x_0 \cos x_0 < 0$

Then $\sin x_0 \cos x_0 < 0 \nmid$ Contradiction with $\sin x_0 \cos x_0 \geq 0$

Then $\sin x + \cos x \geq 1$

Therefore $\forall x \in [0, \frac{\pi}{2}] : \sin x + \cos x \geq 1$

2. Prove by contradiction

Assume $\log_2 3$ is rational

Then $\exists p, q \in \mathbb{Z} : \log_2 3 = \frac{p}{q} \wedge q \neq 0$

Let $p_0, q_0 \in \mathbb{Z}$ such that $\log_2 3 = \frac{p_0}{q_0} \wedge q_0 \neq 0$

Then $\log_2 3 > 0 \nmid$ By definition of logarithm

Then $\frac{p_0}{q_0} > 0$

Then $(p_0 > 0 \wedge q_0 > 0) \vee (p_0 < 0 \wedge q_0 < 0)$

Then $p_0 > 0 \wedge q_0 > 0$

\nmid If p_0 and q_0 are both less than 0, we can erase their negative sign at once

\nmid then they become the same case of p_0 and q_0 are both greater than 0

Then p_0, q_0 are integers which greater or equal to 1 \nmid By assumption

Then $\forall p_0, q_0 \in \mathbb{Z} : (p_0 \geq 1) \wedge (q_0 \geq 1)$

Then $\forall p_0, q_0 \in \mathbb{N} : (p_0 \geq 1) \wedge (q_0 \geq 1)$ # Natural numbers are integers begin with 0

Then $2^{\frac{p_0}{q_0}} = 3$ # By definition of logarithm

Then $2^{p_0} = 3^{q_0}$

Prove 2^{p_0} is even by induction

Basis step: Prove 2^1

$2^1 = 2$

Then 2^1 is even

Inductive step: Prove $\forall k \in \mathbb{N} : [(k \geq 1) \rightarrow (2^k \text{ is even} \rightarrow 2^{k+1} \text{ is even})]$

Let $k \in \mathbb{N}$

Assume $k \geq 1$

Assume 2^k is even

Then $\exists j \in \mathbb{Z} : 2^k = 2j$

Let $j_0 \in \mathbb{Z}$ such that $2^k = 2j_0$

Then $2^{k+1} = 2^k \times 2$

Then $2^{k+1} = 2j_0 \times 2$

Let $j_1 = 2j_0$

Then $j_1 \in \mathbb{Z}$

Then $2^{k+1} = 2j_1$

Then $\exists j \in \mathbb{Z} : 2^{k+1} = 2j$

Then 2^{k+1} is even

Therefore $2^k \text{ is even} \rightarrow 2^{k+1} \text{ is even}$

Therefore $k \geq 1 \rightarrow (2^k \text{ is even} \rightarrow 2^{k+1} \text{ is even})$

Therefore $\forall k \in \mathbb{N} : [(k \geq 1) \rightarrow (2^k \text{ is even} \rightarrow 2^{k+1} \text{ is even})]$

Therefore 2^{p_0} is even # By induction

Prove 3^{q_0} is odd by induction

Basis step: Prove 3^1

$3^1 = 3$

Then 3^1 is odd

Inductive step: Prove $\forall k \in \mathbb{N} : [(k \geq 1) \rightarrow (3^k \text{ is odd} \rightarrow 3^{k+1} \text{ is odd})]$

Let $k \in \mathbb{N}$

Assume $k \geq 1$

Assume 3^k is odd

Then $\exists j \in \mathbb{Z} : 3^k = 2j + 1$

Let $j_0 \in \mathbb{Z}$ such that $3^k = 2j_0 + 1$

Then $3^{k+1} = 3^k \times 3$

Then $3^{k+1} = (2j_0 + 1) \times 3$

Then $3^{k+1} = 6j_0 + 3$

Then $3^{k+1} = 6j_0 + 2 + 1$

Then $3^{k+1} = 2(3j_0 + 1) + 1$

Let $j_1 = 3j_0 + 1$

Then $j_1 \in \mathbb{Z}$

Then $3^{k+1} = 2j_1 + 1$

Then $\exists j \in \mathbb{Z} : 3^{k+1} = 2j + 1$

Then 3^{k+1} is odd

Therefore 3^k is odd $\rightarrow 3^{k+1}$ is odd

Therefore $k \geq 1 \rightarrow (3^k \text{ is odd} \rightarrow 3^{k+1} \text{ is odd})$

Therefore $\forall k \in \mathbb{N} : [(k \geq 1) \rightarrow (3^k \text{ is odd} \rightarrow 3^{k+1} \text{ is odd})]$

Therefore 3^{q_0} is odd \nmid By induction

Then even number = odd number \nmid Contradiction with even \neq odd

Then $\forall p, q \in \mathbb{Z} : \log_2 3 \neq \frac{p}{q}$

Therefore $\log_2 3$ is irrational

Problem 4.

(6 MARKS) Prove the following:

1. (3 MARKS) $\forall x, y \in \mathbb{R} : x^2 + y^2 = (x + y)^2 \Leftrightarrow x = 0 \vee y = 0$.
2. (3 MARKS) $\forall x, y \in \mathbb{R} : x^3 + x^2y = y^2 + xy \Leftrightarrow y = x^2 \vee y = -x$.

Solution

1. Let $x, y \in \mathbb{R}$

First, prove $x^2 + y^2 = (x + y)^2 \rightarrow x = 0 \vee y = 0$

Assume $x^2 + y^2 = (x + y)^2$

Then $(x + y)^2 = x^2 + 2xy + y^2$

Then $x^2 + y^2 = x^2 + 2xy + y^2$

Then $2xy = 0$

Then $xy = 0$

Case 1

Assume $x = 0$

Then $x = 0$

Case 2

Assume $x \neq 0$

Then $\frac{xy}{x} = \frac{0}{x}$

Then $y = 0$

Then $(x = 0) \vee (y = 0)$

Therefore $x^2 + y^2 = (x + y)^2 \rightarrow x = 0 \vee y = 0$

Second, prove $x = 0 \vee y = 0 \rightarrow x^2 + y^2 = (x + y)^2$

Assume $x = 0 \vee y = 0$

Case 1

Assume $x = 0$

Then $xy = 0$

Then $2xy = 0$

Then $x^2 + y^2 = x^2 + y^2 + 0$

Then $x^2 + y^2 = x^2 + y^2 + 2xy$

Then $x^2 + y^2 = (x + y)^2$

Therefore $x = 0 \rightarrow x^2 + y^2 = (x + y)^2$

Case 2

Assume $y = 0$

Then $xy = 0$

Then $2xy = 0$

Then $x^2 + y^2 = x^2 + y^2 + 0$

Then $x^2 + y^2 = x^2 + y^2 + 2xy$

Then $x^2 + y^2 = (x + y)^2$

Therefore $y = 0 \rightarrow x^2 + y^2 = (x + y)^2$

Therefore $x = 0 \vee y = 0 \rightarrow x^2 + y^2 = (x + y)^2$

Therefore $\forall x, y \in \mathbb{R} : x^2 + y^2 = (x + y)^2 \Leftrightarrow x = 0 \vee y = 0$

2. Let $x, y \in \mathbb{R}$

First, prove $x^3 + x^2y = y^2 + xy \rightarrow y = x^2 \vee y = -x$

Assume $x^3 + x^2y = y^2 + xy$

Then $x^2(x + y) = y(x + y)$

Case 1

Assume $x + y \neq 0$

Then $x^2 = y$

Therefore $x^3 + x^2y = y^2 + xy \rightarrow y = x^2$

Case 2

Assume $x + y = 0$

Then $y = -x$

Therefore $x^3 + x^2y = y^2 + xy \rightarrow y = -x$

Therefore $x^3 + x^2y = y^2 + xy \rightarrow y = x^2 \vee y = -x$

Second, prove $y = x^2 \vee y = -x \rightarrow x^3 + x^2y = y^2 + xy$

Assume $y = x^2 \vee y = -x$

Case 1

Assume $y = x^2$

Then $y(x + y) = x^2(x + y)$

Then $xy + y^2 = x^3 + x^2y$

Therefore $y = x^2 \rightarrow x^3 + x^2y = xy + y^2$

Case 2

Assume $y = -x$

Then $x + y = 0$

Then $y(x + y) = y \times 0 = 0$

Then $x^2(x + y) = x^2 \times 0 = 0$

Then $y(x + y) = x^2(x + y) = 0$

Then $xy + y^2 = x^3 + x^2y = 0$

Therefore $y = -x \rightarrow x^3 + x^2y = xy + y^2$

Therefore $y = x^2 \vee y = -x \rightarrow x^3 + x^2y = y^2 + xy$

Therefore $\forall x, y \in \mathbb{R} : x^3 + x^2y = y^2 + xy \Leftrightarrow y = x^2 \vee y = -x$