

CS5489

Lecture 4.1: Duality

Kede Ma

City University of Hong Kong (Dongguan)



香港城市大學（東莞）
City University of Hong Kong
(Dongguan)

Slide template by courtesy of Benjamin M. Marlin
Part of slides are borrowed from Stephen P. Boyd

Outline

1 Review

2 Convex Optimization Problems

3 Duality

Convex Set

- **Line segment** between $\mathbf{x}^{(1)}$ and $\mathbf{x}^{(2)}$: all points

$$\mathbf{x} = \alpha \mathbf{x}^{(1)} + (1 - \alpha) \mathbf{x}^{(2)}$$

with $0 \leq \alpha \leq 1$

- **Convex set**: contains line segment between any two points in the set

$$\mathbf{x}^{(1)}, \mathbf{x}^{(2)} \in \mathcal{X}, 0 \leq \alpha \leq 1 \implies \alpha \mathbf{x}^{(1)} + (1 - \alpha) \mathbf{x}^{(2)} \in \mathcal{X}$$

Convex Function


- $f : \mathbb{R}^N \mapsto \mathbb{R}$ is convex if $\text{dom}(f)$ is a convex set and

$$f(\alpha \mathbf{x} + (1 - \alpha) \mathbf{y}) \leq \alpha f(\mathbf{x}) + (1 - \alpha) f(\mathbf{y})$$


for all $\mathbf{x}, \mathbf{y} \in \text{dom}(f)$, $0 \leq \alpha \leq 1$

- This is also called Jensen's inequality
- f is concave if $-f$ is convex

Convex



Non-Convex



First-Order Condition

- f is **differentiable** if the gradient

$$\nabla f(\mathbf{x}) = \left[\frac{\partial f(\mathbf{x})}{\partial x_1}, \frac{\partial f(\mathbf{x})}{\partial x_2}, \dots, \frac{\partial f(\mathbf{x})}{\partial x_N} \right]^T$$

exists at each $\mathbf{x} \in \text{dom}(f)$

- **1st-order condition:** differentiable f with convex domain is convex iff

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x}), \quad \text{for all } \mathbf{x}, \mathbf{y} \in \text{dom}(f)$$

- First-order approximation of f is global underestimator

Second-Order Condition

- f is **twice differentiable** if the Hessian $\nabla^2 f(\mathbf{x}) \in \mathbb{S}^N$

$$\nabla^2 f(\mathbf{x})_{ij} = \frac{\partial^2 f(\mathbf{x})}{\partial x_i \partial x_j}$$

exists at each $\mathbf{x} \in \text{dom}(f)$

- **2nd-order condition:** twice differentiable f with convex domain is convex iff

$$\nabla^2 f(\mathbf{x}) \succeq 0$$

- This means that the Hessian matrix is positive semidefinite

$$\mathbf{A} \succeq 0 \iff \forall \mathbf{x}, \mathbf{x}^T \mathbf{A} \mathbf{x} \geq 0$$

Outline

1 Review

2 Convex Optimization Problems

3 Duality

Optimization Problem in Standard Form



$$\begin{aligned} & \underset{\mathbf{x}}{\text{minimize}} && f_0(\mathbf{x}) \\ & \text{subject to} && f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, r \\ & && h_i(\mathbf{x}) = 0, \quad i = 1, \dots, s \end{aligned}$$

- $\mathbf{x} \in \mathbb{R}^N$ is the optimization variable
- $f_0 : \mathbb{R}^N \mapsto \mathbb{R}$ is the objective or cost function
- $f_i : \mathbb{R}^N \mapsto \mathbb{R}, i = 1, \dots, r$, are the inequality constraint functions
- $h_i : \mathbb{R}^N \mapsto \mathbb{R}, i = 1, \dots, s$ are the equality constraint functions

■ Optimal value:

$$p^* = \min\{f_0(\mathbf{x}) \mid f_i(\mathbf{x}) \leq 0, i = 1, \dots, r, h_i(\mathbf{x}) = 0, i = 1, \dots, s\}$$

- $p^* = \infty$ if problem is infeasible (no \mathbf{x} satisfies the constraints)
- $p^* = -\infty$ if the problem is unbounded below

Optimal and Locally Optimal Points

- \mathbf{x} is **feasible** if $\mathbf{x} \in \text{dom}(f_0)$ and it satisfies the constraints
- A feasible \mathbf{x} is optimal if $f_0(\mathbf{x}) = p^*$
- \mathbf{x} is locally optimal if there is an $R > 0$ such that \mathbf{x} is optimal for

$$\begin{aligned} & \underset{\mathbf{z}}{\text{minimize}} && f_0(\mathbf{z}) \\ & \text{subject to} && f_i(\mathbf{z}) \leq 0, \quad i = 1, \dots, r \\ & && h_i(\mathbf{z}) = 0, \quad i = 1, \dots, s \\ & && \|\mathbf{z} - \mathbf{x}\|_2 \leq R \end{aligned}$$

- Examples:

- $f_0(x) = -\log x$, $\text{dom}(f_0) = \mathbb{R}_{++}$: $p^* = -\infty$
- $f_0(x) = x \log x$, $\text{dom}(f_0) = \mathbb{R}_{++}$: $p^* = -1/e$, $x = 1/e$ is optimal
- $f_0(x) = x^3 - 3x$: $p^* = -\infty$, locally optimum at $x = 1$

Feasibility Problem



$$\begin{array}{ll}\text{find} & \mathbf{x} \\ \text{subject to} & f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, r \\ & h_i(\mathbf{x}) = 0, \quad i = 1, \dots, s\end{array}$$

can be considered a special case of the general problem with $f_0(\mathbf{x}) = 0$:

$$\begin{array}{ll}\text{minimize}_{\mathbf{x}} & 0 \\ \text{subject to} & f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, r \\ & h_i(\mathbf{x}) = 0, \quad i = 1, \dots, s\end{array}$$

- $p^* = 0$ if constraints are feasible; any feasible \mathbf{x} is optimal
- $p^* = \infty$ if constraints are infeasible

Convex Optimization Problem in Standard Form



$$\begin{aligned} &\text{minimize} && f_0(\mathbf{x}) \\ &\text{subject to} && f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, r \\ & && \mathbf{a}_i^T \mathbf{x} = b_i, \quad i = 1, \dots, s \end{aligned}$$

- f_0, f_1, \dots, f_r are convex
- Equality constraints are affine

■ Often written as

$$\begin{aligned} &\underset{\mathbf{x}}{\text{minimize}} && f_0(\mathbf{x}) \\ &\text{subject to} && f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, r \\ & && \mathbf{A}\mathbf{x} = \mathbf{b} \end{aligned}$$

- Important property: feasible set of a convex optimization problem is convex

Example



$$\begin{aligned} \text{minimize} \quad & f_0(\mathbf{x}) = x_1^2 + x_2^2 \\ \text{subject to} \quad & f_1(\mathbf{x}) = x_1/(1 + x_2^2) \leq 0 \\ & h_1(\mathbf{x}) = (x_1 + x_2)^2 = 0 \end{aligned}$$

- f_0 is convex
- Feasible set $\{(x_1, x_2) | x_1 = -x_2 \leq 0\}$ is convex
- Not a convex problem (in standard form): f_1 is not convex, h_1 is not affine
- Writes it as a convex problem in standard form

$$\begin{aligned} \text{minimize} \quad & f_0(\mathbf{x}) = x_1^2 + x_2^2 \\ \text{subject to} \quad & f_1(\mathbf{x}) = x_1 \leq 0 \\ & h_1(\mathbf{x}) = x_1 + x_2 = 0 \end{aligned}$$

Local and Global Optima in Convex Optimization

- Any locally optimal point is (globally) optimal

Proof. Suppose \mathbf{x} is locally optimal and \mathbf{y} is optimal with $f_0(\mathbf{y}) < f_0(\mathbf{x})$. Locally optimal \mathbf{x} means there is an $R > 0$ such that

$$\mathbf{z} \text{ feasible, } \|\mathbf{z} - \mathbf{x}\|_2 < R \implies f_0(\mathbf{z}) \geq f_0(\mathbf{x}).$$

Consider $\mathbf{z} = \alpha\mathbf{y} + (1 - \alpha)\mathbf{x}$, with $\alpha = R/(2\|\mathbf{y} - \mathbf{x}\|_2)$

- $\|\mathbf{y} - \mathbf{x}\|_2 > R$, so $0 < \alpha < 1/2$
- \mathbf{z} is a convex combination of two feasible points, hence feasible
- $\|\mathbf{z} - \mathbf{x}\|_2 = R/2$ and

$$f_0(\mathbf{z}) \leq \alpha f_0(\mathbf{y}) + (1 - \alpha)f_0(\mathbf{x}) < f_0(\mathbf{x})$$

which contradicts our assumption that \mathbf{x} is locally optimal

Outline

1 Review

2 Convex Optimization Problems

3 Duality

Lagrangian

- Standard form problem (not necessarily convex):

$$\begin{aligned} \min_{\mathbf{x}} \quad & f_0(\mathbf{x}) \\ \text{subject to} \quad & f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, r \\ & h_i(\mathbf{x}) = 0, \quad i = 1, \dots, s \end{aligned}$$

- Variable $\mathbf{x} \in \mathbb{R}^N$, domain \mathcal{X} , optimal value p^*
- **Lagrangian:** $L : \mathbb{R}^N \times \mathbb{R}^r \times \mathbb{R}^s \mapsto \mathbb{R}$, with
 $\text{dom}(L) = \mathcal{X} \times \mathbb{R}^r \times \mathbb{R}^s$

$$L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = f_0(\mathbf{x}) + \sum_{i=1}^r \lambda_i f_i(\mathbf{x}) + \sum_{i=1}^s \nu_i h_i(\mathbf{x})$$

- Weighted sum of objective and constraint functions
- λ_i is Lagrange multiplier associated with $f_i(\mathbf{x}) \leq 0$
- ν_i is Lagrange multiplier associated with $h_i(\mathbf{x}) = 0$

Lagrange Dual Function

- **Lagrange dual function:** $g : \mathbb{R}^r \times \mathbb{R}^s \mapsto \mathbb{R}$,

$$g(\boldsymbol{\lambda}, \boldsymbol{\nu}) = \inf_{\mathbf{x} \in \mathcal{X}} L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = \inf_{\mathbf{x} \in \mathcal{X}} \left(f_0(\mathbf{x}) + \sum_{i=1}^r \lambda_i f_i(\mathbf{x}) + \sum_{i=1}^s \nu_i h_i(\mathbf{x}) \right)$$

g is concave and can be $-\infty$ for some $\boldsymbol{\lambda}$ and $\boldsymbol{\nu}$

- **Lower bound property:** if $\boldsymbol{\lambda} \geq 0$, then $g(\boldsymbol{\lambda}, \boldsymbol{\nu}) \leq p^*$

Proof. If $\tilde{\mathbf{x}}$ is feasible and $\boldsymbol{\lambda} \geq 0$, then

$$f_0(\tilde{\mathbf{x}}) \geq L(\tilde{\mathbf{x}}, \boldsymbol{\lambda}, \boldsymbol{\nu}) \geq \inf_{\mathbf{x} \in \mathcal{X}} L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = g(\boldsymbol{\lambda}, \boldsymbol{\nu})$$

Minimizing over all feasible $\tilde{\mathbf{x}}$ gives $p^* \geq g(\boldsymbol{\lambda}, \boldsymbol{\nu})$

Least-Norm Solution of Linear Equations



$$\begin{array}{ll}\underset{\mathbf{x}}{\text{minimize}} & \mathbf{x}^T \mathbf{x} \\ \text{subject to} & \mathbf{A}\mathbf{x} = \mathbf{b}\end{array}$$

■ Dual function:

- Lagrangian is $L(\mathbf{x}, \boldsymbol{\nu}) = \mathbf{x}^T \mathbf{x} + \boldsymbol{\nu}^T (\mathbf{A}\mathbf{x} - \mathbf{b})$
- To minimize L over \mathbf{x} , set gradient equal to zero:

$$\nabla_{\mathbf{x}} L(\mathbf{x}, \boldsymbol{\nu}) = 2\mathbf{x} + \mathbf{A}^T \boldsymbol{\nu} = 0 \implies \mathbf{x} = -(1/2)\mathbf{A}^T \boldsymbol{\nu}$$

- Plug in L to obtain g :

$$g(\boldsymbol{\nu}) = L((-1/2)\mathbf{A}^T \boldsymbol{\nu}, \boldsymbol{\nu}) = -\frac{1}{4} \boldsymbol{\nu}^T \mathbf{A} \mathbf{A}^T \boldsymbol{\nu} - \mathbf{b}^T \boldsymbol{\nu}$$

a concave function of $\boldsymbol{\nu}$

- Lower bound property: $p^* \geq -\frac{1}{4} \boldsymbol{\nu}^T \mathbf{A} \mathbf{A}^T \boldsymbol{\nu} - \mathbf{b}^T \boldsymbol{\nu}$ for all $\boldsymbol{\nu}$

Standard Form Linear Programming



$$\begin{aligned} & \underset{\mathbf{x}}{\text{minimize}} && \mathbf{c}^T \mathbf{x} \\ & \text{subject to} && \mathbf{Ax} = \mathbf{b}, \quad \mathbf{x} \succeq 0 \end{aligned}$$

■ Dual function:

■ Lagrangian is

$$\begin{aligned} L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) &= \mathbf{c}^T \mathbf{x} + \boldsymbol{\nu}^T (\mathbf{Ax} - \mathbf{b}) - \boldsymbol{\lambda}^T \mathbf{x} \\ &= -\mathbf{b}^T \boldsymbol{\nu} + (\mathbf{A}^T \boldsymbol{\nu} - \boldsymbol{\lambda} + \mathbf{c})^T \mathbf{x} \end{aligned}$$

■ L is affine in \mathbf{x} , hence:

$$g(\boldsymbol{\lambda}, \boldsymbol{\nu}) = \inf_{\mathbf{x}} L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = \begin{cases} -\mathbf{b}^T \boldsymbol{\nu} & \mathbf{A}^T \boldsymbol{\nu} - \boldsymbol{\lambda} + \mathbf{c} = 0 \\ -\infty & \text{otherwise} \end{cases}$$

g is linear on $\{(\boldsymbol{\lambda}, \boldsymbol{\nu}) | \mathbf{A}^T \boldsymbol{\nu} - \boldsymbol{\lambda} + \mathbf{c} = 0\}$, hence concave

■ Lower bound property: $p^* \geq -\mathbf{b}^T \boldsymbol{\nu}$ if $\mathbf{A}^T \boldsymbol{\nu} + \mathbf{c} \succeq 0$

The Dual Problem

■ Lagrange dual problem:

$$\begin{array}{ll}\max_{\lambda, \nu} & g(\lambda, \nu) \\ \text{subject to} & \lambda \geq 0\end{array}$$

- λ, ν are dual feasible if $\lambda \succeq 0, (\lambda, \nu) \in \text{dom}(g)$
 - Finds best lower bound p^* , obtained from Lagrange dual problem
 - A convex optimization problem; optimal value denoted d^*
- ## ■ Weak duality: $d^* \leq p^*$
- Always holds (for convex and nonconvex problems)
 - Can be used to find nontrivial lower bounds for difficult problems
- ## ■ Strong duality: $d^* = p^*$
- Does not hold in general
 - (Usually) holds for convex problems, *e.g.*, SVM

Complementary Slackness

- Assuming strong duality holds, \mathbf{x}^* is primal optimal, $(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*)$ is dual optimal

$$\begin{aligned} f_0(\mathbf{x}^*) = g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^r \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^s \nu_i^* h_i(\mathbf{x}) \right) \\ &\leq f_0(\mathbf{x}^*) + \sum_{i=1}^r \lambda_i^* f_i(\mathbf{x}^*) + \sum_{i=1}^s \nu_i^* h_i(\mathbf{x}^*) \\ &\leq f_0(\mathbf{x}^*) \end{aligned}$$

- Hence, the two inequalities hold with equality
 - \mathbf{x}^* not only minimizes $f_0(\mathbf{x})$, but also minimizes $L(\mathbf{x}, \boldsymbol{\lambda}^*, \boldsymbol{\nu}^*)$
 - $\lambda_i^* f_i(\mathbf{x}^*) = 0$ for $i = 1, \dots, r$ (complementary slackness):

$$\lambda_i^* > 0 \implies f_i(\mathbf{x}^*) = 0, \quad f_i(\mathbf{x}^*) < 0 \implies \lambda_i^* = 0$$

Karush-Kuhn-Tucker (KKT) Conditions

- The following four conditions are called KKT conditions (for a problem with differentiable f_i, h_i)
 - 1 Primal constraints: $f_i(\mathbf{x}) \leq 0, i = 1, \dots, r, h_i(\mathbf{x}) = 0, i = 1, \dots, s$
 - 2 Dual constraints: $\lambda \succeq 0$
 - 3 Complementary slackness: $\lambda_i f_i(\mathbf{x}) = 0, i = 1, \dots, r$
 - 4 Gradient of Lagrangian with respect to \mathbf{x} vanishes:

$$\nabla f_0(\mathbf{x}) + \sum_{i=1}^r \lambda_i \nabla f_i(\mathbf{x}) + \sum_{i=1}^s \nu_i \nabla h_i(\mathbf{x}) = 0$$

- If strong duality holds and \mathbf{x}, λ, ν are optimal, then they must satisfy the KKT conditions