

MATH 240: Discrete Structures 1 Assignment #1

Monorina Mukhopadhyay (ID: 260364335)

September 28, 2017

Problem 1

a Truth Table

P	Q	$\neg P$	$\neg P \vee Q$	$P \implies Q$	$(P \implies Q) \iff (\neg P \vee Q)$
T	T	F	T	T	T
T	F	F	F	F	T
F	T	T	T	T	T
F	F	T	T	T	T

This is a tautology

b Truth Table This is a tautology

P	Q	$\neg P$	$\neg Q$	$\neg P \wedge \neg Q$	$\neg(\neg P \wedge \neg Q)$	$P \vee Q$	$\neg(\neg P \wedge \neg Q) \iff P \vee Q$
T	T	F	F	F	T	T	T
T	F	F	T	F	T	T	T
F	T	T	F	F	T	T	T
F	F	T	T	T	F	F	T

c Truth Table This is a tautology

P	Q	$\neg P$	$\neg Q$	$P \iff Q$	$P \wedge Q$	$\neg P \wedge \neg Q$	$((P \wedge Q) \vee (\neg P \wedge \neg Q))$	$(P \iff Q) \iff ((P \wedge Q) \vee (\neg P \wedge \neg Q))$
T	T	F	F	T	T	F	T	T
T	F	F	T	F	F	F	F	T
F	T	T	F	F	F	F	F	T
F	F	T	T	T	F	T	T	T

d This is a tautology

X	Y	$X \implies Y$	$\neg(X \implies Y)$	$Y \implies X$	$(\neg(X \implies Y)) \implies (Y \implies X)$
T	T	T	F	T	T
T	F	F	T	T	T
F	T	T	F	F	T
F	F	T	F	T	T

e This is a contingency

P	Q	R	$\neg P$	$\neg Q$	$\neg P \wedge \neg Q$	$R \implies Q$	$(\neg P \wedge \neg Q) \implies (R \implies Q)$
T	T	T	F	F	F	T	T
T	T	F	F	F	F	T	T
T	F	T	F	T	F	F	T
T	F	F	F	T	F	T	T
F	T	T	T	F	F	T	T
F	T	F	T	F	F	T	T
F	F	T	T	T	T	F	F
F	F	F	T	T	T	T	T

Problem 2

- a The left side of the equation is: $(X \implies Y) \vee (X \implies Z)$
 $\equiv (\neg X \vee Y) \vee (\neg X \vee Z)$ Condition Identity
 $\equiv (\neg X \vee Y \vee \neg X) \vee (\neg X \vee Y \vee Z)$ Associative Property
 $\equiv (\neg X \vee Y) \vee (\neg X \vee Y \vee Z)$ Idempotent Identity
 $\equiv (\neg X \neg X \vee Y \vee Y) \vee (Z)$ Associative Identity
 $\equiv (\neg X \vee Y) \vee (Z)$ Idempotent Identity
 $\equiv \neg X \vee (Y \vee Z)$ Associative Identity
 $\equiv X \implies (Y \vee Z)$ Conditional Identity
- b Starting from the left side again: $(P \iff Q) \equiv (P \implies Q) \wedge (Q \implies P)$ Biconditional Identity
 $\equiv (P \implies Q) \wedge (\neg Q \vee P)$ Conditional Identity
 $\equiv (P \implies Q) \wedge (\neg(\neg P) \vee \neg Q)$ Commutative and Double Negative Properties
 $\equiv (P \implies Q) \wedge (\neg P \implies \neg Q)$ Conditional Identity

Problem 3

- a In this case, let us assume $P = \text{False}$. Therefore, as shown in class, $(P \Rightarrow Q)$ is always true. Similarly, $P \implies (Q \Rightarrow R)$ is always True. However, we can find R such that $(P \Rightarrow Q) \Rightarrow R$ is not true, that is, when R is False. So, for $P = \text{False}$, $R = \text{False}$, the two expressions are not the same and therefore the expressions are logically different.
- b To show that the two expressions: $(X \wedge Y) \vee Z$ and $X \wedge (Y \vee Z)$ are not equivalent, let $X = \text{False}$. Now, if Z is picked as True, $(X \wedge Y)$ is False, and $(X \wedge Y) \vee Z$ is True. On the other hand, $X \wedge (Y \vee Z)$ is False since X is False. Therefore, the two sides are not equivalent here.

Problem 4

- a $D \Rightarrow H$
- b $\neg P \wedge \neg J$
- c $\forall x(K(x) \wedge (x = k)) \implies S$
- d $B \Rightarrow Q$
- e A
- f (If you jump over buildings, then you must be a superhero) $B \Rightarrow H$
- g (Rephrasing: if x can marry my daughter, then x must be a knight) $\forall x M(x, d) \Rightarrow K(x)$
- h (There exists at least four distinct x_i 's, such that $P(x_i)$ holds and the x_i 's are all distinct). This is written as:
 $(\exists x_1 P(x_1)) \wedge (\exists x_2 P(x_2)) \wedge (\exists x_3 P(x_3)) \wedge (\exists x_4 P(x_4)) \wedge (x_1 \neq x_2 \neq x_3 \neq x_4)$
- i For this, let x be the squart root of 5. For a number to be irrational, it cannot be written as a fraction of two integers, ie, there are not both numbers m, n such that $x \neq \frac{m}{n}$
This can be written as: $\forall x(x \cdot x = 5) \implies \neg[\exists m I(m) \wedge \exists n I(n) \wedge (m = n \cdot x)]$

Problem 5

- a Converse: If Susan needs to take Math 240, then she is a sophomore.
 Contrapositive: If Susan is not a sophomore, then she does not need to take Math 240
- b Converse: If one keeps trying, one will succeed.
 Contrapositive: If one does not keep trying, then one will not succeed
- c Converse: If you've paid your library fines, then you can graduate
 Contrapositive: If you haven't paid your library fines, then you can't graduate.

Problem 6

Knights only tell the truth, and knaves only lie.

Let, P : A is a knight

Q : B is a knight

R: C is a knight

A says: B is a knave (1)

B says: A and C are the same (2)

Formulating (1) into a logical expression: $P \iff \neg Q$ (3)

(If and only if A is a knight, then A is telling the truth and B is a knave)

$Q \iff ((P \wedge R) \vee (\neg P \wedge \neg R))$ (4)

If B is a knight, then either A and C are both knights, or A and C are both knaves

From (3): $P \iff \neg Q \equiv (P \wedge \neg Q) \vee (\neg P \wedge Q)$ (From the biconditional identity)

This means either P or Q is true, but not both.

From (4):

$Q \iff ((P \wedge R) \vee (\neg P \wedge \neg R))$

$\equiv Q \wedge ((P \wedge R) \vee (\neg P \wedge \neg R)) \vee \neg Q \wedge \neg((P \wedge R) \vee (\neg P \wedge \neg R))$ (Using the Biconditional identity)

$\equiv Q \wedge ((P \wedge R) \vee (\neg P \wedge \neg R)) \vee \neg Q \wedge ((P \vee R) \wedge (\neg P \vee \neg R))$ (By DeMorgan's laws)

$\equiv Q \wedge ((P \wedge R) \vee (\neg P \wedge \neg R)) \vee \neg Q \wedge ((\neg P \wedge R) \vee (P \wedge \neg R))$ (By distributive and complement and identity laws)

$\equiv P \wedge ((Q \wedge R) \vee (\neg Q \wedge \neg R)) \vee \neg P \wedge ((Q \wedge \neg R) \vee (\neg Q \wedge R))$ (By the distributive and associative laws)

This means, that if P is true, then either Q and R are both true or Q and R are both false \Rightarrow B and C are the same type if

A is a knight \Rightarrow From (1) B must be a knave and so C must be a knave

Alternatively, if P is false, then either Q is true or R is true but not both \Rightarrow B and C are different types if A is a knave \Rightarrow

Q is true (from (3)) i.e., B is a knight and therefore, C must be a knave.

So, in either case, C is a knave

Problem 7

A $\forall x(P(x) \wedge ((\exists yP(y)) \wedge (\exists yR(x, y)) \implies (\exists yP(y))))$

$\equiv \forall x(P(x) \wedge ((\exists y\neg P(y)) \vee \forall y\neg R(x, y) \vee \exists yP(y)))$ (By the deMorgan's law)

$\equiv \forall x(P(x) \wedge T)$ (By commutative and complement and dominant property)

$\equiv \forall x(P(x))$ (By the dominant property again)

Negating the statement gives: $\neg(\forall xP(x)) \equiv \exists x\neg P(x)$ Match (4)

B $\forall x\forall y(\neg R(x, y) \vee \neg P(x))$

The negation is $\exists x\exists y\neg(\neg R(x, y) \vee \neg P(x)) \equiv \exists x\exists y(R(x, y) \wedge P(x))$, using first deMorgan's law, and then double negative rule.

Match: (3)

C $\forall x(P(x) \iff \exists yR(x, y)) \equiv \forall x((P(x) \wedge \exists yR(x, y)) \vee (\neg P(x) \wedge \forall y\neg R(x, y)))$ (From the biconditional rule).

The negation is $\exists x\neg((P(x) \wedge \exists yR(x, y)) \vee (\neg P(x) \wedge \forall y\neg R(x, y)))$

$\equiv \exists x(\neg(P(x) \wedge \exists yR(x, y)) \wedge \neg(\neg P(x) \wedge \forall y\neg R(x, y)))$ (By deMorgan's Thm.)

$\equiv \exists x((\neg P(x) \vee \forall y\neg R(x, y)) \wedge (P(x) \vee \exists yR(x, y)))$ (By deMorgan's Thm.)

$(P(x) \vee \exists yR(x, y)) \wedge (\neg P(x) \vee \forall y\neg R(x, y))$ (By commutative property)

Match (2)

D $\forall x\forall y(P(x) \wedge R(x, y) \Rightarrow R(y, x))$

$\equiv \forall x\forall y((P(x) \wedge R(x, y)) \Rightarrow R(y, x))$ (By order of precedence of operations)

The negation is: $\exists x \exists y \neg(\neg(P(x) \wedge R(x, y)) \vee R(y, x))$ (By Conditional property)
 $\equiv \exists x \exists y (P(x) \wedge R(x, y) \wedge \neg R(y, x))$ (By deMorgan's thm)
 Match (1)

Problem 8

From the truth table, the equation $P * Q$ is the same as $\neg(P \wedge Q)$ First, I will try to reduce the given expression $\neg(X \Rightarrow (\neg X \vee \neg Y))$ into a simpler form.

$$\begin{aligned} & \neg(X \Rightarrow (\neg X \vee \neg Y)) \\ & \equiv \neg(\neg X \vee (\neg X \vee \neg Y)) && \text{By the conditional identity} \\ & \equiv \neg(\neg X \vee \neg Y) && \text{By the idempotent identity} \\ & \equiv (X \wedge Y) && \text{By DeMorgan's law} \end{aligned}$$

Therefore, we need to write $X \wedge Y$ in terms of $P * Q \equiv \neg(P \wedge Q)$

Proof:

$$\begin{aligned} X \wedge Y & \equiv \neg(\neg(X \wedge Y)) && \text{(Property of negation)} \\ & \equiv \neg((\neg(X \wedge Y)) \wedge (\neg(X \wedge Y))) && \text{(Idempotent property)} \\ & \equiv (\neg(X \wedge Y)) * (\neg(X \wedge Y)) && \text{(From the definition of *)} \\ & \equiv (X * Y) * (X * Y) && \text{(From the definition of *)} \end{aligned}$$