

MATH 240: Discrete Structures 1 Assignment #2

Monorina Mukhopadhyay (ID: 260364335)

October 22, 2017

Problem 1

$$N \equiv 2 \pmod{3} \quad (1)$$

$$N \equiv 1 \pmod{5} \quad (2)$$

$$N \equiv 4 \pmod{7} \quad (3)$$

First, I use the Chinese Remainder Theorem to solve (1) and (2). 1 can be written as

$$1 = 3m_1 + 5m_2$$

$$[b = 5, a = 3] \quad 5 = 3x_1 + 2$$

$$[b = 3, a = 2] \quad 3 = 2x_1 + 1$$

$$1 = 3 - 2 = 3 - (5 - 3) = 2 \cdot 3 - 5$$

From the Chinese Remainder Theorem, $x = 3m_1a_2 + 5m_2a_1 = 3 \cdot 2 \cdot 1 - 5 \cdot 2 = -4$ solves (1) and (2)

Since -4 is less than n_1n_2 , and $x \pmod{y} \equiv x$ if $x < y$, $-4 \equiv -4 \pmod{15} \equiv 11 \pmod{15}$

Now, the two equations to solve are:

$$N \equiv 11 \pmod{15}$$

$$N \equiv 4 \pmod{7}$$

Using the Chinese Remainder Theorem again: $[b = 15, a = 7] \quad 15 = 7 \cdot 2 + 1$

$$1 = 15 - 7 \cdot 2$$

$$N = 4(15) - 7(11)(2) = -94 \equiv -94 \pmod{105} \equiv 11 \pmod{105} = 11$$

Check:

$$11 \pmod{3} \equiv 2 \pmod{3}$$

$$11 \pmod{5} \equiv 1 \pmod{5}$$

$$11 \pmod{7} \equiv 4 \pmod{7}$$

Problem 2

a Need to find x, y such that $60 \mid xy$, $60 \nmid x$ and $60 \nmid y$

By inspection, for any $x < 60$ and $y < 60$, such that $xy = 60k$ (where k is an integer), this will work.

For example, for $x = 15$, $y = 4$: $15 \pmod{60} \not\equiv 0 \pmod{60}$ and $4 \pmod{60} \not\equiv 0 \pmod{60}$ but $15 \cdot 4 \pmod{60} \equiv 0 \pmod{60}$.

The same also holds true for $x = 12$, $y = 10$, where $xy \pmod{60} \equiv 0 \pmod{60}$, but $x \pmod{60} \not\equiv 0 \pmod{60}$ and $y \pmod{60} \not\equiv 0 \pmod{60}$

b x, y integers, p prime and $xy \pmod{p} \equiv 0 \implies xy = pt$

Assume p does not divide $x \implies \gcd(p, x) = 1 \implies 1 = m_1p + m_2x$

$$\implies y = m_1py + m_2xy \implies y = m_1py + m_2pt = p(m_1y + m_2t) \implies p \mid y.$$

Since x was picked randomly, the same holds true if x and y are reversed. Therefore, if a prime number divides the product of two integers, it must divide at least one of the two integers.

Problem 3

Given: $m \mid N$ and $n \mid N \implies N = mt_1$ and $N = nt_2$

Since m and n are relatively prime, $\gcd(m, n) = 1 \implies N = a_1mN + a_2nN = a_1mnt_2 + a_2nmt_1 = (a_1t_2 + a_2t_1)mn \implies mn \mid N$

Problem 4

$a > 2$ and $n \geq 1$, $a, n \in \mathbb{Z}$

Need to prove that: $a - 1 \mid a^n - 1$

Proof by induction:

Base case: $a > 2, n = 1; a^n - 1 = a - 1 \implies (a - 1) \mid (a - 1)$ so base case holds

Induction hypothesis: Assume $a - 1 \mid a^k - 1 \implies a^k - 1 = (a - 1)t$ for an integer t

Induction Step: For $k+1$, $a^{k+1} - 1 = a^{k+1} - a^k + a^k - 1 = a^k(a - 1) + (a^k - 1) = a^k(a - 1) + (a - 1)t = (a - 1)(a^k + t) \implies a - 1 \mid a^{k+1} - 1$

Problem 5

$x \in 0, 1, 2, \dots, 38$ For x to satisfy $x^{39} - x \equiv 0 \pmod{39}$ means that $39 \mid x^{39} - x$.

Since $39 = 13 \cdot 3$, and $\gcd(13, 3) = 1$, this means that $13 \mid x^{39} - x$ and $3 \mid x^{39} - x$

This is easily proved as below:

Let $x^{39} - x = N$. $39 \mid N \implies N = 39k = 13 \cdot 3k = 13(3k)$ and $N = 3(13k) \implies 13 \mid N$ and $3 \mid N$ Solving these two equations using the Chinese Remainder Theorem:

$13 = 3(4) + 1 \implies 1 = 13 - 3(4) \implies x = x^{39} \cdot 13 - 3 \cdot 4 \cdot x^{39} = x^{39} \cdot x = x^{39}$ is only true for $x = 0$ or $x = 1$. For any higher x , x^{39} is always $> x$. So there are only two values of x for which the equation holds true.

Problem 6

Using the dynamic programming algorithm for fast algorithm shown in class: $22^{362} = (22^{181})^2$

$$\begin{aligned} &= (22^2)(22^{180})^2 \\ &= 22^2(22^{90})^4 \\ &= 22^2(22^{45})^8 \\ &= 22^2 22^8 (22^{44})^8 \\ &= 22^2 22^8 (22^{22})^{16} \\ &= 22^2 22^8 (22^{11})^{32} \\ &= 22^2 22^8 (22^{32})(22^{10})^{32} \\ &= 22^2 22^8 22^{32} (22^5)^{64} \\ &= 22^2 22^8 22^{32} 22^{64} (22^4)^{64} \\ &= 22^2 22^8 22^{32} 22^{64} (22^2)^{128} \\ &= 22^2 22^8 22^{32} 22^{64} (22)^{256} \cdot 22^{362} \pmod{12} = 22^2 22^8 22^{32} 22^{64} (22)^{256} \pmod{12}. \end{aligned}$$

$$22^1 \pmod{12} = 10 \pmod{12} = 10.$$

$$22^2 \pmod{12} = (22^1 \cdot 22^1) \pmod{12}$$

$$= (22 \pmod{12})(22 \pmod{12}) \pmod{12} \equiv 10 \cdot 10 \pmod{12} \equiv 4 \pmod{12}$$

$$22^4 \pmod{12} = (22^2 \pmod{12})(22^2 \pmod{12}) \pmod{12}$$

$$\equiv 16 \pmod{12} \equiv 4 \pmod{12}$$

$$22^8 \pmod{12} = (22^4 \pmod{12})(22^4 \pmod{12}) \pmod{12} \equiv 16 \pmod{12} \equiv 4 \pmod{12}$$

\implies , all higher powers of 22 will also reduce to 4 mod 12 since they can always be expressed as $(4 \pmod{12})(4 \pmod{12}) \pmod{12}$.

$$22^{362} \pmod{12} = 22^2 22^8 22^{32} 22^{64} (22)^{256} \pmod{12} \equiv (4 \cdot 4 \cdot 4 \cdot 4 \cdot 4) \pmod{12} \equiv 4^5 \pmod{12} \equiv 4 \cdot 4^2 \cdot 4^2 \pmod{12} \equiv 4 \cdot 4 \cdot 4 \pmod{12} \equiv (4 \pmod{12})(4 \pmod{12}) \equiv 4 \pmod{12}.$$