

Homework 2

Physics 201

Due September 11th, 2020

Exercise 1. Let $p = 0$,

$$\frac{d}{dx} \left(\sum_{j=0}^{\infty} a_j(j)x^{j+1} \right) + \sum_{j=0}^{\infty} a_j x^{j+2} - \sum_{j=0}^{\infty} 2a_j x^j = 0$$

$$\sum_{j=0}^{\infty} a_j(j)(j+1)x^j + \sum_{j=0}^{\infty} a_j x^{j+2} - \sum_{j=0}^{\infty} 2a_j x^j = 0$$

$$a_0(0)(1)(x^0) + a_1(1)(2)(x) - 2a_0(x^0) - 2a_1x + \sum_{j=2}^{\infty} a_j(j)(j+1)x^j + \sum_{j=2}^{\infty} a_{j-2}x^j - \sum_{j=2}^{\infty} 2a_j x^j = 0$$

Cancelling and combining terms, and let $a_0 = 0$,

$$\sum_{j=2}^{\infty} x^j (a_j((j)(j+1) - 2) + a_{j-2})$$

$$\sum_{j=0}^{\infty} x^{j+2} (a_{j+2}((j+2)(j+3) - 2) + a_j)$$

Thus the recursion relation is,

$$a_{j+2} = -\frac{a_j}{(j+2)(j+3) - 2}$$

And the first few terms are, for a given a_1 ,

$$a_3 = -\frac{a_1}{10}$$

$$a_5 = -\frac{a_3}{28} = \frac{a_1}{280}$$

$$a_7 = -\frac{a_5}{54} = -\frac{a_1}{1512}$$

Exercise 2. For ease we will set $x = i\omega t$. Using the different forms given we see,

$$\frac{1}{2}Ae^x + \frac{1}{2}Ae^{-x} + \frac{1}{2i}Be^x + \frac{1}{2i}Be^{-x} = \bar{A}e^x + \bar{B}e^{-x}$$

Thus we can derive \bar{A} from A ,

$$\bar{A} = \frac{1}{2}A \left(\frac{e^x + e^{-x}}{e^x} \right) = \frac{1}{2}A \left(1 + \frac{1}{e^{2x}} \right)$$

and

$$\bar{B} = \frac{1}{2i}B \left(\frac{e^x - e^{-x}}{e^x} \right) = \frac{1}{2i}B \left(1 - \frac{1}{e^{2x}} \right)$$

Exercise 3.

$$i^2 = \sqrt{-1}\sqrt{-1} = -1$$

$$i^3 = i^2 \cdot i = -i$$

$$i^4 = i^2 \cdot i^2 = (-1)(-1) = 1$$

$$i^5 = i^3 \cdot i^2 = (-i)(-1) = i$$

Exercise 4. $z = s_z^{i\phi_z}$, $p = s_p e^{i\phi_p}$, $s_z^2 = a^2 + b^2$, $s_p^2 = u^2 + v^2$.

$$\frac{z}{p} = \frac{s_z}{s_p} e^{i\phi_z} e^{i\phi_p} = \frac{z(p^*)}{s_p^2} = \frac{(a+bi)(u-vi)}{u^2+v^2}$$

$$\implies \Re\left(\frac{z}{p}\right) = \frac{au+bv}{u^2+v^2} = \Im\left(\frac{z}{p}\right) = \frac{ub-av}{u^2+v^2}$$

Exercise 5. a.

$$(z+p)^* = ((a+u) + (b+v)i)^* = ((a+u) - (b+v)i) = (a-bi) + (u-vi) = z^* + p^*$$

b.

$$(zp)^* = ((au-bv) + i(av+bu))^* = ((au-bv) - i(av+bu)) = (a-bi)(u-vi) = z^* p^*$$

c.

Following results from 1.5.3,

$$\left(\frac{z}{p}\right)^* = \left(\frac{(a+bi)(u-vi)}{u^2+v^2}\right)^* = \frac{au+bv-i(ub-av)}{u^2+v^2} = \frac{(-a+bi)(-u-vi)}{u^2+v^2} = \frac{-(z^*)(-p)}{s_p^2} = (z^*)\left(\frac{e^{-i\phi_p}}{s_p}\right) = \frac{z^*}{p^*}$$

Exercise 6.

$$\begin{aligned} \cos(\phi + \theta) &= \frac{1}{2}(e^{i(\phi+\theta)} + e^{-i(\phi+\theta)}) \\ &= \frac{1}{2}(e^{i\phi}e^{i\theta} + e^{-i\phi}e^{-i\theta}) \\ &= \frac{1}{2}((\cos(\phi) + i\sin(\phi))(\cos(\theta) + i\sin(\theta)) + (\cos(\phi) - i\sin(\phi))(\cos(\theta) - i\sin(\theta))) \end{aligned}$$

after cancellation of $i\cos(\phi)\sin(\theta)$ and $i\cos(\theta)\sin(\phi)$ terms

$$\begin{aligned} &= \frac{1}{2}(2\cos(\phi)\cos(\theta) - 2\sin(\phi)\sin(\theta)) \\ &= \cos(\phi)\cos(\theta) - \sin(\phi)\sin(\theta) \end{aligned}$$

Exercise 7. Since all sinh covers all of the even terms and cosh covers all the odd, the exponential expression becomes,

$$\sum_{i=0}^{\infty} \frac{x^i}{i!} = \sum_{i=0}^{\infty} \frac{x^{2i}}{(2i)!} + \sum_{i=0}^{\infty} \frac{x^{2i+1}}{(2i+1)!} \implies e^x = \cosh(x) + \sinh(x)$$


Doing similar to before,

$$e^x = \cosh(x) + \sinh(x)$$

$$\pm e^{-x} = \cosh(x) - \sinh(x)$$

$$\implies \cosh(x) = \frac{e^x + e^{-x}}{2}$$

$$\sinh(x) = \frac{e^x - e^{-x}}{2}$$



Exercise 8.

$$\begin{aligned} \sin(i\theta) &= \frac{1}{2i}(e^{i(i)\theta} - e^{-i(i)\theta}) \\ &= \frac{1}{2i}(e^{-\theta} - e^{\theta}) \\ &= -\frac{1}{i} \frac{1}{2}(e^{\theta} - e^{-\theta}) \\ &= -\frac{\sinh(\theta)}{i} \\ &= i \sinh(\theta) \end{aligned}$$

$$\begin{aligned} \sinh(i\eta) &= \frac{1}{2}(e^{i\eta} - e^{-i\eta}) \\ &= i \frac{1}{2i}(e^{i\eta} - e^{-i\eta}) \\ &= i \sinh(\eta) \end{aligned}$$