

# Homework 1

Physics 201

Due September 4th, 2020

**Exercise 1.** Using the general solution

$$x(t) = A \cos(\omega t) + B \sin(\omega t) + a$$

then using the boundary values,


$$x(0) = x_0 = A + a$$

$$x(t^*) = x_* = (x_0 - a) \cos(\omega t^*) + B \sin(\omega t^*) + a \implies B = -\frac{(x_0 - a) \cos(\omega t^*) - x_* + a}{\sin(\omega t^*)}$$

Thus the final solution is

$$x(t) = (x_0 - a) \cos(\omega t) - \frac{(x_0 - a) \cos(\omega t^*) - x_* + a}{\sin(\omega t^*)} \sin(\omega t) + a$$

**Exercise 2.** If  $\omega t^* = n\pi$ , then there arises division by zero because the  $B$  term has  $\sin(n\pi) = 0$  in the denominator. Thus the solution fails.



**Exercise 3.**

**Exercise 4.**

$$m\ddot{x}(t) = F_0 \cos(\omega t) \implies \ddot{x}(t) = \frac{F_0 \cos(\omega t)}{m}$$

Integrate once

$$\int \ddot{x}(t) dt = \int \frac{F_0 \cos(\omega t)}{m} dt \implies \dot{x}(t) = -\frac{F_0}{m\omega} \sin(\omega t) + A$$

Integrating again

$$\int \dot{x}(t)dt = \int -\frac{F_0}{m\omega} \sin(\omega t) + A dt \implies x(t) = -\frac{F_0}{m\omega^2} \cos(\omega t) + At + B$$

Using the boundary conditions,

$$\begin{aligned} x(0) = x_0 = -\frac{F_0}{m\omega^2} + B &\implies B = x_0 + \frac{F_0}{m\omega^2} \\ \dot{x}(0) = v_0 = A & \end{aligned}$$

Thus the final solution is,

$$x(t) = -\frac{F_0}{m\omega^2} \cos(\omega t) + v_0 t + x_0 + \frac{F_0}{m\omega^2}$$

**Exercise 5.** Since we are using Taylor expansion we need to determine an  $x_0$ . For our case we will use 0. Thus,

$$\sin(\Delta x) = f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x + \frac{1}{2}f''(x_0)\Delta x^2 \implies \sin(\Delta x) \approx \Delta x$$

$$\cos(\Delta x) = f(x_0 + \Delta x) \approx f(x_0) + f'(x_0)\Delta x + \frac{1}{2}f''(x_0)\Delta x^2 \implies \cos(\Delta x) \approx 1 - \frac{1}{2}\Delta x^2$$

Thus we get the table

$\Delta x$	$\sin(\Delta x)$	$\cos(\Delta x)$	$\sin(\Delta x)$ approx	$\cos(\Delta x)$ approx
0.1	0.0998	0.9950	0.1	0.995
0.2	0.1987	0.9801	0.2	0.98
0.4	0.3894	0.9211	0.4	0.92
0.8	0.7174	0.6967	0.8	0.68

**Exercise 6.** For the approximation with  $x_0 = 0$  and  $a = 1$ ,

$$f(x_0 + \Delta x) = (1 + x_0 + \Delta x)^n \approx f(x_0) + f'(x_0)\Delta x = 1 + n\Delta x$$

Thus for the examples,

$$\sqrt{1+x} \approx 1 + \frac{1}{2}\Delta x$$

$$\frac{1}{\sqrt{1+x}} \approx 1 - \frac{1}{2}\Delta x$$

$$\frac{1}{(1+x)^2} \approx 1 - 2\Delta x$$

**Exercise 7.** We will use the function  $f(x) = x^{-2}$  with  $x_0 = 10$  and  $\Delta x = 1$ . So our Taylor expansion is

$$f(x_0 + \Delta x) \approx f(x_0) + \Delta f'(x_0) = 0.01 - 2(0.001)\Delta x$$

Thus we can quickly calculate  $1/121$ ,

$$\frac{1}{121} = f(x_0 + \Delta x) \approx 0.01 - 2(0.001)(1) = 0.008$$

Which is close to the true value of 0.00826.