Lec-8


19/09/26 4:55 PM

189. Rotate Array

Solved

Medium · Topics · Companies · Hint

Given an integer array `nums`, rotate the array to the right by `k` steps, where `k` is non-negative.




Example 1:


```
Input: nums = [1,2,3,4,5,6,7], k = 3
Output: [5,6,7,1,2,3,4]
Explanation:
rotate 1 steps to the right: [7,1,2,3,4,5,6]
rotate 2 steps to the right: [6,7,1,2,3,4,5]
rotate 3 steps to the right: [5,6,7,1,2,3,4]
```


Example 2:

```
Input: nums = [-1,-100,3,99], k = 2
Output: [3,99,-1,-100]
Explanation:
rotate 1 steps to the right: [99,-1,-100,3]
rotate 2 steps to the right: [3,99,-1,-100]
```



```
height = [4,2,0,3,2,5]
```






Example 1:

```
Input: nums = [5,0,3,8,6]
Output: 3
Explanation: left = [5,0,3], right = [8,6]
```

Example 2:

```
Input: nums = [1,1,1,0,6,12]
Output: 4
Explanation: left = [1,1,1,0], right = [6,12]
```



134. Gas Station


Solved

Medium · Topics · Companies

There are `n` gas stations along a circular route, where the amount of gas at the `ith` station is `gas[i]`.

You have a car with an unlimited gas tank and it costs `cost[i]` of gas to travel from the `ith` station to its next `(i + 1)th` station. You begin the journey with an empty tank at one of the gas stations.

Given two integer arrays `gas` and `cost`, return *the starting gas station's index if you can travel around the circuit once in the clockwise direction, otherwise return* `-1`. If there exists a solution, it is **guaranteed** to be **unique**.



Example 1:

```
Input: gas = [1,2,3,4,5], cost = [3,4,5,1,2]
Output: 3
```

