

# Polarization Entanglement Distribution in Ensemble-Based Atomic Memories

Bhaskar Mookerji\*

*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

Jeffrey H. Shapiro†

*Massachusetts Institute of Technology, Research Laboratory of Electronics, Cambridge, Massachusetts 02139, USA*

(Dated: February 22, 2012)

Quantum networks enable the long-distance communication of quantum states through teleportation, but require, in advance, the robust distribution of entanglement between relevant parties. Engineering these networks requires quantum interconnects, which convert quantum states in one physical system to those of another reversibly, and with high fidelity. In the following, we describe implementations of long-distance quantum communication networks using polarization entanglement and atomic ensembles. We concisely describe the interactions of a quantum optical field with a heralding atomic ensemble, accounting for multiple-pair events at entanglement generation, as well as finite transmission and photodetection efficiencies under number-resolving and non-resolving photodetection schemes. Using these results, we perform a detailed quantitative performance analysis of quantum networks that distribute and swap entanglement.

## I. INTRODUCTION

Quantum communication exploits quantum mechanical resources, such as entanglement, to achieve tasks un-realizable by classical means, such as accurate teleportation of quantum states and unconditionally secure private key distribution. For such applications, the fundamental problem of quantum communication is the establishment, over optical channels, of entanglement between distant nodes. However, the rate of entanglement distribution to nodes decreases exponentially with channel length. The possibility of creating *scalable* optical quantum networks requires that we overcome this difficulty by storing and processing quantum information locally in quantum memories: first, as repeaters increasing network scalability, and second, as light-matter interfaces to quantum computers [1].

The aim of this thesis is to address the following open problem: the theoretical limits of atomic-ensemble quantum memories that store polarization entanglement. It is part of a larger research program in RLE’s Optical and Quantum Communications Group investigating the system performance of ensemble-based hybrid systems in quantum communication. This thesis marries the formalism of collective interactions of atomic ensembles and quantum optical fields with a number-state analysis of our architecture’s performance. The remainder of this chapter introduces pre-existing memory and repeater architectures forming the foundation of this work, discusses the importance of polarization entanglement to quantum communication, and outlines the contributions made in this thesis.

To date, the theoretical progress of quantum information science and engineering has surpassed its experimental achievements. This thesis paints a detailed picture

of what future long-distance quantum communication networks might look like, and their limitations. In the course of this work, we have analyzed quantum networks that distribute polarization entanglement using neutral atomic ensembles. This analysis focuses on three areas. We first abstracted quantized light-ensemble interactions within a heralding atomic memory and determined that these interactions preclude a Gaussian state analysis of local entanglement distribution (Chapter ??, Appendix ??). After describing the potential losses in an entanglement distribution architecture, we applied the SU(2) representation of beam splitter operators to model loss in a number state basis, and performed a Gaussian state analysis of polarization entanglement swapping (Chapter II). A number-state analysis of the entanglement distribution captures the joint state of the heralding light and atomic excitations, accounting for imperfections in transmission loss, photodetection efficiency and counting resolution, and multiple-pair events at the downconversion source.

The numerical characterization of entanglement distribution and connection presented in this thesis is so far consistent with our physical intuition of how such networks should behave. In particular, the probability of a successful heralding event is independent of whether a single, significant, uniform efficiency loss is located either before or after an ensemble memory, or during photodetection. With regards to the probability measuring a single photon, all losses are effectively the same. The same is not true for fidelity. A uniform transmission loss between the entanglement source and the ensembles will significantly reduce the likelihood that you’ve stored a polarization Bell state, but significant quantum efficiency losses at photodetection are less likely to diminish that fidelity. Increasing the pump power at the entanglement source increases the likelihood of multiple-pair events, increasing the heralding probability but making the fidelity more sensitive to transmission and photodetection efficiency losses. These results are true for both

---

\* [mookerji@mit.edu](mailto:mookerji@mit.edu)

† [jhs@mit.edu](mailto:jhs@mit.edu)

number-resolving and non-resolving photodetectors. For both entanglement distribution and connection, heralding probabilities for non-resolving detectors are higher, but the fidelities are significantly lower. Many of these same issues appear in entanglement swapping as well.

Using Gaussian-state outputs from the OPA's, we can quantify fidelity loss in entanglement distribution by accounting for multiple-pair effects, fiber propagation loss, photodetection limitations, and phase mismatch between atomic ensembles. A Gaussian-state analysis for finite atomic ensembles, while valid for the DLCZ protocol, is not applicable for the distribution of externally-generated entanglement. The discussion in this thesis focuses on building a new abstraction for quantum memories and a corresponding formalism for its architectural analysis.

The remainder of the thesis is organized as follows:

- **Chapter ??** summarizes the atomic physics concerning the interactions between multi-atom ensembles and quantized light fields. Sections ?? and ?? formally introduce quantized fields and common assumptions regarding their interactions with collective atomic states, particularly in the context of three-level Raman interactions. In Section ??, we apply this formalism to derive the trilinear Hamiltonian describing a heralding DLCZ-type memory.
- **Chapter II** synthesizes the primary theoretical results of this thesis: an architectural analysis of a DLCZ-type quantum communication system, driven by a dual-OFA source, with particular emphasis paid to multiple-pair effects. In Section II.1, we provide an overview of this architecture and discuss relevant experimental parameters for a prototype system. Assuming perfect absorption by the memories, we determine the fidelity and success probability of our heralding architecture in Section ?? by applying a number-state analysis. Section III applies a Gaussian state analysis to polarization entanglement connection.
- **Chapter IV** summarizes our theoretical results and discusses possible future directions for studying architectures for quantum communication with polarization entanglement.
- **Appendix ??** presents a Quantum Monte Carlo simulation of the pump, signal, and idler modes of a trilinear Hamiltonian. We'll use this simulation to: (i) justify an *ansatz* solution in the Chapter II's number-state analysis; and (ii) explain why approximate solutions in the literature fail to adequately describe memory-loading dynamics. Section ?? argues that trilinear Hamiltonians are not analytically diagonalizable using a finite Lie group representation. Section ?? lists the code for the numerical simulation of the trilinear Hamiltonian's associated master equation.




FIG. 1. Components of a quantum repeater node in the MIT-NU architecture. (a) Parametric downconversion creates pairs of polarization-entangled photons, sending the idler photon to atom-trap 1 and the signal photon to atom-trap 2. Each trap contains a single ultra-cold rubidium atom cooled to its hyperfine ground state. In the energy level diagram, the AB-transition absorbs 795 nm photons, and the BD-transition is coherently driven, thereby enabling storage at D. (b) Polarization-entangled photon pairs generated by a pair of type-II optical parametric amplifiers (OPAs) and a polarizing beam splitter (PBS). The polarizations  $\hat{x}$  and  $\hat{y}$  are denoted by arrows and bullets, respectively. Figures taken from [2] and [3].

### I.1. Architectures for Long-Distance Quantum Networks

An illustration of a model quantum communication system is shown in Fig. 1, in the case of a single-trapped atom. Through type-II parametric downconversion, a post-selected maximally-entangled state can be produced of the form,

$$|\psi_1\rangle = \frac{1}{\sqrt{2}} (|\sigma_+\rangle_1 |\sigma_-\rangle_2 + e^{i\phi} |\sigma_-\rangle_1 |\sigma_+\rangle_2) \quad (1)$$

where  $\sigma_+$  ( $\sigma_-$ ) indicates right (left) circular polarization, and  $\phi$  is a phase offset. An arbitrary polarization of any photon entering the cavity can be stored in the basis of right and left circular polarizations, such that

$$|\psi_2\rangle = \alpha |\sigma_+\rangle + \beta |\sigma_-\rangle. \quad (2)$$

Through a Raman  $\Lambda$ -type interaction, a signal or idler photon effectively transfers its entanglement to the degenerate B magnetic hyperfine levels, and subsequently




FIG. 2. Entanglement with the DLCZ protocol. (Left) Weak and strong coherent pulses induce writing and reading through spontaneous Raman transitions, respectively. (Right) Measurement-induced interference results in a single-excitation entangled state. Figures taken from [1] and [4].

to the D levels through a coherently-driven transition. However, the efficient coupling of a single photon to a single trapped atom is a daunting technical task, requiring that the atom be held in an ultrahigh- $Q$  cavity. By using an atomic ensemble, we eliminate the need for such a high-quality cavity because a collective atomic state is easily produced by a single photon. Such a state is one in which a single atom has been excited from its ground state  $|g\rangle$  at A to the metastable state  $|s\rangle$  at D.

To illustrate this behavior for an atomic ensemble, consider  $N_a$  atoms prepared in their ground states, a collective state denoted by  $|0\rangle_a = |g\rangle^{\otimes N_a}$ . Coherently pumping the ensemble creates an inelastic Raman scattering event that is collectively enhanced by constructive interference within the ensemble [5]. The resulting forward-scattered Stokes light arises from coherent spontaneous emission in the ensemble, and the correlated ensemble excitation is

a collective spin state,

$$|1\rangle_a = \hat{S}^\dagger |0\rangle_a = \frac{1}{\sqrt{N_a}} \sum_{i=1}^{N_a} |g\rangle_1 \cdots |s\rangle_i \cdots |g\rangle_{N_a}. \quad (3)$$

where  $\hat{S} = (1/\sqrt{N_a}) \sum_i |g\rangle_i \langle s|$ . Because the excitation is composed of many atoms, the collective spin excitation is protected against the loss of individual atoms in the ensemble, increasing its robustness for storage. In the weak interaction limit, in which most of the atoms remain in their ground state, the spin excitation  $\hat{S}$  is effectively a ladder operator, as  $[\hat{S}, \hat{S}^\dagger] = \sum_i (|g\rangle_i \langle g| - |s\rangle_i \langle s|)/N_a \approx 1$ , and the outgoing Stokes light and spin excitation are in a two-mode squeezed state [4]. Generally, the term with the  $n$ th atom in  $|s\rangle$  acquires the phase  $e^{i(\mathbf{k}_w - \mathbf{k}_s) \cdot \mathbf{x}_n}$ , where  $\mathbf{k}_w$  is the wave vector of pump field,  $\mathbf{k}_s$  is that of the detected Stokes photon, and  $\mathbf{x}_n$  is the  $n$ th atom's position. Collective excitations can be read out very efficiently when converted into single anti-Stokes photons, which are emitted into a well-defined mode because of collective interference. A resonant laser excitation of the collective state in Eqn. 3 results in  $N_A - 1$  atoms in  $|g\rangle$  and one delocalized excitation in  $|e\rangle$ . Through decay to the  $|g\rangle^{\otimes N_a}$ , an anti-Stokes photon is emitted along the  $|e\rangle - |g\rangle$  transition. Denoting  $\mathbf{k}_{as}$  and  $\mathbf{k}_r$  as the wave vectors of the anti-Stokes photon and read laser, respectively, satisfying the phase matching condition  $\mathbf{k}_s + \mathbf{k}_{as} = \mathbf{k}_r + \mathbf{k}_w$  results in a very high probability amplitude for the anti-Stokes photon to be in the  $\mathbf{k}_r + \mathbf{k}_w - \mathbf{k}_s$  direction due to constructive interference [6].

Our analysis merges the approaches of trapped single atoms in cavity quantum electrodynamics (QED) proposed by MIT and Northwestern University (MIT-NU) and the ensemble-based repeater architecture proposed by Duan, Lukin, Cirac, and Zoller (DLCZ) [2, 4]. The MIT-NU and DLCZ protocols both utilize spontaneous Raman transitions to mediate atomic storage. Whereas the MIT-NU protocol has the advantage of storing externally generated entanglement and verifying its success through cycling-transition fluorescence, it is prohibitively difficult to implement because of the strong coupling requirements of cavity QED.

In contrast, the DLCZ protocol creates a collective atomic excitation, as in Eqn. 3, not by an external input photon, but by the ensemble itself interacting with a classical (write) field. The entangled state is generated probabilistically (but heralded) through postselection and measurement quantum interference, as shown in Fig. 2. In the ideal case of low excitation probability, a photodetection event at either of the two detectors projects the two ensembles into a maximally-entangled singlet state of excitations. Although scalably resilient to issues that might plague such protocols, such as propagation loss and photodetector dark counts, DLCZ requires stable phase coherence and number-resolving photodetectors, neither of which are easy to implement in practice. By enabling the storage of externally-generated entanglement in a DLCZ-type protocol, we will address




FIG. 3. DLCZ with quantum field inputs. (Left) Input-output formalism for a single-sided, two-mode ring cavity. (Right) Interaction in a three-mode parametric amplifier

new error models for entanglement fidelity in quantum memories.

We abstracted a model for the interaction of input quantum field into an ensemble-based quantum memory. A basis for this model is inspired by recent experimental work on heralded single-photon atomic memories and interfaces from [7] [6], which utilized two spatially-overlapping atomic ensembles to absorb arbitrarily polarized single photons. Heralding was observed (at rate of  $10^{-6}$ , using pulsed coherent states ( $\bar{N} \approx 500$ ) with an absorption probability  $\alpha = 0.01$ ). Despite operating in an effective single-photon regime, multiple photon inputs were still present, a problem we wish to analyze in the case of a parametric downconverter input. We consider an ensemble of  $\Lambda$ -type atoms confined in a single-sided, low-finesse ring cavity, as shown in Fig. 3. The  $|g\rangle - |e\rangle$  and  $|e\rangle - |s\rangle$  transitions are coupled to the cavity modes  $\hat{a}$  and  $\hat{b}$ , respectively, each with coupling coefficient  $g_c$ . Under the rotating wave approximation, the interaction Hamiltonian for the collective interaction process is given by

$$\hat{H} = \hbar\Gamma (\hat{a}\hat{S}^\dagger\hat{b}^\dagger + \hat{b}\hat{S}\hat{a}^\dagger) \quad (4)$$

where  $\Gamma = g_c^2 N_a / \Delta$  ( $\Delta$  is the detuning from the two-photon resonance). In general, a Hamiltonian that cannot be analytically diagonalized will not have analytical

dynamics [8, 9], and will therefore not be useful in a Gaussian state analysis. A three-mode interaction describes a heralding quantum memory where an input quantum field (pump mode  $\hat{a}$ ) creates a stationary ensemble excitation (idler mode  $\hat{b}$ ) and a heralding Stokes photon (signal mode  $\hat{S}$ ). Parameterizing the strength of this interaction by  $\Gamma$ , the trilinear Hamiltonian describing this process is then

$$\hat{H} = \hat{H}_0 + \hat{H}_{\text{int}} \quad (5)$$

where

$$\hat{H}_0 = \omega_a \hat{a}^\dagger \hat{a} + \omega_b \hat{b}^\dagger \hat{b} + \omega_S \hat{S}^\dagger \hat{S} \quad (6)$$

$$\hat{H}_{\text{int}} = \Gamma \hat{a} \hat{b}^\dagger \hat{S}^\dagger + \Gamma^* \hat{S} \hat{b} \hat{a} \quad (7)$$

with

$$[\hat{H}_0, \hat{H}_{\text{int}}] = 0. \quad (8)$$

assuming that each mode begins from rest, we can also say that

$$\hat{N}_{abc} = 2\hat{a}^\dagger \hat{a} + \hat{b}^\dagger \hat{b} + \hat{S}^\dagger \hat{S} \quad \hat{N}_{bc} = \hat{b}^\dagger \hat{b} - \hat{S}^\dagger \hat{S} \quad (9)$$

are conserved quantities. assuming that each mode begins from rest, we can also say that

$$\hat{N}_{abc} = 2\hat{a}^\dagger \hat{a} + \hat{b}^\dagger \hat{b} + \hat{S}^\dagger \hat{S} \quad \hat{N}_{bc} = \hat{b}^\dagger \hat{b} - \hat{S}^\dagger \hat{S} \quad (10)$$

are conserved quantities. As an aside, it is worth noting a competing approach to photon storage that may serve as a useful comparison in the future, namely, the usage of stimulated Raman transitions and electromagnetically induced transparency (EIT) to increase the coupling of an input quantum field with an atomic ensemble [10–13]. In this approach, an external coherent control field couples the  $|e\rangle - |s\rangle$  transition in a  $\Lambda$ -type atom, adiabatically reducing the group velocity of a single photon wavepacket and trapping it within the ensemble. Such an approach is deterministic, with high throughput, but admits neither easy verification (as in MIT-NU) nor heralding (as in DLCZ). Like the DLCZ protocol above, its compatibility with externally-generated entanglement is an open question.

## I.2. Polarization Quantum Entanglement and Communication

Because we will be concerned with the behavior of atomic ensembles that are illuminated by the entangled signal and idler produced by optical parametric amplification (OPA), it is important to have an appropriate model for such light beams. As our goal is to quantify the effects of multiple-pair emissions from such a source on the resulting stored entanglement, we cannot immediately default to a postulated biphoton picture. Instead,

we shall use the full Gaussian-state description, in which the the input field in this interaction is one of a pair of polarization-entangled light beams generated by the interference of a pair of anti-phased optical parametric amplifiers as shown in Fig. 1. We assume that the signal and idler cavities are matched with identical linewidths  $\Gamma$ , and pumping fractions,  $G^2$ , of oscillation threshold, with no depletion of or excess noise on the pump [3, 14]. Following interference, the output fields are in an entangled, zero-mean Gaussian pure state with the normally-ordered and phase-sensitive correlation functions

$$\begin{aligned}\langle \hat{A}_{k_j}^\dagger(t+\tau) \hat{A}_{k_j}(t) \rangle &= \frac{G\Gamma}{2} \left[ \frac{e^{-(1-G)\Gamma|\tau|}}{1-G} - \frac{e^{-(1+G)\Gamma|\tau|}}{1+G} \right] \\ \langle \hat{A}_{S_j}(t+\tau) \hat{A}_{I_j}(t) \rangle &= \frac{(-1)^{j-1} G\Gamma}{2} \left[ \frac{e^{-(1-G)\Gamma|\tau|}}{1-G} + \frac{e^{-(1+G)\Gamma|\tau|}}{1+G} \right]\end{aligned}\quad (11)$$

where  $\{\hat{A}_{k_j}(t) e^{-j\omega_k t} : k = S(\text{signal}), I(\text{idler}), j = 1, 2\}$  are the positive-frequency, photon-units OPA-output fields. The low-flux output state of this process at a detuning  $\Delta\omega$  is given by expanding out the number ket representations of the OPAs to first order,

$$\begin{aligned}|\psi\rangle_{SI} &= \sum_n \sqrt{\frac{\bar{N}}{(\bar{N}+1)^{n+1}}} |n\rangle_{S_x} |n\rangle_{I_y} \otimes \sum_n (-1)^n \sqrt{\frac{\bar{N}}{(\bar{N}+1)^{n+1}}} |\text{non-orthogonal quantum states required for teleportation } [17]\rangle_{S_y I_x} \\ &\approx \frac{1}{\bar{N}+1} |0\rangle_{S_x} |0\rangle_{I_y} |0\rangle_{S_y} |0\rangle_{I_x} + \sqrt{\frac{\bar{N}}{(\bar{N}+1)^3}} (|1\rangle_{S_x} |1\rangle_{I_y} |0\rangle_{S_y} |0\rangle_{I_x} + |0\rangle_{S_x} |0\rangle_{I_y} |1\rangle_{S_y} |1\rangle_{I_x}) \\ &= |\text{vac}\rangle + \sqrt{\frac{\bar{N}}{(\bar{N}+1)^3}} (|H\rangle_S |V\rangle_I - |V\rangle_S |H\rangle_I),\end{aligned}\quad (12)$$

where  $\bar{N} = 4G^2 / [(1 - G^2 - \Delta\omega^2/\Gamma^2)^2 + 4\Delta\omega^2/\Gamma^2]$  is the average photon number per mode; and  $|H\rangle_S = |1\rangle_{S_x} |0\rangle_{S_y}$ ,  $|V\rangle_S = |0\rangle_{S_x} |1\rangle_{S_y}$ ,  $|H\rangle_I = |1\rangle_{I_x} |0\rangle_{I_y}$ , and  $|V\rangle_I = |0\rangle_{I_x} |1\rangle_{I_y}$ . Following measurement postselection, this state is a maximally entangled singlet state of the form in Eqn. 1 [14], and expansions to higher orders account for multiple-pair effects. A useful property of the full state  $|\psi\rangle_{SI}$  is that its anti-normally ordered characteristic function is a zero-mean, jointly Gaussian distribution that remains Gaussian under linear transformations. Its joint density operator is  $\hat{\rho}_{SI} = \hat{\rho}_{S_x I_y} \otimes \hat{\rho}_{S_y I_x}$ , whose anti-normally ordered characteristic functions are given by


$$\begin{aligned}\chi_A^{\rho_{S_x I_y}}(\zeta_S, \zeta_I) &= \langle e^{-\zeta_S^* \hat{a}_{S_x} - \zeta_I^* \hat{a}_{I_y}} e^{\zeta_S \hat{a}_{S_x}^\dagger + \zeta_I \hat{a}_{I_y}^\dagger} \rangle \\ &= e^{-(1+\bar{N})(|\zeta_S|^2 + |\zeta_I|^2) + 2\bar{N}\text{Re}(\zeta_S \zeta_I)}\end{aligned}\quad (13)$$

and

$$\begin{aligned}\chi_A^{\rho_{S_y I_x}}(\zeta_S, \zeta_I) &= \langle e^{-\zeta_S^* \hat{a}_{S_y} - \zeta_I^* \hat{a}_{I_x}} e^{\zeta_S \hat{a}_{S_y}^\dagger + \zeta_I \hat{a}_{I_x}^\dagger} \rangle \\ &= e^{-(1+\bar{N})(|\zeta_S|^2 + |\zeta_I|^2) - 2\bar{N}\text{Re}(\zeta_S \zeta_I)},\end{aligned}\quad (14)$$


which contain all multiple-pair orders of  $|\psi\rangle_{SI}$ . Following a linear transformation, the output state  $\hat{\rho}_{out}$  can be determined by taking the inverse Fourier transform of the output characteristic function, with that characteristic function being easily calculated from the input characteristic function and the field transformation. In a memory or teleportation architecture, we want the output—represented by a pure or mixed state  $\hat{\rho}_{out}$ —to have the highest possible fidelity with respect to its input state  $\hat{\rho}_{in}$ . The trace separation quantifies this fidelity as  $F(\hat{\rho}) = \text{Tr} [\sqrt{\hat{\rho}_{out}} \hat{\rho}_{in} \sqrt{\hat{\rho}_{out}}]$ , which reduces to a projection overlap  $\langle \psi_{in} | \hat{\rho}_{out} | \psi_{in} \rangle$  when the input is the pure state  $\hat{\rho}_{in} = |\psi_{in}\rangle \langle \psi_{in}|$ .

The most immediate problem affecting the distribution of polarization-entangled photons is photon propagation loss. Although a 1 km length of low-loss, optical telecom fiber ( $\lambda = 1.55 \mu m$ )—a relatively short distance—has a transmission nearing 95%, photon transmission decays exponentially with increasing distances, ruling out direct transmission of entangled photons over hundreds of kilometers. In classical communication, this type of signal attenuation is easily compensated with fiber-based amplifiers. However, for quantum communication, the no-cloning theorem prevents noiseless amplification of the

 Fig. 10(a) shows a 'repeat-and-measure' approach that overcomes the amplifier restriction by incrementally extending entanglement across a larger network [15]. Lettered boxes represent memories, and pairs of memories form nodes on the network. For two nodes, A–B and C–D, that are each separately entangled, a joint Bell measurement between memories B and C will entangle systems A and D. This process, known as *entanglement swapping*, establishes entanglement between two networks links that may have never interacted. One can then establish entanglement between memories A and Z that are separated by a distance  $L$  by independently creating entanglement at  $N$  equally-spaced adjacent nodes out of  $2N$  memories.  $N - 1$  entanglement swapping operations in this network ultimately entangle memories A and Z. For the repeater protocol to work in an asynchronous fashion, these links must be heralding quantum memories.


Once entanglement distribution is successful, we can carry out long-distance quantum communication using teleportation, a protocol described in Fig. 4(b). In qubit teleportation, Charlie (C) sends a message (M) to Bob (B) using Alice (A) as an intermediary. Charlie's message is in the form of an arbitrary, unknown qubit spanned by the orthonormal basis  $\{|0\rangle, |1\rangle\}$  represented by  $|\psi\rangle_C = \alpha |0\rangle_C + \beta |1\rangle_C$ , where  $|\alpha|^2 + |\beta|^2 = 1$ . Together, entanglement repeaters and teleportation form basis for high-fidelity communications of quantum states over long distances.

## Entanglement Creation ??, with architectures for polarization entanglement distribution, entanglement connection, and quantum teleportation. We will first discuss our architecture



**QM** entanglement distribution and protocol. A very basic abstraction for quantum memories. The remainder characterizes figures of merit—fidelity, heralding probability, and protocol success probability—under various transmission loss and photodetection conditions.


## First Entanglement Swapping




### II.1. Architecture Overview and Figures of Merit

We will first discuss our overall architecture and loss model, followed by particular details of the dual-OPA polarization entanglement source and quantum memories. Our architecture for polarization entanglement distribution, shown in Fig. 5, is a modification of the standard DLCZ architecture shown in Fig. 2. In the original DLCZ protocol, both ensembles are coherently pumped and the probability that both ensembles will emit single photons is low compared to that for emission from a single ensemble. Interference at the 50-50 beam splitter in Fig. 2 erases any which-path information for the emission event, so single detection at either photon counter  $D_1$  or  $D_2$  is used to herald entanglement of the two ensembles. The figure caption [11] in Fig. 5 is when a polarization singlet is emitted from the source and the overall system is lossless. The polarizing beam splitters (PBS) then load the signal and idler photons from the singlet into a coherent superposition of excitations of ensembles 1 and 2 and ensembles 3 and 4, respectively. This loading is heralded by the single-photon detections from pair  $(D_1, D_2)$  and  $(D_3, D_4)$ .

## Last Entanglement Swapping



(a) Principle of a quantum repeater architecture. Entanglement is independently created at short distances between nodes AB...YZ. Entanglement is swapped between neighboring links such that locations A and D, for example, share entanglement over an intermediate distance. Swapping occurs over successive long distances of the two ensembles. The separation, A and Z, are entangled.



(b) Optical qubit teleportation. Two nodes sharing entanglement can teleport quantum data over intermediate nodes via a classical communication channel. Figure 6 shows the error modes for the Fig. 5 distribution architecture. The Type-II SPDC source may produce multiple signal-idler pairs, which will be modeled with the full joint Gaussian state description of its output. Propagation losses between the PBS and the atomic ensembles (labelled ‘pre’) and between the atomic ensembles and the 50-50 beam splitter (labelled ‘post’) are modeled by fictitious beam splitters whose vacuum-state input ports inject Gaussian quantum noise. Finite quantum efficiency photodetectors (labelled ‘pho’) are similarly modeled, and we have ignored dark counts, which are known to be alarmingly low at heralding wavelength.

Figures 6 and 7 show silicon Geiger-mode avalanche photodiodes (APDs) [18]. Our number state model for these fictitious beam splitters, described in Section ??, includes the effects of phase differences between input ports, which we can use to characterize the effects of phase mismatch between pairs of ensembles. We will assume that any accumulated phase offsets leading to the ensembles can be incorporated into the pre-transmission efficiency.

The preceding imperfections expose two fundamentally different failure modes for this protocol that affect fidelity and probability of success, respectively. First, is it possible for heralding detections at the APDs to declare the protocol’s success even when the ensembles

FIG. 4. Fundamentals of quantum communication using entanglement connection and teleportation.

## II. HERALDED POLARIZATION ENTANGLEMENT DISTRIBUTION WITH ATOMIC ENSEMBLES

This chapter combines the operating principles of ensemble-based quantum memories, as described in




FIG. 5. Modified DLCZ architecture for distributing polarization entanglement, using a spontaneous parametric downconversion (SPDC) source and interferometer measurement for entanglement verification.




FIG. 6. Hong-Ou-Mandel (HOM) interferometer measurement with loss (labeled).

themselves are not in a polarization singlet state. For example, a multiple-pair event from the entanglement source could lead to multiple heralding photons emitted by a quantum memory. Post-memory attenuation and finite-quantum efficiency photodetection could eliminate all but one of those heralding photons, and the usage of Geiger-mode APDs might completely preclude our ability to distinguish between multiple-photon and single-photon events. A relative phase offset between two ensembles, either because of pump photon phase mismatch or pre-transmission phase accumulation, would similarly affect the final fidelity of the loaded quantum state. Lastly, because post-memory imperfections reduce potential heralding detections, it is possible to declare the protocol a failure—and reduce the probability of success—even when the ensembles are successfully

loaded.

We characterize this architecture's performance of heralding entanglement distribution by determining its fidelity figure-of-merit and measurement statistics under different photodetection schemes: photon-number resolving detection (PNRD), which can distinguish between single-photon and multi-photon photodetection events; and non-resolving single-photon detection (NRPD), which is unable to exclude multi-photon error events. For these two schemes, the projective measurement operators  $\hat{M}_j$  ( $j = 1, 2, 3, 4$ ) represent four successful heralding outcomes at the photodetectors  $D_i$  ( $i = 1, 2, 3, 4$ ). The input to each 50-50 beam splitter in Fig. 5 is a superposition state of linearly co-polarized heralding photons from the atomic memories, so we expect that successful entanglement distribution to yield

single clicks at the signal and idler photodetector pairs. This DLCZ-like protocol for heralded polarization entanglement distribution therefore requires the occurrence of a single detection on either photodetectors  $D_1$  or  $D_2$ , as well as a *corresponding event* on either  $D_4$  or  $D_3$ . In the following, successful heralding is given by the projective measurements

$$\hat{M}_j^{\text{PNRD}} = \begin{cases} (|1\rangle_{11}\langle 1|) \otimes (|0\rangle_{22}\langle 0|) \otimes (|1\rangle_{33}\langle 1|) \otimes (|0\rangle_{44}\langle 0|) & j = 1 \\ (|0\rangle_{11}\langle 0|) \otimes (|1\rangle_{22}\langle 1|) \otimes (|1\rangle_{33}\langle 1|) \otimes (|0\rangle_{44}\langle 0|) & j = 2 \\ (|0\rangle_{11}\langle 0|) \otimes (|1\rangle_{22}\langle 1|) \otimes (|0\rangle_{33}\langle 0|) \otimes (|1\rangle_{44}\langle 1|) \\ (|1\rangle_{11}\langle 1|) \otimes (|0\rangle_{22}\langle 0|) \otimes (|0\rangle_{33}\langle 0|) \otimes (|1\rangle_{44}\langle 1|) \end{cases} \quad (15)$$

$$\hat{M}_j^{\text{NRPD}} = \begin{cases} (\hat{I}_1 - |0\rangle_{11}\langle 0|) \otimes (|0\rangle_{22}\langle 0|) \otimes (\hat{I}_3 - |0\rangle_{33}\langle 0|) \\ (|0\rangle_{11}\langle 0|) \otimes (\hat{I}_2 - |0\rangle_{22}\langle 0|) \otimes (\hat{I}_3 - |0\rangle_{33}\langle 0|) \\ (|0\rangle_{11}\langle 0|) \otimes (\hat{I}_2 - |0\rangle_{22}\langle 0|) \otimes (|0\rangle_{33}\langle 0|) \otimes (\hat{I}_4 - |0\rangle_{44}\langle 0|) \\ (\hat{I}_1 - |0\rangle_{11}\langle 0|) \otimes (|0\rangle_{22}\langle 0|) \otimes (|0\rangle_{33}\langle 0|) \otimes (\hat{I}_4 - |0\rangle_{44}\langle 0|) \end{cases} \quad (16)$$

where  $|n\rangle_i$  ( $i = 0, 1$ ) are the vacuum and single-photon states, and  $\hat{I}_i$  is the identity operator for the  $\hat{a}_i$  mode measured at photodetector  $D_i$  ( $i = 1, 2, 3, 4$ ).

The heralding probability  $P_{\text{herald}}$  is the probability a heralding—a ( $D_1, D_2$ ) single click and a ( $D_3, D_4$ ) single click. The success probability  $P_{\text{success}}$  is the probability that heralding has occurred and the four ensembles have loaded a polarization Bell state. The fidelity  $F_j$  is the projection of the post-heralding ensemble state onto the appropriate Bell state for  $\hat{M}_j$  heralding, i.e.,

$$|\psi_j\rangle = \frac{1}{\sqrt{2}} \left( |1\rangle_S^1 |0\rangle_S^2 |0\rangle_S^3 |1\rangle_S^4 + (-1)^j |0\rangle_S^1 |1\rangle_S^2 |1\rangle_S^3 |0\rangle_S^4 \right) \quad (17)$$

Following a photodetection measurement, the joint density operator of the four atomic ensembles is determined by applying  $\hat{M}_j$  and tracing over the optical modes:


$$\hat{\rho}_{\text{post}}^j = \frac{1}{P_j^{\text{herald}}} \text{tr}_{1,2,3,4} \left( \hat{\rho}_{\text{out}} \hat{M}_j \right) \quad (18)$$

where  $\hat{\rho}_{\text{out}}$  is the joint density operator of the heralding light fields and the ensembles

$$P_j^{\text{herald}} = \text{tr} \left( \hat{\rho}_{\text{out}} \hat{M}_j \right). \quad (19)$$

If  $|\psi_j\rangle$  is the entangled state of the four ensembles as heralded by  $\hat{M}_j$ , and the entanglement storage fidelity is  $F_j$ , we find that the success probability is

$$P_{\text{success}} = \sum_{j=1}^4 P_j^{\text{herald}} F_j = \sum_{j=1}^4 \text{tr} \left( \hat{\rho}_{\text{out}} \hat{M}_j \right) \langle \psi_j | \hat{\rho}_{\text{post}}^j | \psi_j \rangle. \quad (20)$$



**FIG. 3.** Beam splitter action on an input density operator  $\hat{\rho}_{\text{in}}$  with two input ports and two output ports. Labeled at each port are the input and output bases used in Eqns. 24 and 25. The boson annihilation operations at the input and output are  $\hat{a}_i$  and  $\hat{b}_i$ , respectively, and the accompanying index  $i$  represents signal ( $i = 1$ ) and auxiliary modes ( $i = 2$ ).

We will determine these measurement statistics using a full-number-state analysis. Because the successful heralding outcomes defined by  $\hat{M}_j$  are symmetric, all the fidelities  $F_j$  are equal to each other. We will, therefore, calculate only  $F_1$  without any loss of generality. In what immediately follows, we will formalize our approach for calculating the joint density operator of the light-ensemble system following entanglement distribution.

## II.2. Photon Number Probability Distributions

Our first task is to concisely represent the effects of propagation loss on light beams of arbitrary statistical composition, as shown by the unitary beam splitter transformation in Fig. 7. The matrix representation of linear-loss beam splitters in Fig. 6 corresponds to the  $SU(2)$  Lie group representation from angular momentum quantization [49]. We can use this equivalence to concisely relate the input and output field density operators from the loss using a joint photon-number probability distribution.

A two-port beam splitter with quantum efficiency  $\eta$  and input-field phase shifts  $\phi_t$  and  $\phi_r$  is described by the general  $SU(2)$  beam splitter operator,

$$\hat{B}(\eta, \phi_t, \phi_r) = e^{-i(\phi_t - \phi_r)\hat{L}_3} e^{-2i\cos^{-1}(\sqrt{\eta})\hat{L}_2} e^{-i(\phi_t + \phi_r)\hat{L}_3}, \quad (21)$$

where the  $\{\hat{L}_i\}$  are the Schwinger angular momentum operators for a two-dimensional quantum harmonic oscillator:

$$\hat{L}_1 = \frac{1}{2} \left( \hat{a}_1^\dagger \hat{a}_2 + \hat{a}_2^\dagger \hat{a}_1 \right) \quad \hat{L}_2 = \frac{1}{2i} \left( \hat{a}_1^\dagger \hat{a}_2 - \hat{a}_2^\dagger \hat{a}_1 \right) \quad \hat{L}_3 = \frac{1}{2} \left( \hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 \right) \quad (22)$$

Given a joint input state—either a pure state  $|\psi\rangle_{\text{in}}$ , or a pure or mixed state  $\hat{\rho}_{\text{in}}$ —of the signal and auxiliary inputs, the output state of the beam splitter is

$$|\psi_{\text{out}}\rangle = \hat{B}^\dagger(\eta, \phi_t, \phi_r) |\psi_{\text{in}}\rangle, \quad \hat{\rho}_{\text{out}} = \hat{B}^\dagger(\eta, \phi_t, \phi_r) \hat{\rho}_{\text{in}} \hat{B}(\eta, \phi_t, \phi_r) \quad (23)$$

In the number-state representation, as shown in Fig. 7, the beam splitter output of a general joint input state

$$\hat{\rho}_{\text{in}} = \sum_{n_1, n_2=0}^{\infty} \sum_{n'_1, n'_2=0}^{\infty} \rho_{\text{in}}(n_1, n_2; n'_1, n'_2) |n_1, n_2\rangle \langle n'_1, n'_2| \quad (24)$$

is,

$$\hat{\rho}_{\text{out}} = \sum_{N_1, N_2=0}^{\infty} \sum_{N'_1, N'_2=0}^{\infty} \rho_{\text{out}}(N_1, N_2; N'_1, N'_2) |N_1, N_2\rangle \langle N'_1, N'_2|, \quad (25)$$

The output state (Eqn. 25) follows from the input state (Eqn. 24) by first applying the unitary beam splitter transformation in Eqn. 21, and then inserting the identity operator in the  $|N_1, N_2\rangle$  basis. The output matrix elements in Eqn. 25 are

$$\rho_{\text{out}}(N_1, N_2; N'_1, N'_2) = \sum_{n_1, n_2=0}^{\infty} \sum_{n'_1, n'_2=0}^{\infty} B_{N_1, N_2}^{n_1, n_2}(\eta, \phi_t, \phi_r) [B_{N'_1, N'_2}^{n'_1, n'_2}(\eta, \phi_t, \phi_r)]^* \quad (26)$$

where

$$\begin{aligned} B_{N_1, N_2}^{n_1, n_2}(\eta, \phi_t, \phi_r) &= \langle N_1, N_2 | \hat{B}^\dagger(\eta, \phi_t, \phi_r) | n_1, n_2 \rangle \\ &= R_{N_1, N_2}^{n_1, n_2}(\eta) e^{i\phi_t(N_1 - n_2) + i\phi_r(N_1 - n_1)}. \end{aligned} \quad (27)$$

The transformation coefficients,

$$R_{N_1, N_2}^{n_1, n_2}(\eta) = \left| B_{N_1, N_2}^{n_1, n_2}(\eta, \phi_t, \phi_r) \right| = \langle N_1, N_2 | e^{-2i\cos^{-1}(\sqrt{\eta})\hat{L}_2} | n_1, n_2 \rangle \quad (28)$$

can be calculated in terms of the Jacobi polynomials  $P_n^{(\alpha, \beta)}(x)$  as

$$R_{N_1, N_2}^{n_1, n_2}(\eta) = \sqrt{\frac{N_1! N_2!}{n_1! n_2!} \eta^{N_1 - n_2} (1 - \eta)^{N_1 - n_1}} P_{N_2}^{(N_1 - n_1, N_1 - n_2)}(2\eta - 1) \quad (29)$$

where

$$\begin{aligned} P_n^{(\alpha, \beta)}(x) &= \frac{(-1)^n}{2^n n!} (1 - x)^{-\alpha} (1 + x)^{-\beta} \left( \frac{d}{dx} \right)^n [(1 - x)^{n+\alpha}] \\ &\quad \alpha, \beta > -1, \end{aligned} \quad (30)$$

The restriction  $N_1 \geq n_1, n_2$  ensures the orthogonality of the Jacobi polynomials in Eqn. 29 over the range of quantum efficiency  $0 \leq \eta \leq 1$ ; we will extend the range of this coefficient shortly. The  $R$  coefficient characterizes the output photon-number probability distribution for the joint input state  $|n_1, n_2\rangle$ . For a number-diagonal joint input state, it can be shown that

$$P_{\text{out}}(N_1, N_2) = \langle N_1, N_2 | \hat{\rho}_{\text{out}} | N_1, N_2 \rangle = \sum_{\substack{n_1=0 \\ n_2=N_1+N_2-n_1}}^{N_1+N_2} P_{\text{out}}(N_1, N_2; n_1, n_2) \quad (31)$$

where the conditional and joint probabilities are

$$P_{\text{out}}(N_1, N_2 | n_1, n_2) = \left[ R_{N_1, N_2}^{n_1, n_2}(\eta) \right]^2 \quad \text{and} \quad P_{\text{in}}(n_1, n_2) = \rho_{\text{in}}(n_1, n_2) \quad (32)$$

respectively.

The beam splitter's physical symmetries allow us to extend the Jacobi polynomials over the full range photon-number output probabilities:

$$R_{N_1, N_2}^{n_1, n_2}(\eta) = \begin{cases} (-1)^{N_1 - n_1} R_{n_1, n_2}^{N_1, N_2}(\eta) & n_2 \leq N_1 < n_1 \\ R_{n_2, n_1}^{N_2, N_1}(\eta) & n_1 \leq N_1 < n_2 \\ (-1)^{N_1 - n_1} R_{N_2, N_1}^{n_2, n_1}(\eta) & n_1 > N_1, n_2 > N_1 \\ 0 & n_1 + n_2 \neq N_1 + N_2 \end{cases} \quad (33)$$

We have made the unitary constraint explicit, as the Jacobi polynomials are not *ad hoc* restricted to events for which  $n_1 + n_2 = N_1 + N_2$ . Eqn. 26 simplifies because the unitarity requirement  $\rho_{\text{in}}(n_1, n_2; n'_1, n'_2) = \rho_{\text{in}}(n'_1, n'_2; n_1, n_2)$  implies  $n_1 + n_2 = N_1 + N_2$  and  $n'_1 + n'_2 = N'_1 + N'_2$ . We thus eliminate the second and fourth summations in Eqn. 26 because  $n_2 = N_1 + N_2 - n_1$  and  $n'_2 = N'_1 + N'_2 - n'_1$ , and restrict the remaining first and third summation there to  $n_1 \in \{0, 1, \dots, N_1 + N_2\}$  and  $n'_1 \in \{0, 1, \dots, N'_1 + N'_2\}$ , respectively.

### III. ENTANGLEMENT CONNECTION

After successfully distributing entanglement to a pair of nodes, resulting in local entanglement, we will want to extend our quantum communication capabilities over distances prohibited by direct transmission. In this section, we use post-selected, polarization-entangled ensembles to accomplish a basic task in long-distance quantum communication, namely entanglement swapping. The high level concepts underlying this procedure was described in Section ?? of this thesis' introduction, and is a modification of the DLCZ protocol's application to quantum communication [4].

Fig. 8 outlines a procedure for accomplishing polarization entanglement connection with atomic ensembles. Polarization entanglement is generated independently, at two different nodes, as described in Section II.1 and Fig. 4(a). A Bell-state measurement between the  $(S_y^A, S_x^A)$  and  $(S_y^B, S_x^B)$  ensemble pairs establishes polarization entanglement between the remaining idler ensemble pairs  $(I_y^A, I_x^A)$  and  $(I_y^B, I_x^B)$ . Coherent, on-resonance pulses at each of the signal ensembles reads a Dicke excitation out of the  $|s\rangle - |e\rangle$  atomic transition into a well-defined spatial mode. Loss-modeling beam splitters before interference ( $\eta_{\text{pre}}$ ) and at photodetection ( $\eta_{\text{pho}}$ ) characterize the quantum efficiency losses, although to maintain consistency with [20], we calculate fidelity with respect to  $\eta_{\text{meas}} = \eta_{\text{pre}} \eta_{\text{pho}}$ . In the following, we determine the fidelity and probability of success when ensembles A and B are independently in singlet or Gaussian states.

Although a full Bell state measurement is not possible using linear optics [21], the observation of single clicks




FIG. 8. Polarization entanglement connection. We assume that ensembles A and B are independently in polarization singlet states or Gaussian states following entanglement distribution in Section II.1. The anti-Stokes photons from reading the A and B signal ensembles are interfered at the 50-50 beam splitter. Photon detections at  $D_A$  and  $D_B$  heralds entanglement connection between the idler ensembles A and B.

at both  $D_A$  and  $D_B$ , when these are unity quantum efficiency photon-number resolving detectors, uniquely heralds the measurement of a singlet state and successful completion of entanglement connection protocol when the A and B ensembles were both in their singlet states. Two signal ensemble pairs in independent singlet states can be separated into four orthogonal basis states

$$|\psi\rangle^A \otimes |\psi\rangle^B = \frac{1}{\sqrt{2}} \left( |1\rangle_{S_y^A} |0\rangle_{S_x^A} |0\rangle_{I_y^A} |1\rangle_{I_x^A} - |0\rangle_{S_y^A} |1\rangle_{S_x^A} |1\rangle_{I_y^A} |0\rangle_{I_x^A} \right) \otimes \frac{1}{\sqrt{2}} \left( |1\rangle_{S_y^B} |0\rangle_{S_x^B} |0\rangle_{I_y^B} |1\rangle_{I_x^B} - |0\rangle_{S_y^B} |1\rangle_{S_x^B} |1\rangle_{I_y^B} |0\rangle_{I_x^B} \right) \exp \left[ - (1 + \bar{N}) \left( |\zeta_{S_y^A}|^2 + |\zeta_{S_x^A}|^2 + |\zeta_{I_y^A}|^2 + |\zeta_{I_x^A}|^2 \right) - (1 + \bar{N}) \left( |\zeta_{S_y^B}|^2 + |\zeta_{S_x^B}|^2 + |\zeta_{I_y^B}|^2 + |\zeta_{I_x^B}|^2 \right) + 2\text{Re} \left( \tilde{N} \zeta_{S_x^A} \zeta_{I_y^A} \right) - 2\text{Re} \left( \tilde{N} \zeta_{S_y^A} \zeta_{I_x^A} \right) + 2\text{Re} \left( \tilde{N} \zeta_{S_x^B} \zeta_{I_y^B} \right) - 2\text{Re} \left( \tilde{N} \zeta_{S_y^B} \zeta_{I_x^B} \right) \right], \quad (34)$$

where each  $|\phi_{ij}\rangle$  ( $i, j = x, y$ ) labels a joint state with signal-photon polarizations  $i$  and  $j$  in each path prior to interference. Because of photon-twinning at interference, only two of these orthogonal states— $|\phi_{xy}\rangle$  and  $|\phi_{yx}\rangle$ —contribute to the probability of a heralding event: either  $(\hat{x}, \hat{y})$ - or  $(\hat{y}, \hat{x})$ -polarized photon pairs at  $(D_A, D_B)$ , with equal probability. As such, the entanglement connection fidelity will be unity in both the PNRD and the NRPD cases—

$$F_C = \frac{P_{\text{success}}}{P_{\text{herald}}} = \frac{P_{xy} + P_{yx}}{P_{xy} + P_{yx}} = 1 \quad (35)$$

—independent of pre-transmission and photodetection quantum efficiency losses.

Now let us consider performance when ensembles A and B are modeled as being in Gaussian states parameterized by an average spin excitation number  $\bar{N}$ . In the absence of any important nonlinear elements in Fig. 8,

we will perform a characteristic function analysis instead of a number-state analysis. The joint density operator of these ensembles is  $\hat{\rho}_{SI}^{\text{in}} = \hat{\rho}_{SI}^A \otimes \hat{\rho}_{SI}^B$ , where  $\hat{\rho}_{SI}^i = \hat{\rho}_{S_x^i I_y^i} \otimes \hat{\rho}_{S_y^i I_x^i}$  ( $i = A, B$ ), which is represented by the anti-normally ordered characteristic function,

$$\begin{aligned} \chi_A^{\text{in}}(\zeta) = & \langle D_A \left( \hat{a}_{S_y^A}, \zeta_{S_y^A} \right) D_A \left( \hat{a}_{S_x^A}, \zeta_{S_x^A} \right) D_A \left( \hat{S}_{I_y^A}, \zeta_{I_y^A} \right) D_A \left( \hat{S}_{I_x^A}, \zeta_{I_x^A} \right) \\ & \cdot D_A \left( \hat{a}_{S_y^B}, \zeta_{S_y^B} \right) D_A \left( \hat{a}_{S_x^B}, \zeta_{S_x^B} \right) D_A \left( \hat{S}_{I_y^B}, \zeta_{I_y^B} \right) D_A \left( \hat{S}_{I_x^B}, \zeta_{I_x^B} \right) \\ & \exp \left[ - (1 + \bar{N}) \left( |\zeta_{S_y^A}|^2 + |\zeta_{S_x^A}|^2 + |\zeta_{I_y^A}|^2 + |\zeta_{I_x^A}|^2 \right) - (1 + \bar{N}) \left( |\zeta_{S_y^B}|^2 + |\zeta_{S_x^B}|^2 + |\zeta_{I_y^B}|^2 + |\zeta_{I_x^B}|^2 \right) + 2\text{Re} \left( \tilde{N} \zeta_{S_x^A} \zeta_{I_y^A} \right) - 2\text{Re} \left( \tilde{N} \zeta_{S_y^A} \zeta_{I_x^A} \right) + 2\text{Re} \left( \tilde{N} \zeta_{S_x^B} \zeta_{I_y^B} \right) - 2\text{Re} \left( \tilde{N} \zeta_{S_y^B} \zeta_{I_x^B} \right) \right], \end{aligned} \quad (36)$$

where  $\tilde{N} = \sqrt{\bar{N}(\bar{N}+1)}$ ,  $D_A(\hat{a}_i, \zeta_i) = e^{-\zeta_i^* \hat{a}_i} e^{\zeta_i \hat{a}_i^\dagger}$  is the antinormally-ordered displacement operator, and

$$\zeta = [\zeta_S, \zeta_I]^T = \left[ \zeta_{S_y^A}, \zeta_{S_y^B}, \zeta_{S_x^A}, \zeta_{S_x^B}, \zeta_{I_y^A}, \zeta_{I_y^B}, \zeta_{I_x^A}, \zeta_{I_x^B} \right]^T. \quad (37)$$

The optical modes reaching the detectors  $D_i$  in Fig. 6 are

$$\hat{\mathbf{a}}_S^{\text{out}} = \sqrt{\eta^{\text{pho}} \eta^{\text{pre}}} \mathbf{B} \hat{\mathbf{a}}_{\text{in}}^S + \sqrt{\eta^{\text{pho}} (1 - \eta^{\text{pre}})} \mathbf{B} \hat{\mathbf{a}}_v^{\text{pre}} + \sqrt{1 - \eta^{\text{pho}}} \hat{\mathbf{a}}_v^{\text{pho}} \quad (38)$$

where we have defined the operator-valued vectors

$$\begin{aligned}\hat{\mathbf{a}}_S^{\text{out}} &= \left[ \hat{a}'_{S_y^A}, \hat{a}'_{S_y^B}, \hat{a}'_{S_x^A}, \hat{a}'_{S_x^B} \right]^T \\ \hat{\mathbf{a}}_S^{\text{in}} &= \left[ \hat{a}_{S_y^A}, \hat{a}_{S_y^B}, \hat{a}_{S_x^A}, \hat{a}_{S_x^B} \right]^T \\ \hat{\mathbf{a}}^{\text{pre}} &= [\hat{a}_1^{\text{pre}}, \dots, \hat{a}_4^{\text{pre}}]^T \\ \hat{\mathbf{a}}^{\text{pho}} &= \left[ \hat{a}_1^{\text{pho}}, \dots, \hat{a}_4^{\text{pho}} \right]^T\end{aligned}\quad (39)$$

with the linear transformation of the signal modes:

$$\mathbf{B} = \begin{bmatrix} \mathbf{M}_{2 \times 2} & \mathbf{0}_{2 \times 2} \\ \mathbf{0}_{2 \times 2} & \mathbf{M}_{2 \times 2} \end{bmatrix} \quad \mathbf{M}_{2 \times 2} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}. \quad (40)$$

All of the idler modes  $\hat{\mathbf{S}}_I^{\text{in}} = \left[ \hat{S}_{I_y^A}, \hat{S}_{I_y^B}, \hat{S}_{I_x^A}, \hat{S}_{I_x^B} \right]^T$  remain unchanged. The Gaussian mixed-state of the Stokes light arriving at the detectors and the idler ensemble excitations is given by the antinormally-ordered characteristic function

$$\begin{aligned}\chi_A^{\rho_{\text{out}}}(\zeta, \tilde{\zeta}) &= \langle D_A(\hat{\mathbf{a}}_S^{\text{out}}, \zeta_S) D_A(\hat{\mathbf{S}}_I^{\text{in}}, \zeta_I) \rangle \\ &= \chi_A^{\rho_{\text{in}}} \left( \left[ \sqrt{\eta_{\text{meas}}} \tilde{\zeta}_{S_y^A}, \sqrt{\eta_{\text{meas}}} \tilde{\zeta}_{S_y^B}, \sqrt{\eta_{\text{meas}}} \tilde{\zeta}_{S_x^A}, \sqrt{\eta_{\text{meas}}} \tilde{\zeta}_{S_x^B}, \zeta_{I_y^A}, \zeta_{I_y^B}, \zeta_{I_x^A}, \zeta_{I_x^B} \right]^T \right) \\ &\cdot \exp \left[ - \sum_{\{\tilde{\zeta}_i\}} \eta^{\text{pho}} (1 - \eta^{\text{pre}}) |\tilde{\zeta}_i|^2 - \sum_{\{\zeta_i\}} (1 - \eta^{\text{pho}})^2 |\zeta_i|^2 \right]\end{aligned}\quad (41)$$

where the scaled  $\tilde{\zeta}$  result from the transformation of the beam splitter transformation in Eqn. 40:

$$\tilde{\zeta} = \mathbf{B}^\dagger \zeta = \begin{bmatrix} \tilde{\zeta}_{S_y^A} \\ \tilde{\zeta}_{S_y^B} \\ \tilde{\zeta}_{S_x^A} \\ \tilde{\zeta}_{S_x^B} \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} \zeta_{S_y^A} + \zeta_{S_y^B} \\ \zeta_{S_y^A} - \zeta_{S_y^B} \\ \zeta_{S_x^A} + \zeta_{S_x^B} \\ \zeta_{S_x^A} - \zeta_{S_x^B} \end{bmatrix}. \quad (42)$$

Rewriting Eqn. 41 in terms of  $\zeta$ , the characteristic function is now

$$\begin{aligned}\chi_A^{\rho_{\text{out}}}(\zeta) &= \exp \left[ - (1 + \eta_{\text{meas}} \bar{N}) \left( |\zeta_{S_y^A}|^2 + |\zeta_{S_x^A}|^2 + |\zeta_{S_y^B}|^2 + |\zeta_{S_x^B}|^2 \right) \right. \\ &\quad \left. - (1 + \bar{N}) \left( |\zeta_{I_y^A}|^2 + |\zeta_{I_x^A}|^2 + |\zeta_{I_y^B}|^2 + |\zeta_{I_x^B}|^2 \right) \right. \\ &\quad \left. + \tilde{N} \sqrt{2\eta_{\text{meas}}} \left[ \text{Re} \left( \zeta_{S_x^A} \zeta_{I_y^A} \right) + \text{Re} \left( \zeta_{S_x^B} \zeta_{I_y^A} \right) \right. \right. \\ &\quad \left. \left. + \text{Re} \left( \zeta_{S_x^A} \zeta_{I_y^B} \right) - \text{Re} \left( \zeta_{S_x^B} \zeta_{I_y^B} \right) - \text{Re} \left( \zeta_{S_y^A} \zeta_{I_x^A}^{\text{post}} \right) + \text{Re} \left( \zeta_{S_y^B} \zeta_{I_x^A}^{\text{post}} \right) \right] \right. \\ &\quad \left. + \text{Re} \left( \zeta_{S_x^B} \zeta_{I_x^B}^{\text{post}} \right) - \text{Re} \left( \zeta_{S_y^B} \zeta_{I_x^B}^{\text{post}} \right) \right]\end{aligned}\quad (43)$$

A useful property of Gaussian antinormally-ordered characteristic functions is that they can be renormalized into a probability density function, whose moments can be calculated. We find the heralding probability and fidelity by re-expressing Eqn. 43 as

$$\chi_A^{\rho_{\text{out}}}(\zeta) = \frac{\pi^8 p_{\mathbf{Z}}(\zeta)}{D_1} \quad (44)$$

where  $p_{\mathbf{Z}}(\zeta)$  is the probability density function for a zero-mean Gaussian random vector  $\zeta$  with covariance matrices

$$\langle \zeta \zeta^\dagger \rangle = \frac{1}{D_1} \begin{bmatrix} (1 + \eta_{\text{meas}} \bar{N}) \mathbf{I}_{4 \times 4} & \mathbf{0}_{4 \times 4} \\ \mathbf{0}_{4 \times 4} & (1 + \bar{N}) \mathbf{I}_{4 \times 4} \end{bmatrix} \quad \langle \zeta \zeta^T \rangle = \frac{1}{D_1} \begin{bmatrix} \mathbf{0}_{4 \times 4} & \mathbf{0}_{4 \times 4} \\ \mathbf{0}_{4 \times 4} & \mathbf{0}_{4 \times 4} \end{bmatrix} \quad (45)$$

for

$$\mathbf{N}_{4 \times 4} = \tilde{N} \sqrt{2\eta_{\text{meas}}} \begin{bmatrix} 0 & 0 & -1 & -1 \\ 0 & 0 & -1 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 \end{bmatrix} \quad (46)$$

and determinant  $D_1 = (1 + \eta_{\text{meas}} \bar{N})^4 (1 + \bar{N})^4 - 256 \bar{N}^8 \eta_{\text{meas}}^4$ . The output density operator of the Stokes field can be expressed as the operator-valued Fourier transformation of Eqn. 43:

$$\begin{aligned}\hat{\rho}_{\text{out}} &= \int \prod_{i=x,y} \frac{d^2 \zeta_{I_i^A}}{\pi^2} \frac{d^2 \zeta_{I_i^B}}{\pi^2} D_N \left( \hat{S}_{I_i^A}, \zeta_{I_i^A} \right) D_N \left( \hat{S}_{I_i^B}, \zeta_{I_i^B} \right) \\ &\cdot \int \prod_{i=x,y} \frac{d^2 \zeta_{S_i^A}}{\pi^2} \frac{d^2 \zeta_{S_i^B}}{\pi^2} \chi_A^{\rho_{\text{out}}}(\zeta) D_N \left( \hat{a}'_{S_i^A}, \zeta_{S_i^A} \right) D_N \left( \hat{a}'_{S_i^B}, \zeta_{S_i^B} \right)\end{aligned}\quad (47)$$

where  $D_N(\hat{a}_i, \zeta_i) = e^{-\zeta_i \hat{a}_i^\dagger e^{\hat{a}_i^\dagger \hat{a}_i}}$  is the normally-ordered displacement operator. To perform trace operations on the operator-valued Fourier transform in Eqn. 47, we know that

$$\langle 0 | D_N(\hat{a}_i, \zeta_i) | 0 \rangle = 1 \quad \langle 1 | D_N(\hat{a}_i, \zeta_i) | 1 \rangle = 1 - |\zeta_i|^2 \quad (48)$$

and that

$$\text{tr} [D_N(\hat{a}_i, \zeta_i)] = \pi \delta(\zeta_i) \quad \text{tr} [D_N(\hat{a}_i, \zeta_i) (\hat{I} - |0\rangle\langle 0|)] = \pi \delta(\zeta_i) - 1 \quad (49)$$

Performing trace operations on the  $\hat{a}_i$  mode of an output density operator is equivalent to setting  $\zeta_i = 0$  in its characteristic function, so any Gaussian moment calculations in the following may involve a marginal distribution of that specified by Eqn. 45.

We will first consider the particular case of a **y**-polarized click at  $D_A$  and a **x**-polarized click at  $D_B$  and then extrapolate to other possibilities: a **x**-polarized click at  $D_A$  and a **y**-polarized click at  $D_B$ , and co-polarized clicks at both  $D_A$  and  $D_B$ . Applying these trace identities in a PNRD scheme, the post-measurement state  $\rho_{\text{post}}^{\text{out}}$  is

$$\begin{aligned}&+ \text{Re} \left( \frac{\gamma}{P_{\text{herald}}^{\text{yx}}} \int_{S_y^A}^{S_x^B} \int_{I_y^A}^{I_x^B} \prod_{i=x,y} \frac{d^2 \zeta_{I_i^A}}{\pi^2} \frac{d^2 \zeta_{I_i^B}}{\pi^2} D_N \left( \hat{S}_{I_i^A}, \zeta_{I_i^A} \right) D_N \left( \hat{S}_{I_i^B}, \zeta_{I_i^B} \right) \right. \\ &\cdot \left. \int \frac{d^2 \zeta_{S_y^A}}{\pi^2} \frac{d^2 \zeta_{S_x^B}}{\pi^2} \chi_A^{\rho_{\text{out}}}(\zeta) \left( 1 - |\zeta_{S_y^A}|^2 \right) \left( 1 - |\zeta_{S_x^B}|^2 \right) \right),\end{aligned}\quad (50)$$

where the heralding probability is

$$P_{\text{herald}}^{\text{yx}} = \int \frac{d^2 \zeta_{S_y^A}}{\pi^2} \frac{d^2 \zeta_{S_x^B}}{\pi^2} \chi_A^{\rho_{\text{out}}} \left( \left[ \zeta_{S_y^A}, \zeta_{S_y^B}, \zeta_{S_x^A}, \zeta_{S_x^B}, 0, 0, 0, 0 \right]^T \right) \left( 1 - |\zeta_{S_y^A}|^2 \right) \left( 1 - |\zeta_{S_x^B}|^2 \right) \quad (51)$$




FIG. 9. Figures of merit for Gaussian state entanglement connection (PNRD and NRPD) for  $N = 0.01\text{--}0.05$  ( $\Delta N = 0.01$ ).

Applying the Gaussian moment factoring theorem, this

heralding probability is

$$\begin{aligned} P_{\text{herald}}^{yx} &= \frac{1}{D_2} \langle (1 - |\zeta_{S_y^A}|^2) (1 - |\zeta_{S_x^B}|^2) \rangle \\ &= \frac{1}{D_2^3} [D_2^2 - 2(1 + \eta_{\text{meas}} \bar{N}) D_2 + (1 + \eta_{\text{meas}} \bar{N})^2] \\ &= \frac{\eta_{\text{meas}}^2 \bar{N}^2 (\eta_{\text{meas}} \bar{N}(\eta_{\text{meas}} \bar{N} + 3) + 3)^2}{(\eta_{\text{meas}} \bar{N} + 1)^{10}}, \end{aligned} \quad (52)$$

where  $\langle \rangle$  denotes ensemble averaging treating  $\zeta$  as a complex-valued Gaussian random vector whose probability density function is a marginal distribution of Eqn. 45, with  $D_2 = (1 + \eta_{\text{meas}} \bar{N})^4$ . Note that the second-moments of the signal modes reaching the photodetectors are all identical, so the heralding probabilities for each of the four heralding probabilities mentioned earlier are equal. Therefore,  $P_{\text{herald}} = 4P_{\text{herald}}^{yx}$  for the PNRD case. Applying the trace identities in Eqn 49, the post-measurement joint density operator for the NRPD scheme is

$$\begin{aligned} \hat{\rho}_{\text{post}}^{yx} &= \frac{1}{P_{\text{herald}}^{yx}} \int \prod_{i=x,y} \frac{d^2 \zeta_{I_i^A}}{\pi^2} \frac{d^2 \zeta_{I_i^B}}{\pi^2} D_N \left( \hat{S}_{I_i^A}, \zeta_{I_i^A} \right) D_N \left( \hat{S}_{I_i^B}, \zeta_{I_i^B} \right) \\ &\cdot \int \frac{d^2 \zeta_{S_y^A}}{\pi^2} \frac{d^2 \zeta_{S_x^B}}{\pi^2} \chi_A^{\rho_{\text{out}}}(\zeta) [\pi \delta(\zeta_{S_y^A}) - 1] [\pi \delta(\zeta_{S_x^B}) - 1]. \end{aligned} \quad (53)$$

Tracing out the idler excitation modes, the NRPD heralding probability is

$$\begin{aligned} P_{\text{herald}}^{yx} &= \\ &\int \prod_{i=x,y} \frac{d^2 \zeta_{S_i^A}}{\pi^2} \frac{d^2 \zeta_{S_i^B}}{\pi^2} \chi_A^{\rho_{\text{out}}} \left( \left[ \zeta_{S_y^A}, \zeta_{S_y^B}, \zeta_{S_x^A}, \zeta_{S_x^B}, 0, 0, 0, 0 \right]^T \right) [\pi \delta(\zeta_{S_y^A}) - 1] [\pi \delta(\zeta_{S_x^B}) - 1]. \end{aligned} \quad (54)$$

which yields

$$P_{\text{herald}}^{yx} = \frac{\eta_{\text{meas}}^2 \bar{N}^2}{(\eta_{\text{meas}} \bar{N} + 1)^4}. \quad (55)$$

This PNRD heralding probability and PNRD fidelity, as well as the NRPD heralding probability and fidelity are shown in Fig. 9. The results for heralding probability are consistent with our understanding of multiple-excitation effects, as described previously in the context of entanglement distribution in Section ??: reading from ensembles with a higher average spin excitation  $N$  will yield more anti-Stokes photons, resulting in a higher likelihood of a heralding event when the photodetectors cannot resolve photon number.

The fidelity, when a **y**-polarized  $D_A$  click and a **x**-

polarized  $D_B$  click provide the herald, is given by

$$\begin{aligned}
F_{yx} &= \langle \psi_1 | \hat{\rho}_{\text{post}}^{yx} | \psi_1 \rangle \\
&= \frac{1}{P_{\text{herald}}^{yx}} \int \prod_{i=x,y} \frac{d^2 \zeta_{I_i^A}}{\pi^2} \frac{d^2 \zeta_{I_i^B}}{\pi^2} \left[ 1 - \frac{|\zeta_{I_x^A} \zeta_{I_y^B} - \zeta_{I_y^A} \zeta_{I_x^B}|^2}{2} \right] \\
&\quad \cdot \int \frac{d^2 \zeta_{S_y^A}}{\pi^2} \frac{d^2 \zeta_{S_x^B}}{\pi^2} \chi_A^{\rho_{\text{out}}}(\zeta) \left( 1 - |\zeta_{S_y^A}|^2 \right) \left( 1 - |\zeta_{S_x^B}|^2 \right) \\
&= \frac{1}{D_1 P_{\text{herald}}^{yx}} \langle \left( 1 - |\zeta_{S_y^A}|^2 \right) \left( 1 - |\zeta_{S_x^B}|^2 \right) \left( 1 - \frac{|\zeta_{I_x^A} \zeta_{I_y^B} - \zeta_{I_y^A} \zeta_{I_x^B}|^2}{2} \right) \rangle \quad (56)
\end{aligned}$$

The necessary higher-order moments for this calculation are, from Gaussian moment factoring

$$\begin{aligned}
\langle |\zeta_{I_x^A} \zeta_{I_y^B} - \zeta_{I_y^A} \zeta_{I_x^B}|^2 \rangle &= \langle |\zeta_{I_x^A}|^2 \rangle \langle |\zeta_{I_y^B}|^2 \rangle + \langle |\zeta_{I_y^A}|^2 \rangle \\
\langle |\zeta_{S_y^A}|^2 |\zeta_{I_x^A} \zeta_{I_y^B} - \zeta_{I_y^A} \zeta_{I_x^B}|^2 \rangle &= \langle |\zeta_{S_y^A}|^2 \rangle \langle |\zeta_{I_x^A}|^2 \rangle \langle |\zeta_{I_y^B}|^2 \rangle \\
&\quad + \langle |\zeta_{S_y^A} \zeta_{I_x^A}|^2 \rangle \langle |\zeta_{I_y^B}|^2 \rangle \\
\langle |\zeta_{S_x^B}|^2 |\zeta_{I_x^A} \zeta_{I_y^B} - \zeta_{I_y^A} \zeta_{I_x^B}|^2 \rangle &= \langle |\zeta_{S_x^B}|^2 \rangle \langle |\zeta_{I_x^A}|^2 \rangle \langle |\zeta_{I_y^B}|^2 \rangle \\
&\quad + \langle |\zeta_{S_x^B} \zeta_{I_y^B}|^2 \rangle \langle |\zeta_{I_x^A}|^2 \rangle \\
\langle |\zeta_{S_y^A}|^2 |\zeta_{S_x^B}|^2 |\zeta_{I_x^A} \zeta_{I_y^B} - \zeta_{I_y^A} \zeta_{I_x^B}|^2 \rangle &= \langle |\zeta_{S_y^A}|^2 \rangle \langle |\zeta_{S_x^B}|^2 \rangle \langle |\zeta_{I_x^A}|^2 \rangle \\
&\quad + \langle |\zeta_{S_y^A}|^2 \rangle \langle |\zeta_{S_x^B}|^2 \rangle \langle |\zeta_{I_y^A}|^2 \rangle \\
&\quad + \langle |\zeta_{S_y^A} \zeta_{I_x^A}|^2 \rangle \langle |\zeta_{S_x^B} \zeta_{I_y^B}|^2 \rangle \\
&\quad + \langle |\zeta_{S_y^B} \zeta_{I_x^A}|^2 \rangle \langle |\zeta_{S_x^B}|^2 \rangle \langle |\zeta_{I_y^B}|^2 \rangle \\
&\quad + \langle |\zeta_{S_x^B} \zeta_{I_y^B}|^2 \rangle \langle |\zeta_{I_x^A}|^2 \rangle \langle |\zeta_{S_y^B}|^2 \rangle \\
&\quad - \left[ \langle \zeta_{S_y^A} \zeta_{I_x^A} \rangle \langle \zeta_{S_y^A}^* \zeta_{I_x^B}^* \rangle \langle \zeta_{S_x^B} \right. \\
&\quad \left. + \langle \zeta_{S_y^A} \zeta_{I_x^B} \rangle \langle \zeta_{S_y^A}^* \zeta_{I_x^B}^* \rangle \right] \tag{57}
\end{aligned}$$

Applying the covariance matrix in Eqn. 45 to these higher-order moment expressions, the PNRD fidelity is given by

$$F_{yx} = \left[ \frac{1}{D_1^3} \left[ D_1^2 - 2(1 + \eta_{\text{meas}} \bar{N}) 1_2 + (1 + \eta_{\text{meas}} \bar{N})^2 \right] \right. \\ \left. - \frac{1}{2D_1} \left( \frac{2(1 + \bar{N})^2}{D_1^2} - \frac{4(1 + \eta_{\text{meas}} \bar{N})(1 + \bar{N})^2}{D_1^3} \right) \right. \\ \left. + \frac{2(1 + \eta_{\text{meas}} \bar{N})^2 (1 + \bar{N})^2 + 8\eta \tilde{N}^2 (1 + \eta_{\text{meas}} \bar{N})^2}{D_1^4} \right] \quad (58)$$

The  $F_{xy}$  fidelity is easily seen to be the same as the  $F_{yx}$  fidelity, as in the heralding probability calculation presented earlier. While the  $F_{xx}$  and  $F_{yy}$  fidelities will also equal, they will differ from the cross-polarized fidelities.

To find  $F_{yy}$ , we need to evaluate

$$F_{yy} = \frac{1}{D_1 P_{\text{herald}}^{yy}} \left\langle \left(1 - |\zeta_{S_y^A}|^2\right) \left(1 - |\zeta_{S_y^B}|^2\right) \cdot \left(1 - \frac{|\zeta_{I_x^A} \zeta_{I_y^B} - \zeta_{I_y^A} \zeta_{I_x^B}|^2}{2}\right) \right\rangle. \quad (59)$$

The calculation is identical to that for  $F_{yx}$  save for the eighth-order moment, which factors into

$$\begin{aligned}
& |\zeta_{S_y^B}|^2 \left( |\zeta_{I_x^A}|^2 |\zeta_{I_y^B}|^2 - |\zeta_{I_y^A}|^2 |\zeta_{I_x^B}|^2 \right) = \\
& + |\zeta_{S_y^B} \zeta_{I_x^A}|^2 (|\zeta_{S_y^A}|^2 |\zeta_{I_y^B}|^2 + |\zeta_{S_y^B} \zeta_{I_x^A}|^2 |\zeta_{S_y^B}|^2) \\
& + |\zeta_{S_y^A} \zeta_{I_x^B}|^2 (|\zeta_{I_y^A}|^2 |\zeta_{S_y^B}|^2 + |\zeta_{S_y^B} \zeta_{I_x^B}|^2 |\zeta_{S_y^A}|^2). \tag{60}
\end{aligned}$$

Comparing Eqn. 60 and 57, it immediately becomes obvious that the  $F_{yy}$  and  $F_{yx}$  differ in their contribution to the overall entanglement fidelity. Prior to our Gaussian state analysis, we assumed that ensembles A and B had loaded pure singlet states prior to entanglement swapping. Their pure singlet states yield a perfect fidelity in both NRPD and PNRD photodetection schemes with the heralding of cross-polarized photons at detectors  $D_A$  and  $D_B$ . However, the measurement of co-polarized photons is indistinguishable from a cross-polarized photon measurement, even though entanglement need not be successfully swapped when the herald comes from co-polarized photons.<sup>12</sup> As such, we define the fidelity of entanglement as

Figs. 9(b) and 9(c)] show the fidelity of entanglement following a swapping operation. As in polarization entanglement distribution, reading and producing multiple-excitations increases heralding probability at the expense of the entanglement fidelity. In this regard, the fidelity plots for emphasize some significant differences between the PNRD and NRPD cases. From the post-measurement density operator, this NRPD fidelity is given by

$$F_{yx} = \langle \psi_1 | \hat{\rho}_{\text{post}}^{yx} | \psi_1 \rangle$$

$$\frac{8\eta(1+\bar{N})\bar{N}^2}{P_{\text{herald}}^{yx}} \int \prod_{i=x,y} \frac{d^2\zeta_{I_i^A}}{\pi^2} \frac{d^2\zeta_{I_i^B}}{\pi^2} \left[ 1 - \frac{|\zeta_{I_x^A}\zeta_{I_y^B} - \zeta_{I_y^A}\zeta_{I_x^B}|^2}{2} \right]$$

$$\frac{(1+\bar{N})}{\left(\frac{d^2\zeta_{S_y^A}}{\pi^2} \frac{d^2\zeta_{S_x^B}}{\pi^2} \chi_A^{\rho_{\text{out}}}(\zeta)\right)} \left[ \pi\delta\left(\zeta_{S_y^A}\right) - 1 \right] \left[ \pi\delta\left(\zeta_{S_x^B}\right) - 1 \right]. \quad (62)$$

Unlike the difference in moment factoring present in the PNRD fidelity calculation, the NRPD fidelity is identical for cross-polarized and co-polarized photodetection events, as is evident from the NRPD POVM applied in Eqn. 62. This calculation is only dependent on

the second-moments of the idler ensembles and the normalizations inherent in the marginal probability distributions following photodetection, each of which is independent of photon polarization. Equation 62 thus simplifies to

$$F_{yx} = \frac{1}{P_{\text{herald}}^{yx}} \left[ \frac{(\bar{N} + 1)^6 (\eta_{\text{meas}} \bar{N} + 1)^7 - 2(\bar{N} + 1)^6 (\eta_{\text{meas}} \bar{N})^8}{(\bar{N} + 1)^{10} (\eta_{\text{meas}} \bar{N} + 1)^9} \right] \quad (63)$$

which is plotted in Fig. 9(b). The PNRD case lets us actively discard garbage photodetection events that might result from multiple-excitations present at entanglement distribution. In this case, we know that having matching, *single* counts at  $D_A$  and  $D_B$  will very likely correspond to a successful entanglement swapping operation with photon-number resolving detectors. On the other hand, high  $\bar{N}$ , even in the presence of high measurement efficiency, severely affects our NRPD entanglement fidelity, because NRPD cannot distinguish false heralding events. Also surprising is the ratio of the fidelity of entanglement for cross-polarized and co-polarized photodetection events, as shown in Fig. 9(c). These fidelities are very close to each other, although they begin to diverge as  $\bar{N}$  increases. That they are so close in value is somewhat surprising because cross-polarized photodetection was predicted for unity entanglement fidelity in the case of a pure singlet distribution, whereas co-polarized photodetection would not necessarily be indicative of a singlet state remaining in the idler ensembles.

#### IV. SUMMARY AND OUTLOOK

Two areas in this thesis are particularly ripe for extension and exploration: further investigation into the mechanisms and imperfections of ensemble memories that distribute polarization entanglement, and formalizing the number state analysis of systems with loss-modeling beam splitters. Several possible extensions are relatively straightforward, such as the inclusion of phase offsets between orthogonally-polarized paths in during entanglement distribution. Others require a more careful consideration of the underlying physics of quantum memories. For example, the performance analysis of polarization entanglement distribution presented in this thesis omits the presence of spin decoherence in the atomic ensembles. In quantifying the singlet-storage fidelity, there is a tradeoff between the time scale of decoherence of Dicke excitations, the time it takes for a single heralding photon to reach a photodetector, and the post-memory transmission efficiency (which scales with the post-transmission

length). Spin decoherence of the ensemble may therefore be a significant factor limiting the optimal physical distance that local entanglement distribution can cover. Another possible departure would be to discard the DLCZ approach entirely, using a stimulated Raman or EIT approach, instead of a spontaneous Raman process. The ~~Hamiltonian in this approach~~<sup>2</sup> would be amenable to a traditional Gaussian state analysis, and would allow for deterministic control of read and write processes in entanglement distribution. As of the writing of this thesis, however, there is currently no accepted means of non-destructive verification of successful entanglement distribution in a coherently-controlled  $\Lambda$ -level atomic ensemble. This open problem makes certain long-distance quantum communication tasks, such as repeated entanglement connection, difficult to accomplish using a driven Raman process.

Unrelated to the question of memories is formalizing the architectural analysis of loss-modeling beam splitters. The SU(2) number-state representation for *single* beam splitters is discussed in great detail by [19]. Summation compositions of beam splitter coefficients determined important architectural figures of merit—heralding probability, fidelity, and success probability—in the course of this work, but were calculated numerically. With a number-state basis, what general properties or figures of merit of a quantum network can we determine if we limit ourselves to linear optical elements (e.g., 50-50 and polarizing beam splitters, loss elements), photodetectors, and photon-number conserving nonlinear optical elements (e.g., heralding quantum memories or Kerr crystals)? In particular, do any of the properties of a beam splitter coefficient introduced in Chapter II permit any useful, analytical simplifications when using the joint density operator of a network to calculate a figure of merit? Is a number-state analysis consistent with a SPDC Gaussian state analysis when nonlinear elements are excluded and we are limited entirely to the propagation of optical fields? Answers to these questions could possibly alleviate the exhausting notational and computational difficulties currently necessitated by number-state analysis.

#### ACKNOWLEDGMENTS

We thank F. Wong, V. Vuletic, Z. Dutton, S. Johnson, M. Tobenkin, R. Nair, and B. J. Yen for useful discussions and advice. This work was supported by the DARPA Quantum Entanglement Science and Technology (QuEST) program.

[1] H. J. Kimble, *Nature* **453**, 1023 (2008).

[2] S. Lloyd, M. S. Shahriar, J. H. Shapiro, and P. R. Hem-

mer, *Phys. Rev. Lett.* **87**, 167903 (2001).

[3] J. H. Shapiro and N. C. Wong, *Journal of Optics B:*

- Quantum and Semiclassical Optics** **2**, L1 (2000).
- [4] L.-M. Duan, M. D. Lukin, J. I. Cirac, and P. Zoller, **Nature** **414**, 413 (2001).
  - [5] R. H. Dicke, **Phys. Rev.** **93**, 99 (1954).
  - [6] H. Tanji, S. Ghosh, J. Simon, B. Bloom, and V. Vuletić, **Phys. Rev. Lett.** **103**, 043601 (2009).
  - [7] J. Simon, H. Tanji, J. K. Thompson, and V. Vuletić, **Phys. Rev. Lett.** **98**, 183601 (2007).
  - [8] S. M. Barnett and P. M. Radmore, *Methods in Theoretical Quantum Optics* (Oxford Science Publications, 1997).
  - [9] A. B. Klimov and S. M. Chumakov, *A Group-Theoretical Approach to Quantum Optics: Models of Atom-Field Interactions* (Wiley, 2009).
  - [10] M. Fleischhauer and M. D. Lukin, **Phys. Rev. A** **65**, 022314 (2002).
  - [11] M. D. Lukin, **Rev. Mod. Phys.** **75**, 457 (2003).
  - [12] A. V. Gorshkov, A. André, M. Fleischhauer, A. S. Sørensen, and M. D. Lukin, **Phys. Rev. Lett.** **98**, 123601 (2007).
  - [13] A. V. Gorshkov, A. André, M. D. Lukin, and A. S. Sørensen, **Phys. Rev. A** **76**, 033804 (2007).
  - [14] J. H. Shapiro, **New Journal of Physics** **4**, 47 (2002).
  - [15] N. Sangouard, C. Simon, H. de Riedmatten, and N. Gisin, ArXiv e-prints 0906.2699 (2009), arXiv:0906.2699 [quant-ph].
  - [16] J. H. Shapiro, (2008).
  - [17] W. K. Wootters and W. H. Zurek, **Nature** **299**, 802 (1982).
  - [18] O. Thomas, Z. L. Yuan, J. F. Dynes, A. W. Sharpe, and A. J. Shields, **Applied Physics Letters** **97** (2010), cited By (since 1996) 1.
  - [19] R. A. Campos, B. E. Saleh, and M. C. Teich, **Phys. Rev. A** **40**, 1371 (1989).
  - [20] M. Razavi and J. H. Shapiro, **Phys. Rev. A** **73**, 042303 (2006).
  - [21] N. Lütkenhaus, J. Calsamiglia, and K.-A. Suominen, **Phys. Rev. A** **59**, 3295 (1999).