

Simulation of Biological Neural Networks: Neuron Models

Abigail Morrison


January 21st, 2016

Overview

- Review of spatial structure and dynamics of a physiological neuron
- Modelling approaches: detail or abstraction?
- Representation of physiological neuron features in a point neuron model
- Advantages of the point neuron modelling approach
- Simulating a mathematical model
- Choosing a model

The physiological neuron: spatial structure


There are thousands of different neuron types with different structural features



based on drawings by Cajal (1852 -1934)

The physiological neuron: a “typical” cell

However, they all function in much the same way:



1. local input (dendrite)
2. integration (soma)
3. long range conducting (axon)
4. output (synapses)


The physiological neuron: resting potential

K^+/Na^+ pump pushes Na^+ out of the cell and K^+ into it

- This creates a concentration gradient
- a 'force' which acts outwards on K^+

K^+ leaks out, dissipating the gradient

- This causes a net negative charge on the inside of the membrane
- electrical force now acts inwards on K^+



Equilibrium reached when diffusion and electrical forces balanced: the Nernst Potential


$$E_{eq,K^+} = \frac{RT}{F} \ln \frac{[K^+]_o}{[K^+]_i}$$

Resting potential is the weighted average of the equilibrium potentials for each ion

$$E_m = \frac{P_{K^+}}{P_{tot}} E_{eq,K^+} + \frac{P_{Na^+}}{P_{tot}} E_{eq,Na^+} + \frac{P_{Cl^-}}{P_{tot}} E_{eq,Cl^-}$$


The physiological neuron: action potential

Action potential - early phase




sodium influx through
voltage-gated sodium channels

voltage-gated potassium
channels still closed




Action potential - late phase



reduced sodium influx through
voltage-gated sodium channels

potassium efflux through
voltage-gated potassium
channels



Essential features of neurons

- Connectivity / morphology
- Internal dynamics that integrate inputs
- Generation of action potentials

Modelling neurons

If we want to investigate the behaviour of neurons outside of the electrophysiology lab, we need a model that captures these essential features

There are two main approaches:

- Detailed (biophysical) neuron models
- Reduced (abstract) neuron models


Here we focus on the latter approach, specifically on point neuron models.

Detailed neuron models



de Schutter & Bower (1994)


Gerstner & Kistler (2001)



- reconstruct neuron morphology from images
- decompose into many compartments
- define properties of each compartment
- simulate using e.g. NEURON or GENESIS


Point neuron models: spatial structure

Morphology of the real neuron is reduced to a directed graph




Point neuron models: spatial structure

Morphology of the real neuron is reduced to a directed graph



Point neuron models: spatial structure

Morphology of the real neuron is reduced to a directed graph



Point neuron models: dynamics

- One isopotential compartment
- Reduction of dynamics to a few differential equations
- Some examples:

Integrate-and-fire (Lapicque, 1907)

$$C \frac{dU}{dt} = I$$

Hodgkin-Huxley (1952)

$$C \frac{dU}{dt} = - \sum_i I_i(t, U)$$

FitzHugh-Nagumo (1962)

$$\begin{aligned} \frac{dU}{dt} &= U - U^3 - w + I \\ \tau \frac{dw}{dt} &= U - a - bw \end{aligned}$$


Point neuron models: dynamics

The leaky integrate-and-fire model:

- Kirchhoff's current law:

$$C \frac{dU}{dt} + \frac{U}{R} = I$$

$$C \frac{dU}{dt} = -\frac{U}{R} + I$$



- Threshold behaviour: $U > \Theta \quad ? \quad U \leftarrow 0$
- Additional simplifications:
 - linear subthreshold integration
 - time invariant parameters

Comparison of detailed and point neuron models

Detailed neuron models:

- have physical extent
- give a good approximation of the electrical properties of a physiological neuron
- can be made arbitrarily complex

Point neuron models:

- have no physical extent
- represent the dynamics in an extremely reduced fashion
- are very simple

Comparison of detailed and point neuron models

Detailed neuron models:

- have physical extent
- give a good approximation of the electrical properties of a physiological neuron
- can be made arbitrarily complex

Point neuron models:

- have no physical extent
- represent the dynamics in an extremely reduced fashion
- are very simple

Why would anyone want to use a point neuron model?

Why use point neuron models?


- Much less initial investment
- Analysis
- Network simulation
- Not as bad an approximation as one might think

Why use point neuron models?

- Much less initial investment
- Analysis
- Network simulation
- Not as bad an approximation as one might think

Point neurons: not as bad as all that

Point neurons can exhibit a wide range of dynamics:



Kobayashi et al. (2009);
see also Izhikevich (2004)

Point neurons: not as bad as all that

SUBMISSION

This contest is currently not open for submission.

SUB NAVIGATION

DIADEM

Spike Time Prediction

2009

Challenge A

Submissions

Challenge B

Challenge C

Challenge D

2007

2008

CHALLENGE A


Predict the spike timing of a regular spiking L5 pyramidal cell responding to in-vivo-like current injection.

Experimental Methods

The experiments were performed by Thomas Berger and Richard Naud in the laboratory of Henry Markram at the EPFL. A 14-day-old Wistar rat was decapitated and its brain was quickly transferred to a slicing chamber filled with iced artificial cerebrospinal fluid (ACSF). 300 μ m thick slices of the primary somatosensory neocortex were prepared using a HR2 vibratome (Sigmund Elektronik, Heidelberg, Germany). Slices were incubated at 36°C for 45 min and left at room temperature until recording. The ACSF contained (in mM): 125 NaCl, 2.5 KCl, 25 D-glucose, 25 NaHCO₃, 1.25 NaH₂PO₄, 2 CaCl₂, and 1 MgCl₂.

Somatic recordings were performed at 33-35°C with a Axopatch 200B amplifier (Molecular Devices, Union City, CA) in the current clamp mode. Voltage traces were filtered with a 2.4 kHz Bessel filter. The amplifier was connected to a ITC-18 acquisition board (Instrutech Co, Port Washington, NY), which was in turn connected to a PC or Macintosh running a custom written routine under IgorPro (Wavemetrics, Portland, OR). Patch pipettes were pulled with a Flaming/Brown micropipette puller P-97 (Sutter Instruments Co, Novato, CA) and had an initial resistance of < 4 M Ω . Pipettes were filled with intracellular solution (ICS) containing (in mM): 110 potassium gluconate, 10 KCl, 4 ATP-Mg, 10 Na-Phosphocreatine, 0.3 Na-GTP, 10 HEPES, 30 Mannitol, and 8 Biocytin. Chemicals were provided by SIGMA or MERCK. The liquid junction potential between the ACSF and the ICS was around 12 mV and not corrected for.

The current-clamp stimulus has two parts. The first part is 17.5 seconds of various stimulus waveforms frequently used to calibrate neuron models. It consists of a series of four step current with a duration of 2 seconds and an inter-step rest time of two seconds (one hyperpolarizing and 3 depolarizing steps). The steps are followed by an injection of white noise of two seconds. The white noise injection can be used to remove the artifact



Point neurons: not as bad as all that

About INCF

Resources

Community

PEOPLE

EVENTS

TRAINING

JOB BOARD

TRAVEL G

SUBMISSION

This contest is currently not open
for submission.

SUB NAVIGATION

DIADEM

Spike Time Prediction

2009

Challenge A

Submissions

Challenge B

Challenge C

Challenge D

2007

2008

CHALLENGE A


Predict the spike timing of a re

Experimental Methods

The experiments were performed on a
14-day-old Wistar rat was decapitated and
fluid (ACSF). 300 mm thick slices were
Elektronik, Heidelberg, Germany. The
contained (in mM): 125 NaCl, 2

Somatic recordings were performed in
Union City, CA) in the current-clamp mode.
amplifier was connected to a IT-1600 (World
in turn connected to a PC or Macintosh (Apple
Portland, OR). Patch pipettes were pulled from
Instruments Co, Novato, CA) and had an initial resistance of $< 4 \text{ M}\Omega$. Pipettes were filled with intracellular
solution (ICS) containing (in mM): 110 potassium gluconate, 10 KCl, 4 ATP-Mg, 10 Na-Phosphocreatine,
0.3 Na-GTP, 10 HEPES, 30 Mannitol, and 8 Biocytin. Chemicals were provided by SIGMA or MERCK. The
liquid junction potential between the ACSF and the ICS was around 12 mV and not corrected for.

The current-clamp stimulus has two parts. The first part is 17.5 seconds of various stimulus waveforms
frequently used to calibrate neuron models. It consists of a series of four step current with a duration of 2
seconds and an inter-step rest time of two seconds (one hyperpolarizing and 3 depolarizing steps). The
steps are followed by an injection of white noise of two seconds. The white noise injection can be used to remove the artifact



Point neurons: not as bad as all that

About INCF

Resources

Community

PEOPLE

EVENTS

TRAINING

JOB BOARD

TRAVEL G

SUBMISSION

This contest
for submiss

SUB NAME

DIADEM

Spike Tim

2009

Challe

Sub


Challe

Challe

Challe


2007

2008



liquid junction potential between the ACSF and the ICS was around 12 mV and not corrected for.

The current-clamp stimulus has two parts. The first part is 17.5 seconds of various stimulus waveforms frequently used to calibrate neuron models. It consists of a series of four step current with a duration of 2 seconds and an inter-step rest time of two seconds (one hyperpolarizing and 3 depolarizing steps). The steps are followed by an injection of white noise of two seconds. The white noise injection can be used to remove the artefact



Point neurons: not as bad as all that

- Point neurons won all challenges in all years, despite behaviourally relevant incentives
- The MAT2 model (Kobayashi et al., 2009) only has one dynamic variable more than the standard leaky I&F model
- The arbitrary complexity of biophysical neuron models makes them a parameter fitting nightmare
- Save complexity for when you really need it