


## Part II Pathology lectures: Dr Donald Palmer Thymus Biology and Central Tolerance

The thymus is the unique site of T cell development. Stem cells from the bone marrow migrate to the thymus where they receive the appropriate signals to become immune competent T cells. These signals are from the thymic microenvironment, includes both cell-cell and soluble factors. During development, T cells are required to pass several checkpoints and under several rounds of selection to become restricted and self-tolerant CD4<sup>+</sup> and CD8<sup>+</sup> T cells. The thymus is responsible for the formation of self-restricted, immunocompetent T cells. A schematic diagram of T cell development is below.



MHC class I and MHC class II are required for positive selection of CD8<sup>+</sup> and CD4<sup>+</sup> single positive (SP) thymocytes respectively. Positive selection occurs when thymocytes expressing a functional TCR are able to recognise self-MHC and peptide with an appropriate affinity (Naeher *et al.*, 2007). Thymocytes that bind with a high affinity are killed by apoptosis, protecting from autoreactive T cells, and those that bind with a low affinity die by neglect. After successful positive selection the cells become SP (CD4<sup>+</sup>CD8<sup>-</sup> or CD8<sup>+</sup>CD4<sup>+</sup>) and migrate to the medullary region via a chemokine gradient. Once in the medullary region, these cells undergo a process called negative selection which forms part of central tolerance, preventing the circulation of autoreactive T cells in the periphery. Negative selection occurs primarily by mTEC which are able to express peripheral, tissue-specific antigens. This phenomenon occurs via the expression of autoimmune regulator (AIRE) gene by mTEC, which acts as a novel transcription regulator allowing the production of gene products that are tissue-specific antigens.

### At the end of these two lectures, students should be able to:

1. Describe how T cells recognise antigen.
2. Describe the basic structure of the T cell receptor and how T cell receptor diversity is achieved.
3. Describe function of the thymus.
4. Describe the developmental pathway of T cells.
5. Describe the role of the microenvironment in T cell differentiation.
6. Describe how tolerance is generated.

### References:

- Janeway's Immunobiology: 7<sup>th</sup> ed. (Garland Sci). Development and Survival of Lymphocytes: <http://www.garlandscience.com/textbooks/0815341237/pdf/chapter07.pdf>
- Anderson G & Jenkinson EJ. (2001) Lymphostromal interactions in thymic development and function. *Nature Immunol.* **1**: 31-40.
- German RN (2002) T-cell development and the CD4-CD8 lineage decision. *Nature Rev Immunol* **2**:309-322.
- Ritter MA & Palmer DB (1998) The human thymic microenvironment: New approaches to functional analysis. *Sem Immunol.* **11**: 13-21.
- Berg LJ & Kang J (2001) Molecular determinants of TCR expression and selection. *Curr. Opin. Immunol.* **13**: 232-241.

Kruisbeek AM *et al.* (2000) Branching out to gain control: how the pre-TCR is linked to multiple functions. *Immunol. Today* **21**: 637-644.

Aw D, Silva, AB & Palmer DB (2007) Immunosenescence: emerging challenges for an ageing population. *Immunology* **120**: 435-46.

Taams LS *et al.* (2006) Regulatory T cells in human disease and their potential for therapeutic manipulation. *Immunology* **118**:1-9