

Prescription Summary Report

Generated on 07 October 2026

Medicines

Medicine	Strength	How often	Check
Dolo 650	650 mg tablet	1-1-1 after food, 3 days (Morning, afternoon and night, after food, for 3 days)	In our verified list
Pan 40	40 mg tablet	1-0-0 before breakfast, 5 days (Morning, before breakfast, for 5 days)	In our verified list
Azithral 500	500 mg tablet	1-0-0 after food, 3 days (Morning, after food, for 3 days)	In our verified list
Cetzine	10 mg tablet	0-0-1 at bedtime, 5 days (Night, at bedtime, for 5 days)	In our verified list

About your medicines

Dolo 650 (Paracetamol 650 mg)

Used for: Pain relief and fever reduction

Precautions:

- Do not exceed recommended daily dose
- Avoid alcohol while taking
- Check liver health if you have liver disease

Common side effects:

- Nausea
- Stomach upset
- Rash

See a doctor if:

- Severe skin rash
- Yellowing of skin or eyes
- Persistent vomiting

Source: our verified medicine list

Pan 40 (Pantoprazole 40 mg)

Used for: Acid reflux and stomach ulcers

Precautions:

- Take before a meal
- Tell doctor if you have liver disease
- Do not crush or chew

Common side effects:

- Headache
- Diarrhoea
- Stomach pain

See a doctor if:

- Severe stomach pain
- Vomiting blood
- Persistent diarrhea

Source: our verified medicine list

Azithral 500 (Azithromycin 500 mg)

Used for: Bacterial infections such as respiratory, skin, and ear infections

Precautions:

- Take with a full glass of water
- Do not take with antacids containing aluminum or magnesium
- Complete the full course even if you feel better
- Tell doctor if you have liver disease

Common side effects:

- Nausea or vomiting
- Diarrhea
- Stomach pain
- Headache

See a doctor if:

- Severe diarrhea or bloody stools
- Allergic rash or swelling
- Chest pain or irregular heartbeat
- Persistent vomiting

Source: our verified medicine list

Cetzine (Cetirizine 10 mg)

Used for: Allergy symptoms like sneezing and itchy eyes

Precautions:

- May cause drowsiness, avoid driving if sleepy
- Take with water, can be taken with or without food

Common side effects:

- Sleepiness
- Dry mouth
- Headache

See a doctor if:

- Severe rash or swelling
- Difficulty breathing
- Fast or irregular heartbeat

Source: our verified medicine list

Summary in simple words

Dolo 650: 1-1-1 after food, 3 days

Pan 40: 1-0-0 before breakfast, 5 days

Azithral 500: 1-0-0 after food, 3 days

Cetzine: 0-0-1 at bedtime, 5 days

Please follow your doctor's advice and ask the doctor or pharmacist if you are unsure.

Good questions to ask your doctor or pharmacist

- What is each medicine for, and what should I watch out for?
- Should I take it before or after food?
- What should I do if I miss a dose?
- When should I come back if I do not feel better?

This report is generated by an AI assistant for a hackathon. Always consult a human doctor or pharmacist before making any medical decisions.