

Package ‘survtd’

April 25, 2017

Version 0.0.0.9002

Title Survival analysis with time-dependent covariates

Depends R (>= 3.1.0), lme4, survival, timereg, mice

Imports boot, ipw, stringr, MASS, Matrix

Description Fit the Cox proportional hazards model or the semi-parametric additive hazards model with time-fixed covariates of any type and continuous time-dependent covariates subject to measurement error, missing data, or simply observed in discrete time: these issues are all handled with a two-stage joint modelling approach using multiple imputation (see Moreno-Betancur et al. 2016). The package also provides a naive two-stage approach (Naive2S, for comparison) and an easy-to-use implementation of the last observation carried forward (LOCF) approach which is the traditional method to incorporate time-dependent covariates of any type in hazard models. Data simulation functions also included.

License GPL (>= 3)

LazyData TRUE

RoxygenNote 6.0.1

Author Margarita Moreno-Betancur [cre, aut],
Samuel L Brilleman [ctb]

Maintainer Margarita Moreno-Betancur <margarita.moreno@mcri.edu.au>

R topics documented:

simjm	1
simjm_benchmark	3
survtd	4

Index	8
--------------	----------

simjm	<i>Simulate data from a joint model of multiple continuous time-dependent markers and a time-to-event</i>
-------	---

Description

simjm simulates data from a Cox proportional hazards or semi-parametric additive model with two time-fixed covariates (Z1 and Z2) and three time-dependent covariates (Yij_1, Yij_2, Yij_3). The user can specify various characteristics of these distributions.

Usage

```
simjm(n = 200, surv_model = "Cox", marker_model = "RE", MErr = "High",
      Miss = "Low", effects = "Weak", corr = "Low")
```

Arguments

n	Size of dataset to be generated
surv_model	Model for the time-to-event. Options are "Cox" for Cox proportional hazard model and "Add" for semi-parametric additive hazards model.
marker_model	Model for the multiple markers. Options are "RE" for the correlated random effects model and "PN" for the product normal model.
MErr	Degree of measurement error in the multiple markers. Options are "Low", "Mod" and "High".
Miss	Degree of missing at-risk measurements. Options are "None", "Low" and "High".
effects	Strength of effects of the markers on the hazard. Options are "Null", "Weak" and "Strong".
corr	Degree of marginal pairwise correlations between the markers. Options are "Low", "Mod" and "High".

Details

The function can be used to generate data from any of the scenarios considered in the the main simulation settings of Moreno-Betancur et al. (2016). See that reference for details.

Value

A data.frame as required by [survtd](#). That is, in the long format, with one row per individual and per visit time at which any of the time-dependent covariates were measured, with the corresponding measurements. The dataset also includes a variable that uniquely identifies observations from the same individual; a variable that indicates the timing of each measurement visit; and the fixed variables (time-to-event, event indicator, time-fixed covariates) which are constant across rows of the same individual. The final columns of the dataset (from `Xij_1` onwards) are to recover the true values of the markers as per the data generation model for use with function [simjm_benchmark](#). Specifically, the variables in the dataset are:

ID Unique identifier of observations from the same individual.

tt Time to event, possibly right-censored.

event Indicator of event, with event=1 if an event occurred at tt and event=0 if the individual is censored.

Z1 Time-fixed binary covariate.

Z2 Time-fixed continuous covariate.

tj Timing of the measurement visit.

Yij_1 Measured value of marker 1 at time tj

Yij_2 Measured value of marker 2 at time tj

Yij_3 Measured value of marker 3 at time tj

Xij_1 True value of marker 1 at time tj

Xij_2 True value of marker 2 at time tj

Xij_3 True value of marker 3 at time tj

fixed_1 Time-fixed part of the linear predictor of the linear mixed model from which Y_{ij_1} is generated.

tim_1 Time-dependent part of the linear predictor of the linear mixed model from which Y_{ij_1} is generated, excluding terms for other markers in the case of product-normal model.

fixed_2 Time-fixed part of the linear predictor of the linear mixed model from which Y_{ij_2} is generated.

tim_2 Time-dependent part of the linear predictor of the linear mixed model from which Y_{ij_2} is generated, excluding terms for other markers in the case of product-normal model.

fixed_3 Time-fixed part of the linear predictor of the linear mixed model from which Y_{ij_3} is generated.

tim_3 Time-dependent part of the linear predictor of the linear mixed model from which Y_{ij_3} is generated, excluding terms for other markers in the case of product-normal model.

References

Moreno-Betancur M, Carlin JB, Brilleman SL, Tanamas S, Peeters A, Wolfe R (2016). Survival analysis with time-dependent covariates subject to measurement error and missing data: Two-stage joint model using multiple imputation. Submitted.

Examples

```
dat<-simjm(n=200,surv_model="Cox",marker_model="PN",
           MErr="High",Miss="None",effects="Weak",corr="Low")

head(dat)

dat<-simjm(n=200,surv_model="Add",marker_model="RE",
           MErr="Low",Miss="High",effects="Strong",corr="Mod")

head(dat)
```

simjm_benchmark

Benchmark "True" analysis of data simulated by simjm function

Description

simjm_benchmark fits Cox proportional hazards and semi-parametric additive hazards models to data generated by [simjm](#) using the perfect data, i.e. the true values of the time-dependent markers.

Usage

```
simjm_benchmark(data, surv_model = "Cox", marker_model = "RE",
                corr = "Low")
```

Arguments

<code>data</code>	A <code>data.frame</code> produced by <code>simjm</code> .
<code>surv_model</code>	Time-to-event model that was used to generate data. Options are "Cox" for Cox proportional hazard model and "Add" for semi-parametric additive hazards model.
<code>marker_model</code>	Multiple marker model that was used to generate data. Options are "RE" for the correlated random effects model and "PN" for the product normal model.
<code>corr</code>	Degree of marginal pairwise correlations between the markers that was used to generate data. Options are "Low", "Mod" and "High".

Details

The function is used to perform what is referred to as the "True" analysis in Moreno-Betancur et al. (2016) which is based on perfect data, i.e. the true values of the markers at each of the event times as drawn from the simulation model. This provides a benchmark for analyses performed with error-polluted and incomplete data using `survtd`.

Value

Returns regression coefficient estimates for each covariate based on the "True" analysis, along with 95 confidence intervals and p-values.

References

Moreno-Betancur M, Carlin JB, Brilleman SL, Tanamas S, Peeters A, Wolfe R (2016). Survival analysis with time-dependent covariates subject to measurement error and missing data: Two-stage joint model using multiple imputation. Submitted.

Examples

```
dat<-simjm(n=200,surv_model="Cox",marker_model="PN",
           MErr="High",Miss="None",effects="Weak",corr="Low")

simjm_benchmark(dat,surv_model="Cox",marker_model="PN",corr="Low")
```

survtd

Fit survival hazard models with time-dependent covariates

Description

`survtd` fits Cox proportional hazards and semi-parametric additive hazards models with time-fixed covariates (any type) and time-dependent covariates either with a multiple imputation-based two-stage joint modelling approach to deal with missing data and measurement error (only continuous time-dependent covariates), a naive two-stage approach (only continuous time-dependent covariates) or the LOCF approach (any type).

Usage

```
survtd(formula, data, id, visit.time, model = "Cox", method = "MICEa",
        M = 5, G = 5, time.trend = as.formula("~x"))
```

Arguments

formula	A formula object, with the response on the left of a <code>~</code> operator, and the terms on the right as regressors. The response must be a survival object as returned by the <code>Surv</code> function of type "right" (other types are not supported at this stage). Time-dependent regressors are specified by the wrapper <code>td()</code> (see Examples section below). This development version does not support interactions nor terms constructed using <code>I()</code> . The user thus needs to create the necessary interaction/derived variables in the dataset before using this function.
data	A <code>data.frame</code> in which to interpret the variables named in the formula. The dataset needs to be in long format, with one row per individual and per visit time at which any of the time-dependent covariates were measured, with the corresponding measurements. The dataset must also include a variable that uniquely identifies observations from the same individual (see parameter <code>id</code> below); a variable that indicates the timing of each measurement visit (see parameter <code>visit.time</code> below); and other fixed variables (time-to-event, event indicator, time-fixed covariates) which are constant across rows of the same individual.
id	The name of the variable in <code>data</code> that uniquely identifies observations from the same individual.
visit.time	The name of the variable in <code>data</code> that indicates the timing of each visit.
model	Indicates which hazard model to fit. Options are "Cox" for the Cox proportional hazard model and "Add" for the semi-parametric additive hazards model.
method	Indicates which method to use to fit the model. Options are "MICEa", "MICEb", "LOCF" and "Naive2S". Both "MICEa" and "MICEb" use a two-stage joint modelling approach based on multiple imputation to incorporate time-varying continuous covariates (see Details section below). Method "LOCF" uses the Last Observation Carried Forward (LOCF) approach. Method "Naive2S" is the naive two-stage approach (see Details section below).
M	Number of imputations to perform for methods "MICEa" and "MICEb" (ignored if <code>method="LOCF"</code> or <code>"Naive2S"</code>).
G	Number of iterations to perform in multiple imputation by chained equations (MICE) algorithm for methods "MICEa" and "MICEb" (ignored if <code>method="LOCF"</code> or <code>"Naive2S"</code>).
time.trend	Formula object with empty left hand side and right hand side expressing a polynomial of "x" which determines the way time is modeled in the fixed effects part of the linear mixed model for each time-dependent marker (ignored if <code>method="LOCF"</code>). Default is a linear trend (i.e. include <code>visit.time</code> as predictor in the model). The random effects part includes a random intercept and slope. Future development plans are to allow a different <code>time.trend</code> argument for each marker, the possibility to include natural cubic splines in this argument and more general random effects structures.

Details

The `survtd` function can be used to fit the Cox proportional hazards or semi-parametric additive hazard models with time-dependent covariates.

Methods "MICEa" and "MICEb" can be used to fit models with time-fixed covariates of any type and multiple continuous time-dependent covariates. To deal with the discrete-time observation of the time-varying covariates, particularly measurement error and missing data, a two-stage joint modeling approach is used that is based on multiple imputation. Details are provided in Moreno-Betancur et al. (2016), but briefly, in Stage 1 the true (error-corrected) values of the time-dependent

covariates at each event time at which an individual is at risk are multiply imputed by drawing iteratively from smoothed trajectories based on interdependent linear mixed models using Multiple Imputation by Chained Equations (MICE). An adaptation of the `mice` function from the *mice* package is used for this step. In Stage 2, the time-to-event model is fitted to each of the imputed datasets (see below) and estimates are pooled using Rubin's MI formulas.

The two MICE methods differ in the way information about the event occurrence is included in the imputation models: the "MICEb" method includes the event indicator, while the "MICEa" approach is based on a more refined approximation including a modified event indicator (see Moreno-Betancur et al. 2016). Hence, the approach "MICEa" is generally preferable.

The "Naive2S" method can be used to fit models with time-fixed covariates of any type and multiple continuous time-dependent covariates using the naive two-stage approach. This method singly imputes each continuous marker at each event time from its estimated trajectory obtained from a linear mixed model including fixed and random effects for time and fixed effects for the time-fixed covariates appearing in the formula argument. The method thus ignores: the uncertainty in these imputed values, the interrelations between the time-dependent markers and the relation between the time-dependent markers and the time-to-event process (see Moreno-Betancur et al. 2016).

The "LOCF" method can be used with any type of time-fixed and time-varying covariates (categorical or continuous). This approach uses the last available measurement of the time-varying covariate to singly impute its value at each event time at which an individual is at risk. It can perform very poorly if the observations are not synchronously updated across individuals (e.g. due to missing data) and if the time-varying covariates are measured with error (see Moreno-Betancur et al. 2016).

Once the values of the time-dependent covariates are imputed at each of the event-times at which the individual is at risk according to either of the four methods, the function uses `coxph` from the *survival* package to fit the Cox model and `aalen` from the *timereg* package to fit the additive model.

Value

This development version returns a data frame with regression coefficient estimates for each covariate based on the method chosen, along with standard errors, 95% confidence intervals and p-values. This will change in future versions when a proper class of objects and summary and other such methods are developed.

References

Moreno-Betancur M, Carlin JB, Brilleman SL, Tanamas S, Peeters A, Wolfe R (2016). Survival analysis with time-dependent covariates subject to measurement error and missing data: Two-stage joint model using multiple imputation. Submitted.

Examples

```
## Example with additive model ##

dat<-simjm(n=200,surv_model="Add",marker_model="RE",
          MErr="Low",Miss="High",effects="Strong",corr="Mod")

survtd(Surv(time=tt,event=event)~Z1+Z2+td(Yij_1)+td(Yij_2)+td(Yij_3),
       data=dat, id="ID", visit.time="tj", model="Add",
       method="MICEa", M=5, G=5,time.trend=as.formula("~x+I(x^2)"))

survtd(Surv(time=tt,event=event)~Z1+Z2+td(Yij_1)+td(Yij_2)+td(Yij_3),
       data=dat, id="ID", visit.time="tj", model="Add",
       method="Naive2S", M=5, G=5,time.trend=as.formula("~x+I(x^2)"))
```

```
survtd(Surv(time=tt,event=event)~Z1+Z2+td(Yij_1)+td(Yij_2)+td(Yij_3),  
      data=dat, id="ID", visit.time="tj", model="Add",  
      method="LOCF", M=5, G=5)
```

```
simjm_benchmark(dat,surv_model="Add",marker_model="RE",corr="Mod")
```

Index

simjm, [1](#), [3](#), [4](#)
simjm_benchmark, [2](#), [3](#)
survtd, [2](#), [4](#), [4](#)