

# Fuel analysis for Oak Symposium: Duff and litter

Morgan Gray

2022-10-25

---

## Overview

The aim of this research was to understand the impact of the 2017 Tubbs fire on fuel loads. The objective of this research was to evaluate the impact of the 2017 Tubbs fire on duff and litter fuel loads over time. To meet this objective, we compared the pre- and post-fire levels of duff and litter at nine long-term vegetation monitoring plots. Surveys were conducted at four time points: 2016 (1y pre-fire), 2017 (0y post-fire), 2019 (2y post-fire), and 2021 (4y post-fire). We conducted surveys along two transects per plot; each transect had two quadrats. As a result, each time point had 36 data points: 9 plots x 2 transects x 2 quadrats.

Using the duff and litter data set as input, we asked the following questions:

1. Did the total fuel load (all duff and litter combined) differ between years?
2. Did fuel load differ between years for duff and/or litter? If so, which years were different?

**Duff and litter data** We answered these questions using a time series of plot-level data for (1) the total amount of duff and litter and (2) the mean amount of each fuel class (duff, litter). We defined “total (or all) duff and litter” as the combined total of the duff and litter within a plot.

First, we calculated the plot-level mean ( $n = 4$  quadrats) for each fuel class (duff, litter) at each time point. A subset of the data frame with the plot-level mean for each fuel class by year is shown below.

fuel_type	fuel_class	plot_id	year	value
Duff & litter	Duff	RxF01	2016	1.125
Duff & litter	Litter	RxF01	2016	0.625
Duff & litter	Duff	RxF02	2016	1.562
Duff & litter	Litter	RxF02	2016	2.375
Duff & litter	Duff	RxF03	2016	1.062
Duff & litter	Litter	RxF03	2016	1.938

Next, we summed the means for the duff and litter fuel classes to find the plot-level total at each time point. Below is a subset of the data frame for the plot-level values of duff and litter by year.

fuel_type	fuel_class	plot_id	year	value
Duff & litter	All	RxF01	2016	1.750
Duff & litter	All	RxF02	2016	3.938
Duff & litter	All	RxF03	2016	3.000
Duff & litter	All	RxF04	2016	2.625
Duff & litter	All	RxF05	2016	1.625
Duff & litter	All	RxF06	2016	3.013

**Statistical methods** Statistically significant differences were identified using repeated measures analysis of variance (ANOVA) and post hoc comparisons. P-values were adjusted using the Bonferroni multiple testing correction method.

The repeated-measures ANOVA is used for analyzing data where same subjects are measured more than once. This test is also referred to as a within-subjects ANOVA or ANOVA with repeated measures. The “within-subjects” term means that the same individuals (here, individuals are plots) are measured on the same outcome variable under different time points. The main goal of a repeated measures ANOVA is to evaluate if there is a statistically significant interaction effect between within-subjects factors in explaining a continuous outcome variable. The repeated measures ANOVA makes the following assumptions about the data:

- No significant outliers in any cell of the design
- Normality: the outcome (or dependent) variable should be approximately normally distributed in each cell of the design
- Assumption of sphericity: the variance of the differences between groups should be equal

We assessed outliers using the the interquartile range (IQR;  $IQR = Q3 - Q1$ ). Values above  $Q3 + 1.5 \times IQR$  or below  $Q1 - 1.5 \times IQR$  are considered as outliers. Values above  $Q3 + 3 \times IQR$  or below  $Q1 - 3 \times IQR$  are considered as extreme points (or extreme outliers).  $Q1$  and  $Q3$  are the first and third quartile, respectively. Extreme outliers can be due to data entry errors, measurement errors, or unusual values. The outlier may be included if one believes the result will not be substantially affected; this can be evaluated by comparing the result of the ANOVA with and without the outlier.

We assessed normality by visual inspection of a QQ plot for each time point. A QQ plot draws the correlation between a given data and the normal distribution. We also conducted the Shapiro-Wilk test for each time point. Using this method, normally distributed data will have p-value  $> 0.05$ .

The assumption of sphericity was checked during the computation of the ANOVA test using the R function `anova_test()` [rstatix package]. The Mauchly’s test was internally used to assess the sphericity assumption, and the Greenhouse-Geisser sphericity correction was automatically applied to factors violating the sphericity assumption.

---

## Q1: Did total fuel load differ between years?

The first question we asked was whether there was a significant main effect of year on total fuel load. A one-way repeated measures ANOVA was used to determine whether fuel load was significantly different between the four time points.


### Summary statistics for the plot-level total amount of duff and litter by year

The following table summarizes the plot-level total amount of duff and litter for each time point.

data_type	fuel_class	year	mean	sd	n
Duff & litter	All	2016	3.113	1.012	9
Duff & litter	All	2017	0.729	0.420	9
Duff & litter	All	2019	1.861	1.154	9
Duff & litter	All	2021	0.926	0.500	9

## Visualization of the plot-level total amount of duff and litter by year

A box plot of total duff and litter by year showed lower values in post-fire years. Specifically, we observed a decrease in the depth of all duff and litter immediately following the fire (2017; 0-y post-fire). The mean depth of duff and litter did not return to pre-fire levels in subsequent years.



## Check assumptions


**Check for outliers** There was one extreme outlier in the data set. Specifically, the value for plot RxF03 in 2017 was 1.738.

data_type	fuel_class	year	plot_id	value	is_outlier	is_extreme
Duff & litter	All	2017	RxF03	1.738	TRUE	TRUE

**Check for normality** Plot-level duff and litter load was normally distributed at each time point ( $p > 0.01$ ), except for year 2017, as assessed by Shapiro-Wilk's test.

data_type	fuel_class	year	is_normal	p	statistic
Duff & litter	All	2017	FALSE	0.0095626	0.771

The QQ plot showed one point in 2017 that was distant from the reference line.



### Repeated measures ANOVA test

Total duff and litter load differed between years. We found a significant main effect of year on duff and litter load (p-adjusted < 0.001). The mean fuel load for all duff and litter (duff and litter combined) was significantly different between the four years;  $F(3, 24) = 14.029$ , p-adjusted (p-adj.) =  $1.73e-05$ , ges = 0.59. The generalized effect size (ges) is the amount of variability due to the within-subjects factor.

data_type	fuel_class	effect	p_adj	statistic	d_fn	d_fd	ges	method
Duff & litter	All	year	1.73e-05	14.029	3	24	0.59	me


### Post hoc tests

We conducted post hoc pairwise comparisons between the levels of the within-subjects factor (here, year). The result of paired t-tests between years showed a significant difference between pre- and post-fire mean duff and litter levels in 2017 (p-adj. <0.001) and 2021 (p-adj <0.01).

The results from the pairwise comparisons are shown below as (1) a boxplot of total duff and litter by year updated with p-values, and (2) a table of test results.

## Duff & litter (all)

Anova,  $F(3,24) = 14.03$ ,  $p = <0.0001$ ,  $\eta_g^2 = 0.59$



data_type	fuel_class	group1	group2	p_adj	p_adj_sig	statistic	df	method
Duff & litter	All	2016	2017	0.000828	***	6.802	8	pwc
Duff & litter	All	2016	2021	0.003000	**	5.548	8	pwc
Duff & litter	All	2017	2019	0.205000	ns	-2.552	8	pwc
Duff & litter	All	2016	2019	0.267000	ns	2.380	8	pwc
Duff & litter	All	2019	2021	0.440000	ns	2.061	8	pwc
Duff & litter	All	2017	2021	1.000000	ns	-0.828	8	pwc

## Q2: Did fuel load differ between years by fuel class?

Next, we investigated whether there was a significant change in plot-level fuel load over time when accounting for fuel class (duff, litter). A two-way repeated measures ANOVA was used to determine whether there was a significant interaction between year and fuel class on fuel load.

Here, the effect of year on fuel load was our focal variable of primary concern. However, the effect of year may differ between duff and litter, so fuel\_class was considered a moderator variable.

### Summary statistics for duff and litter by year

The following table summarizes duff and litter by class for each time point.

data_type	fuel_class	statistic	2016	2017	2019	2021
Duff & litter	duff	mean	1.569	0.438	0.085	0.146
Duff & litter	duff	sd	0.485	0.132	0.085	0.133
Duff & litter	litter	mean	1.543	0.292	1.776	0.781
Duff & litter	litter	sd	0.745	0.316	1.134	0.467

### Visualization of duff and litter by year


A box plot of duff and litter by year showed differing post-fire trends. Although duff and litter both showed a decrease in depth immediately following the fire (2017; 0-y post-fire), the subsequent annual trends differed. The temporal trend for total duff and litter was more similar to that of the litter fuel class, and showed less similarity to that of the duff fuel class. These results suggest that litter was an influential driver of the post-fire patterns observed for duff and litter as a group.

#### Duff

- Duff depth remained low in all post-fire years.
- Values in 2019 and 2021 were lower than those for 2017.

#### Litter

- Litter depth returned to pre-fire levels 2-y after the fire (2019), then showed a second decrease in 2021 (4-y post-fire).
- This post-fire increase and decrease in litter depth was similar to the temporal trend observed for duff and litter overall.




## Check assumptions

**Check for outliers** There were two extreme outliers in the data set. Specifically, the values for duff at plots RxF03 and RxF04 in 2021 were 0.112 and 0.500, respectively.

data_type	fuel_class	year	plot_id	value	is_outlier	is_extreme
Duff & litter	duff	2021	RxF03	0.112	TRUE	TRUE
Duff & litter	duff	2021	RxF04	0.500	TRUE	TRUE

**Check for normality** Plot-level litter and duff values were normally distributed at each time point ( $p > 0.01$ ), except for year duff in 2021 and litter in 2017, as assessed by Shapiro-Wilk's test.

data_type	fuel_class	year	is_normal	p	statistic
Duff & litter	duff	2021	FALSE	0.0000005	0.406
Duff & litter	litter	2017	FALSE	0.0065427	0.757



## Repeated measures ANOVA test





There was a statistically significant two-way interaction between duff and litter and year,  $F(2, 14) = 10.382$ ,  $p_{\text{adj.}} = 0.006$ ,  $\text{ges} = 0.329$ . A significant two-way interaction indicates that the impact of fuel class on fuel load depends on year (and vice versa).


data_type	effect	p_adj	p_adj_sig	statistic	d_fn	d_fd	ges	method
Duff & litter	year	5.19e-05	***	14.029	3.00	24.00	0.451	aov2
Duff & litter	fuel_class	5.82e-04	***	41.884	1.00	8.00	0.211	aov2
Duff & litter	fuel_class:year	6.00e-03	**	10.382	1.75	13.99	0.329	aov2

## Post hoc tests

We found a significant main effect of year on fuel load for both duff and litter. One thing the pairwise comparisons by fuel class had in common: For both duff and litter, the fuel load in 2016 was significantly different from that in 2017 (duff:  $p < 0.001$ ; litter:  $p < 0.01$ ).

### Duff and litter

Year  1-y pre-fire (2016)  0-y post-fire (2017)  2-y post-fire (2019)  4-y post-fire (2021)







data_type	fuel_class	effect	p_adj	p_adj_sig	statistic	d_fn	d_fd	ges	method
Duff & litter	duff	year	1.07e-05	***	62.909	1.29	10.3	0.853	me
Duff & litter	litter	year	4.00e-03	**	6.830	3.00	24.0	0.423	me


The following sections present the results from post hoc tests by fuel class for (1) the main effect of fuel class on fuel load at each time point, and (2) significant pairwise differences between years.

**Duff vs. Year** There was a statistically significant main effect of year on duff;  $F(3, 24) = 62.909$ ,  $p\text{-adj.} = 1.07e-05$ ,  $ges = 0.853$ . Pairwise comparisons showed that the mean duff depth was significantly different between all years except 2019/2021. This result indicated that duff depth differed between 2016 (pre-fire) and all post-fire years, as well as between 2017 (0y post-fire) and subsequent years.



## Duff





Year  1-y pre-fire (2016)  0-y post-fire (2017)  2-y post-fire (2019)  4-y post-fire (2021)




data_type	fuel_class	group1	group2	p_adj	p_adj_sig	statistic	df	method
Duff & litter	Duff	2016	2019	9.24e-05	****	9.232	8	pwc
Duff & litter	Duff	2016	2021	2.08e-04	***	8.263	8	pwc
Duff & litter	Duff	2016	2017	7.38e-04	***	6.909	8	pwc
Duff & litter	Duff	2017	2019	1.00e-03	**	6.541	8	pwc
Duff & litter	Duff	2017	2021	5.00e-03	**	5.126	8	pwc
Duff & litter	Duff	2019	2021	1.00e+00	ns	-1.226	8	pwc

**Litter vs. Year** There was a statistically significant main effect of year on litter;  $F(1, 10) = 6.83$ ,  $p\text{-adj.} = 0.004$ ,  $ges = 0.423$ . Pairwise comparisons showed that the mean litter depth was significantly different between 2016 and 2017 ( $p\text{-adj.} < 0.01$ ), as well as 2017 and 2019 ( $p\text{-adj.} < 0.05$ ).


## Litter

Year  1-y pre-fire (2016)  0-y post-fire (2017)  2-y post-fire (2019)  4-y post-fire (



data_type	fuel_class	group1	group2	p_adj	p_adj_sig	statistic	df	method
Duff & litter	Litter	2016	2017	0.006	**	5.058	8	pwc
Duff & litter	Litter	2017	2019	0.046	*	-3.536	8	pwc
Duff & litter	Litter	2017	2021	0.248	ns	-2.429	8	pwc
Duff & litter	Litter	2016	2021	0.273	ns	2.367	8	pwc
Duff & litter	Litter	2019	2021	0.335	ns	2.236	8	pwc
Duff & litter	Litter	2016	2019	1.000	ns	-0.471	8	pwc

**Duff vs. Litter** We found a significant difference in mean depth between duff and litter in 2019 and in 2021.



data_type	year	group1	group2	p_adj	p_adj_signif	statistic	df
Duff & litter	2016	duff	litter	0.919	ns	0.105	8
Duff & litter	2017	duff	litter	0.108	ns	1.810	8
Duff & litter	2019	duff	litter	0.002	**	-4.526	8
Duff & litter	2021	duff	litter	0.004	**	-4.046	8

---

Date of last revision  
2022-10-25

---