

4.3 Measurement and expectation values

Slides: Video 4.3.5 Time evolution and the Hamiltonian

Text reference: Quantum Mechanics for Scientists and Engineers

Section 3.11





Measurement and expectation values



Time evolution and the Hamiltonian

Quantum mechanics for scientists and engineers

David Miller

Time evolution and the Hamiltonian

Taking Schrödinger's time dependent equation

$$\hat{H}\Psi(\mathbf{r},t) = i\hbar \frac{\partial \Psi(\mathbf{r},t)}{\partial t}$$

and rewriting it as

$$\frac{\partial \Psi(\mathbf{r},t)}{\partial t} = -\frac{i\hat{H}}{\hbar} \Psi(\mathbf{r},t)$$

and presuming \hat{H} does not depend explicitly on time

i.e., the potential $V(\mathbf{r})$ is constant

could we somehow legally write

$$\Psi(\mathbf{r},t_1) = \exp\left(-\frac{i\hat{H}(t_1 - t_0)}{\hbar}\right) \Psi(\mathbf{r},t_0)$$

Time evolution and the Hamiltonian

Certainly,

if the Hamiltonian operator \hat{H} here was replaced by a constant number

we could perform such an integration of

$$\frac{\partial \Psi(\mathbf{r}, t)}{\partial t} = -\frac{i\hat{H}}{\hbar} \Psi(\mathbf{r}, t)$$

to get

$$\Psi(\mathbf{r}, t_1) = \exp\left(-\frac{i\hat{H}(t_1 - t_0)}{\hbar}\right) \Psi(\mathbf{r}, t_0)$$

Time evolution and the Hamiltonian

If, with some careful definition, it was legal to do this

then we would have an operator that

gives us the state at time t_1 directly from that at time t_0

To think about this “legality”

first we note that, because \hat{H} is a linear operator

for any number a

$$\hat{H} [a\Psi(\mathbf{r}, t)] = a\hat{H}\Psi(\mathbf{r}, t)$$

Since this works for any function $\Psi(\mathbf{r}, t)$

we can write as a shorthand

$$\hat{H}a \equiv a\hat{H}$$

Time evolution and the Hamiltonian

Next we have to define what we mean by an operator raised to a power

By \hat{H}^2 we mean $\hat{H}^2\Psi(\mathbf{r},t) = \hat{H}[\hat{H}\Psi(\mathbf{r},t)]$

Specifically, for example, for the energy eigenfunction $\psi_n(\mathbf{r})$

$$\hat{H}^2\psi_n(\mathbf{r}) = \hat{H}[\hat{H}\psi_n(\mathbf{r})] = \hat{H}[E_n\psi_n(\mathbf{r})] = E_n\hat{H}\psi_n(\mathbf{r}) = E_n^2\psi_n(\mathbf{r})$$

We can proceed inductively to define all higher powers

$$\hat{H}^{m+1} \equiv \hat{H}[\hat{H}^m]$$

which will give, for the an energy eigenfunction

$$\hat{H}^m\psi_n(\mathbf{r}) = E_n^m\psi_n(\mathbf{r})$$

Time evolution and the Hamiltonian

Now let us look at the time evolution of some wavefunction $\Psi(\mathbf{r}, t)$ between times t_0 and t_1

Suppose the wavefunction at time t_0 is $\psi(\mathbf{r})$

which we expand in the energy eigenfunctions $\psi_n(\mathbf{r})$

as
$$\psi(\mathbf{r}) = \sum_n a_n \psi_n(\mathbf{r})$$

Then we know

multiplying by the complex exponential factors for the time-evolution of each basis function

$$\Psi(\mathbf{r}, t_1) = \sum_n a_n \exp\left[-\frac{iE_n(t_1 - t_0)}{\hbar}\right] \psi_n(\mathbf{r})$$

Time evolution and the Hamiltonian

In

$$\Psi(\mathbf{r}, t_1) = \sum_n a_n \exp\left[-\frac{iE_n(t_1 - t_0)}{\hbar}\right] \psi_n(\mathbf{r})$$

noting that $\exp(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$

we can write the exponentials as power series

so

$$\Psi(\mathbf{r}, t_1) = \sum_n a_n \left[1 + \left(-\frac{iE_n(t_1 - t_0)}{\hbar} \right) + \frac{1}{2!} \left(-\frac{iE_n(t_1 - t_0)}{\hbar} \right)^2 + \dots \right] \psi_n(\mathbf{r})$$

Time evolution and the Hamiltonian

In

$$\Psi(\mathbf{r}, t_1) = \sum_n a_n \left[1 + \left(-\frac{iE_n(t_1 - t_0)}{\hbar} \right) + \frac{1}{2!} \left(-\frac{iE_n(t_1 - t_0)}{\hbar} \right)^2 + \dots \right] \psi_n(\mathbf{r})$$

because we showed that $\hat{H}^m \psi_n(\mathbf{r}) = E_n^m \psi_n(\mathbf{r})$
we can substitute to obtain

$$\Psi(\mathbf{r}, t_1) = \sum_n a_n \left[1 + \left(-\frac{i\hat{H}(t_1 - t_0)}{\hbar} \right) + \frac{1}{2!} \left(-\frac{i\hat{H}(t_1 - t_0)}{\hbar} \right)^2 + \dots \right] \psi_n(\mathbf{r})$$

Time evolution and the Hamiltonian

With

$$\Psi(\mathbf{r}, t_1) = \sum_n a_n \left[1 + \left(-\frac{i\hat{H}(t_1 - t_0)}{\hbar} \right) + \frac{1}{2!} \left(-\frac{i\hat{H}(t_1 - t_0)}{\hbar} \right)^2 + \dots \right] \psi_n(\mathbf{r})$$

because the operator \hat{H} and all its powers commute with scalar quantities (numbers) we can rewrite

$$\begin{aligned} \Psi(\mathbf{r}, t_1) &= \left[1 + \left(-\frac{i\hat{H}(t_1 - t_0)}{\hbar} \right) + \frac{1}{2!} \left(-\frac{i\hat{H}(t_1 - t_0)}{\hbar} \right)^2 + \dots \right] \sum_n a_n \psi_n(\mathbf{r}) \\ &= \left[1 + \left(-\frac{i\hat{H}(t_1 - t_0)}{\hbar} \right) + \frac{1}{2!} \left(-\frac{i\hat{H}(t_1 - t_0)}{\hbar} \right)^2 + \dots \right] \Psi(\mathbf{r}, t_0) \end{aligned}$$

Time evolution and the Hamiltonian

So, provided we define the exponential of the operator in terms of a power series, i.e.,

$$\exp\left[-\frac{i\hat{H}(t_1-t_0)}{\hbar}\right] \equiv \left[1 + \left(-\frac{i\hat{H}(t_1-t_0)}{\hbar}\right) + \frac{1}{2!}\left(-\frac{i\hat{H}(t_1-t_0)}{\hbar}\right)^2 + \dots\right]$$

then we can write our preceding expression as

$$\Psi(\mathbf{r}, t_1) = \exp\left(-\frac{i\hat{H}(t_1-t_0)}{\hbar}\right) \Psi(\mathbf{r}, t_0)$$

Time evolution and the Hamiltonian

Hence we have established that

there is a well-defined operator that

given the quantum mechanical wavefunction or
"state" at time t_0

will tell us what the state is at a time t_1

$$\Psi(\mathbf{r}, t_1) = \exp\left(-\frac{i\hat{H}(t_1 - t_0)}{\hbar}\right) \Psi(\mathbf{r}, t_0)$$

