



Sums, products and power series

Background mathematics review

David Miller



Sums, products and power series



Sum, factorial and product notations

Background mathematics review

David Miller

Summation notation

If we want to add a set of numbers $a_1, a_2, a_3,$
and a_4 , we can write

$$S = a_1 + a_2 + a_3 + a_4$$

or we can use summation notation

$$S = \sum_{j=1}^4 a_j \equiv \sum_{j=1,2,3,4} a_j \equiv \sum_{j=1,\dots,4} a_j$$

If the range of j is obvious

$$S = \sum_j a_j$$

Here j is an "index"

plural of index –
"indexes"
or
"indices"

Example – arithmetic series

For a set of numbers spaced by a constant amount,
an “arithmetic progression or sequence”

i.e., where the n th number is $a_n = a_1 + (n-1)d$

e.g., the numbers 4, 7, 10, and 13

i.e., $a_1 = 4$, $a_2 = 7$, $a_3 = 10$ and $a_4 = 13$

with, therefore $d = 3$ and $m = 4$ terms in the
progression

gives the “series” (sum of the terms)

$$a_1 + a_2 + a_3 + a_4 = 4 + 7 + 10 + 13 = 34$$

Example – arithmetic series

For a set of numbers spaced by a constant amount,
an “arithmetic progression or sequence”

i.e., where the n th number is $a_n = a_1 + (n-1)d$

e.g., the numbers 4, 7, 10, and 13

i.e., $a_1 = 4$, $a_2 = 7$, $a_3 = 10$ and $a_4 = 13$

with, therefore $d = 3$ and $m = 4$ terms in the
progression

gives the “series” (sum of the terms)

$$a_1 + a_2 + a_3 + a_4 \equiv \sum_{j=1}^4 a_j = \sum_{j=1}^m [a_1 + (j-1)d]$$

Example – arithmetic series

For a set of numbers spaced by a constant amount,
an “arithmetic progression or sequence”

i.e., where the n th number is $a_n = a_1 + (n-1)d$

e.g., the numbers 4, 7, 10, and 13

i.e., $a_1 = 4$, $a_2 = 7$, $a_3 = 10$ and $a_4 = 13$

with, therefore $d = 3$ and $m = 4$ terms in the
progression

gives the “series” (sum of the terms)

$$a_1 + a_2 + a_3 + a_4 = \sum_{j=1}^m \left[a_1 + (j-1)d \right] = m \frac{(a_1 + a_4)}{2}$$

Example – arithmetic series

For a set of numbers spaced by a constant amount,
an “arithmetic progression or sequence”

i.e., where the n th number is $a_n = a_1 + (n-1)d$

e.g., the numbers 4, 7, 10, and 13

i.e., $a_1 = 4$, $a_2 = 7$, $a_3 = 10$ and $a_4 = 13$

with, therefore $d = 3$ and $m = 4$ terms in the
progression

gives the “series” (sum of the terms)

$$a_1 + a_2 + a_3 + a_4 = \sum_{j=1}^m \left[a_1 + (j-1)d \right] = m \frac{(a_1 + a_m)}{2} = 4 \times \frac{(4+13)}{2} = 34$$

Example – geometric series

With a constant ratio between successive terms,

a “geometric progression or sequence”

i.e., where the n th term is $a_n = a_1 r^{n-1}$

e.g., the numbers 3, 6, 12, and 24

i.e., $a_1 = 3$, $a_2 = 6$, $a_3 = 12$ and $a_4 = 24$

with, therefore $r = 2$ and $m = 4$ terms in the progression

gives the “series” (sum of the terms)

$$a_1 + a_2 + a_3 + a_4 = 3 + 6 + 12 + 24 = 45$$

Example – geometric series

With a constant ratio between successive terms,

a “geometric progression or sequence”

i.e., where the n th term is $a_n = a_1 r^{n-1}$

e.g., the numbers 3, 6, 12, and 24

i.e., $a_1 = 3$, $a_2 = 6$, $a_3 = 12$ and $a_4 = 24$

with, therefore $r = 2$ and $m = 4$ terms in the progression

gives the “series” (sum of the terms)

$$a_1 + a_2 + a_3 + a_4 \equiv \sum_{j=1}^4 a_j = \sum_{j=1}^m a_1 r^{(j-1)}$$

Example – geometric series

With a constant ratio between successive terms,

a “geometric progression or sequence”

i.e., where the n th term is $a_n = a_1 r^{n-1}$

e.g., the numbers 3, 6, 12, and 24

i.e., $a_1 = 3$, $a_2 = 6$, $a_3 = 12$ and $a_4 = 24$

with, therefore $r = 2$ and $m = 4$ terms in the progression

gives the “series” (sum of the terms)

$$a_1 + a_2 + a_3 + a_4 \equiv \sum_{j=1}^4 a_j = \sum_{j=1}^m a_1 r^{(j-1)} = a_1 \frac{1 - r^m}{1 - r}$$

Example – geometric series

With a constant ratio between successive terms,

a “geometric progression or sequence”

i.e., where the n th term is $a_n = a_1 r^{n-1}$

e.g., the numbers 3, 6, 12, and 24

i.e., $a_1 = 3$, $a_2 = 6$, $a_3 = 12$ and $a_4 = 24$

with, therefore $r = 2$ and $m = 4$ terms in the progression

gives the “series” (sum of the terms)

$$a_1 + a_2 + a_3 + a_4 \equiv \sum_{j=1}^4 a_j = \sum_{j=1}^m a_1 r^{(j-1)} = a_1 \frac{1-r^m}{1-r} = 3 \times \frac{1-16}{1-2} = 3 \times 15 = 45$$

Summation over multiple indexes

We can extend the summation notation

Suppose we have two lists of numbers

a_1, a_2, a_3, a_4 and b_1, b_2, b_3

and we want to add up all the products

$$\begin{aligned} R = & a_1b_1 + a_1b_2 + a_1b_3 \\ & + a_2b_1 + a_2b_2 + a_2b_3 \\ & + a_3b_1 + a_3b_2 + a_3b_3 \\ & + a_4b_1 + a_4b_2 + a_4b_3 \end{aligned}$$

Summation over multiple indexes

Then we can write

$$\begin{aligned} R = & a_1b_1 + a_1b_2 + a_1b_3 \\ & + a_2b_1 + a_2b_2 + a_2b_3 \\ & + a_3b_1 + a_3b_2 + a_3b_3 \\ & + a_4b_1 + a_4b_2 + a_4b_3 \end{aligned} = \sum_{j=1}^4 \left(\sum_{k=1}^3 a_j b_k \right)$$

and, because order of addition does not matter
in various equivalent notations

$$R = \sum_{j=1}^4 \sum_{k=1}^3 a_j b_k = \sum_{k=1}^3 \sum_{j=1}^4 a_j b_k \equiv \sum_{j,k} a_j b_k \equiv \sum_{j,k} a_j b_k$$

Factorial notation

Quite often, we need a convenient way of writing the product of successive integers

e.g., $1 \times 2 \times 3 \times 4$

We can write this as

$$1 \times 2 \times 3 \times 4 \equiv 4!$$

called “four factorial”

and using the “exclamation point” “!”

The notation is obvious for most other cases

Note, though, that we choose

$$0! = 1$$

Product notation

Generally, when we want to write the product of various successive terms

$$a_1 \times a_2 \times a_3 \times a_4$$

by analogy with the summation notation
we can use the "product" notation

$$a_1 \times a_2 \times a_3 \times a_4 \equiv \prod_{j=1}^4 a_j$$

For example, for all integers $n \geq 1$

$$n! \equiv \prod_{p=1}^n p$$





Sums, products and power series



Power series

Background mathematics review

David Miller

Analytic functions and power series

For a very broad class of the functions in physics

we presume they are “analytic”

(except possibly at some “singularities”)

i.e., at least for some range of values of the argument x near some point x_o

the function $f(x)$ can be arbitrarily well approximated by a “power series”

i.e., with $x_o = 0$ for simplicity we have

$$f(x) \equiv a_o + a_1x + a_2x^2 + a_3x^3 \cdots$$

which may be an infinite series

*The “ellipsis”
“...” means we
omit writing
some terms
explicitly*

Analytic functions and power series

For a very broad class of the functions in physics

we presume they are “analytic”

(except possibly at some “singularities”)

i.e., at least for some range of values of the argument x near some point x_o

the function $f(x)$ can be arbitrarily well approximated by a “power series”

i.e., generally

$$f(x) \equiv a_o + a_1(x - x_o) + a_2(x - x_o)^2 + a_3(x - x_o)^3 \cdots$$

Maclaurin series

Taking the simplest case of $x_o = 0$ first

$$f(x) \equiv a_o + a_1x + a_2x^2 + a_3x^3 \dots$$

Obviously at $x = 0$

$$f(0) = a_o \text{ so, trivially, } a_o = f(0)$$

Now,

$$f'(x) \equiv \frac{df}{dx} = a_1 + 2a_2x + 3a_3x^2 \dots$$

so

$$f'(0) \equiv \left. \frac{df}{dx} \right|_0 = a_1$$

Maclaurin series

Continuing

$$f''(x) \equiv \frac{d^2 f}{dx^2} = 2a_2 + 3 \times 2 \times a_3 x \cdots = 2! \times a_2 + 3! \times a_3 x \cdots$$

so

$$f''(0) \equiv \left. \frac{d^2 f}{dx^2} \right|_0 = 2! a_2 \cdots$$

and

$$f'''(0) \equiv \left. \frac{d^3 f}{dx^3} \right|_0 = 3! a_3 \cdots$$

and so on

Maclaurin and Taylor series

Continuing gives the Maclaurin series

$$f(x) = f(0) + \frac{x}{1!} \frac{df}{dx} \Big|_0 + \frac{x^2}{2!} \frac{d^2 f}{dx^2} \Big|_0 + \dots + \frac{x^n}{n!} \frac{d^n f}{dx^n} \Big|_0 + \dots$$

Repeating the same procedure around $x = x_o$ with

$$f(x) \equiv a_o + a_1(x - x_o) + a_2(x - x_o)^2 + a_3(x - x_o)^3 \dots$$

gives the Taylor series

$$f(x) = f(x_o) + \frac{(x - x_o)}{1!} \frac{df}{dx} \Big|_{x_o} + \frac{(x - x_o)^2}{2!} \frac{d^2 f}{dx^2} \Big|_{x_o} + \dots + \frac{(x - x_o)^n}{n!} \frac{d^n f}{dx^n} \Big|_{x_o} + \dots$$

Example power series expansions

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

$$\exp(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

Power series for approximations

Maclaurin and Taylor series allow approximations for small ranges of the argument about a point

Examples for small x

Approximations to “first order in x ”

$$1/(1+x) \simeq 1-x \quad \exp(x) \simeq 1+x \quad \sin x \simeq x$$

$$\sqrt{1+x} \simeq 1+x/2 \quad \ln(1+x) \simeq x \quad \tan x \simeq x$$

Note lowest order dependence on x for $\cos x$ is “second order”

$$\cos x \simeq 1-x^2/2$$

