


9.3 Applying time-dependent perturbation theory

Slides: Video 9.3.3 Refractive index

Text reference: Quantum Mechanics
for Scientists and Engineers

Section 7.3





Applying time-dependent perturbation
theory



Refractive index

Quantum mechanics for scientists and engineers

David Miller

Refractive index

In classical electromagnetism

when we apply an electric field E to a material

we can imagine that the field pulls the average position of the negatively charged electrons

away from the average position of the positively charged nuclei of the atoms

creating what is called a

“polarization” P

Refractive index

At least for applied fields that are small compared to the fields in the atoms

we would expect that this "polarization" P
is proportional to the applied field E
with a proportionality constant
the "susceptibility" χ

For historical reasons

a fundamental constant is also involved
the "electric constant" ϵ_o
so we write $P = \epsilon_o \chi E$

Refractive index

In a classical view of electromagnetic waves
the fact that we are moving these electrons
with the oscillating field
means that we have a new oscillating
current
which itself generates new waves

These new waves interfere with the original waves
with the net result that the effective (phase)
velocity of the total wave is changed
formally, divided by the "refractive index" n_r

Refractive index

If we follow through the classical
electromagnetism of this problem

for example, using Maxwells' equations

we can conclude

at least for non-absorbing materials

that the refractive index is given by

$$n_r = \sqrt{1 + \chi}$$

Hence, if we can calculate the proportionality
between P and E

we can calculate n_r

Calculation of refractive index

Consider a system with a single electron
or in which our interactions are only with a single
electron

Classically the dipole moment μ_{dip}
associated with moving a single electron
through a distance z is
by definition

$$\mu_{dip} = -ez$$

where the minus sign is because the electron
charge is negative

Calculation of refractive index

The polarization P is formally

the dipole moment per unit volume

and so the quantum mechanical expectation value of the polarization is

$$\langle P \rangle = \frac{-e \langle z \rangle}{V}$$

where V is the volume of the system

Our quantum mechanical task of calculating refractive index

reduces essentially to calculating $\langle P \rangle$

Calculation of refractive index

Since we are working in first-order perturbation theory
we write the (approximate) total state of the system

$$|\Psi\rangle = |\Phi^{(0)}\rangle + |\Phi^{(1)}\rangle$$

Note we use the full time-dependent state vectors (kets)

$|\Phi^{(0)}\rangle$ is the unperturbed state vector

$|\Phi^{(1)}\rangle$ is the first-order (time-dependent) correction

where
$$|\Phi^{(1)}\rangle = \sum_n a_n^{(1)}(t) \exp(-i\omega_n t) |\psi_n\rangle$$

with unperturbed energy eigenvalues E_n

and eigenfunctions $|\psi_n\rangle$

Calculation of refractive index

With the state vector $|\Psi\rangle = |\Phi^{(0)}\rangle + |\Phi^{(1)}\rangle$

the expectation value of the polarization would be

$$\begin{aligned}\langle P \rangle &= -\frac{e}{V} \langle \Psi | z | \Psi \rangle \\ &= -\frac{e}{V} \left[\langle \Phi^{(0)} | z | \Phi^{(0)} \rangle + \langle \Phi^{(1)} | z | \Phi^{(0)} \rangle + \langle \Phi^{(0)} | z | \Phi^{(1)} \rangle + \langle \Phi^{(1)} | z | \Phi^{(1)} \rangle \right]\end{aligned}$$

The first term $-e \langle \Phi^{(0)} | z | \Phi^{(0)} \rangle$

is just the static dipole moment of the material in its unperturbed state

so we will not consider it further

Calculation of refractive index

$$\langle P \rangle = -\frac{e}{V} \langle \Psi | z | \Psi \rangle$$

$$= -\frac{e}{V} \left[\langle \Phi^{(0)} | z | \Phi^{(0)} \rangle + \langle \Phi^{(1)} | z | \Phi^{(0)} \rangle + \langle \Phi^{(0)} | z | \Phi^{(1)} \rangle + \langle \Phi^{(1)} | z | \Phi^{(1)} \rangle \right]$$

The fourth term $-e \langle \Phi^{(1)} | z | \Phi^{(1)} \rangle$

is second order in the perturbation

and hence, in this first order calculation

we drop it also

Calculation of refractive index

So we are left with

$$\langle P \rangle = -\frac{e}{V} \left[\langle \Phi^{(1)} | z | \Phi^{(0)} \rangle + \langle \Phi^{(0)} | z | \Phi^{(1)} \rangle \right]$$

But because z is a Hermitian operator

$$\langle \Phi^{(1)} | z | \Phi^{(0)} \rangle = \langle \Phi^{(0)} | z | \Phi^{(1)} \rangle^*$$

therefore

$$\langle P \rangle = -\frac{2e}{V} \operatorname{Re} \left[\langle \Phi^{(0)} | z | \Phi^{(1)} \rangle \right]$$

Calculation of refractive index

For the sake of definiteness

we presume we start in eigenstate m , i.e.,

$$|\Phi^{(0)}\rangle = \exp(-i\omega_m t) |\psi_m\rangle$$

Hence, using the first-order perturbation expansion

$$|\Phi^{(1)}\rangle = \sum_n a_n^{(1)}(t) \exp(-i\omega_n t) |\psi_n\rangle$$

then, from $\langle P \rangle = -\frac{2e}{V} \operatorname{Re} \left[\langle \Phi^{(0)} | z | \Phi^{(1)} \rangle \right]$

we have $\langle P \rangle = -\frac{2e}{V} \operatorname{Re} \left[\sum_n a_n^{(1)}(t) \exp(i\omega_{mn} t) \langle \psi_m | z | \psi_n \rangle \right]$

Calculation of refractive index

We are interested here in the steady-state situation
with a continuous oscillating field

so we take $\hat{H}_p(t) = eE(t)z = \hat{H}_{po} [\exp(-i\omega t) + \exp(i\omega t)]$
as valid for all times

We can rewrite $\dot{a}_q^{(1)}(t) = \frac{1}{i\hbar} \exp(i\omega_{qm}t) \langle \psi_q | \hat{H}_p(t) | \psi_m \rangle$ as

$$\dot{a}_q^{(1)}(t) = \frac{eE_o}{i\hbar} \langle \psi_q | z | \psi_m \rangle \exp(i\omega_{qm}t) [\exp(-i\omega t) + \exp(i\omega t)]$$

to obtain

$$a_q^{(1)}(t) = -\frac{eE_o}{\hbar} \langle \psi_q | z | \psi_m \rangle \left[\frac{\exp[i(\omega_{qm} - \omega)t]}{(\omega_{qm} - \omega)} + \frac{\exp[i(\omega_{qm} + \omega)t]}{(\omega_{qm} + \omega)} \right]$$

Calculation of refractive index

Substituting

$$a_q^{(1)}(t) = -\frac{eE_o}{\hbar} \langle \psi_q | z | \psi_m \rangle \left[\frac{\exp[i(\omega_{qm} - \omega)t]}{(\omega_{qm} - \omega)} + \frac{\exp[i(\omega_{qm} + \omega)t]}{(\omega_{qm} + \omega)} \right]$$

into

$$\langle P \rangle = -\frac{2e}{V} \operatorname{Re} \left[\sum_n a_n^{(1)}(t) \exp(i\omega_{mn}t) \langle \psi_m | z | \psi_n \rangle \right]$$

gives

$$\begin{aligned} \langle P \rangle &= \frac{2e^2 E_o}{\hbar V} \operatorname{Re} \sum_n \left| \langle \psi_m | z | \psi_n \rangle \right|^2 \exp(i\omega_{mn}t) \\ &\times \left[\frac{\exp[i(\omega_{nm} - \omega)t]}{(\omega_{nm} - \omega)} + \frac{\exp[i(\omega_{nm} + \omega)t]}{(\omega_{nm} + \omega)} \right] \end{aligned}$$

Calculation of refractive index

Noting $\omega_{mn} = -\omega_{nm}$ and taking the real part gives

$$\begin{aligned}\langle P \rangle &= \frac{2e^2 E_o}{\hbar V} \sum_n \left| \langle \psi_m | z | \psi_n \rangle \right|^2 \left[\frac{\cos(-\omega t)}{(\omega_{nm} - \omega)} + \frac{\cos(\omega t)}{(\omega_{nm} + \omega)} \right] \\ &= \frac{2e^2 E_o \cos(\omega t)}{\hbar V} \sum_n \left| \langle \psi_m | z | \psi_n \rangle \right|^2 \left[\frac{1}{(\omega_{nm} - \omega)} + \frac{1}{(\omega_{nm} + \omega)} \right]\end{aligned}$$

and so we have, from $P = \epsilon_o \chi E$

$$\chi = \frac{e^2}{\epsilon_o \hbar V} \sum_n \left| \langle \psi_m | z | \psi_n \rangle \right|^2 \left[\frac{1}{(\omega_{nm} - \omega)} + \frac{1}{(\omega_{nm} + \omega)} \right]$$

Calculation of refractive index

From
$$\chi = \frac{e^2}{\epsilon_o \hbar V} \sum_n \left| \langle \psi_m | z | \psi_n \rangle \right|^2 \left[\frac{1}{(\omega_{nm} - \omega)} + \frac{1}{(\omega_{nm} + \omega)} \right]$$

and using $n_r = \sqrt{1 + \chi}$

we can now deduce the refractive index n_r

completing our calculation of refractive index

Hence, knowing the unperturbed energy eigenfunctions and eigenvalues for some quantum mechanical system

we can calculate the refractive index

using time-dependent perturbation theory

Absorption and refractive index

Note a key difference between

absorption and
refractive index

For absorption

the frequency ω must match the transition
frequency ω_{nm} very closely

for that particular transition to give rise
to absorption of photons

Absorption and refractive index

For the refractive index

the contribution of a particular possible
transition $|\psi_m\rangle \rightarrow |\psi_n\rangle$

to the susceptibility

and hence to the refractive index

is finite

even when the frequencies do not
match exactly or even closely

That contribution to the susceptibility rises
steadily as ω rises towards ω_{nm}

Absorption and refractive index

Note that

if we have an absorbing transition at some
frequency ω_{nm}
it contributes to refractive index at all
frequencies

Refractive index (in a region where the material is
transparent)

arises entirely because of the possibility of
absorption at other frequencies

If there is a refractive index different from unity
then there must be absorption possible at
some other frequency or frequencies

Absorption and refractive index

The relation between refractive index and absorption is known from classical physics

through the Kramers-Kronig relations

The derivation of those relations is entirely mathematical

shedding no light on

the physical mechanism whereby
absorption and refractive index are
related

Absorption and refractive index

With our quantum mechanical expressions
we can understand any particular aspect in
the relation between the two processes

In the quantum mechanical picture, we find
even though we are in a transparent
spectral region
there are finite occupation probabilities
for all of the states of the system
which are essential if the material has
a polarization

Absorption and refractive index

The polarization arises because the charges in the material

change their physical wavefunctions in response to the field

mixing in other states of the system in response to the perturbation

If we examined the expectation value of the energy of the material

we would also find quite real energy stored in the material as a result

even though the material is transparent

