

Introductory Statistics for the Life and Biomedical Sciences

Derivative of
OpenIntro Statistics
Third Edition

Original Authors

David M Diez
Christopher D Barr
Mine Çetinkaya-Rundel

Contributing Authors

David Harrington
[Briefly Describe Contribution]

Julie Vu
[Briefly Describe Contribution]

Alice Zhao
[Briefly Describe Contribution]

Copyright © 2015. Third Edition.

This textbook is available under a Creative Commons license. Visit openintro.org for a free PDF, to download the textbook's source files, or for more information about the license.

Contents

1	Introduction to data	8
1.1	Case study	9
1.2	Data basics	12
1.3	Data collection principles	17
1.4	Examining numerical data	28
1.5	Considering categorical data	40
1.6	Exercises	47

Preface

This book provides an introduction to statistics and its applications in the life sciences, and biomedical research. It is based on the freely available *OpenIntro Statistics, Third Edition*, and, like *OpenIntro* it may be downloaded as a free PDF at **Need location**. The text adds substantial new material, revises or eliminates sections from *OpenIntro*, and re-uses some material directly. Readers need not have read *OpenIntro*, since this book is intended to be used independently. We have retained some of the exercises from *OpenIntro* that may not come directly from medicine or the life sciences but illustrate important ideas or methods that are commonly used in fields such as biology.

Introduction to Statistics for the Life and Biomedical Sciences is intended for graduate and undergraduate students interested in careers in biology or medicine, and may also be profitably read by students of public health. It covers many of the traditional introductory topics in statistics used in those fields, but also adds some newer methods being used in molecular biology. Statistics has become an integral part of research in medicine and biology, and the tools for displaying, summarizing and drawing inferences from data are essential both for understanding the outcomes of studies and for incorporating measures of uncertainty into that understanding. An introductory text in statistics for students considering careers in medicine, public health or the life sciences should be more than the usual introduction with more examples from biology or medical science. Along with the value of careful, robust analyses of experimental and observational data, it should convey some of the excitement of discovery that emerges from the interplay of science with data collection and analysis. We hope we have conveyed some of that excitement here.

We have tried to balance the sometimes competing demands of mastering the important technical aspects of methods of analysis with gaining an understanding of important concepts. The examples and exercises include opportunities for students to build skills in conducting data analyses and to state conclusions with clear, direct language that is specific to the context of a problem. We also believe that computing is an essential part of statistics, just as mathematics was when computing was more difficult or expensive. The text includes many examples where software is used to aid in the understanding of the features of a data as well as exercises where computing is used to help illustrate the notions of randomness and variability. Because they are freely available, we use the R statistical language with the *R Studio* interface. Information on downloading R and *R Studio* is may be found in the Labs at openintro.org. Nearly all examples and exercises can be adapted to either SAS, Stata or other software, but we have not done that.

Textbook overview

The chapters of this book are as follows:

- 1. Introduction to data.** Data structures, variables, summaries, graphics, and basic data collection techniques.
- 2. Probability (special topic).** The basic principles of probability. An understanding of this chapter is not required for the main content in Chapters ??-??.
- 3. Distributions of random variables.** Introduction to the normal model and other key distributions.
- 4. Foundations for inference.** General ideas for statistical inference in the context of estimating the population mean.
- 5. Inference for numerical data.** Inference for one or two sample means using the normal model and t distribution, and also comparisons of many means using ANOVA.
- 6. Inference for categorical data.** Inference for proportions using the normal and chi-square distributions, as well as simulation and randomization techniques.
- 7. Introduction to linear regression.** An introduction to regression with two variables. Most of this chapter could be covered after Chapter 1.

8. Multiple and logistic regression. An introduction to multiple regression and logistic regression for an accelerated course.

The remainder of this section requires revision

OpenIntro Statistics was written to allow flexibility in choosing and ordering course topics. The material is divided into two pieces: main text and special topics. The main text has been structured to bring statistical inference and modeling closer to the front of a course. Special topics, labeled in the table of contents and in section titles, may be added to a course as they arise naturally in the curriculum.

Examples, exercises, and appendices

Examples and within-chapter exercises throughout the textbook may be identified by their distinctive bullets:

- **Example 0.1** Large filled bullets signal the start of an example.

Full solutions to examples are provided and often include an accompanying table or figure.

- **Guided Practice 0.2** Large empty bullets signal to readers that an exercise has been inserted into the text for additional practice and guidance. Students may find it useful to fill in the bullet after understanding or successfully completing the exercise. Solutions are provided for all within-chapter exercises in footnotes.¹

There are exercises at the end of each chapter that are useful for practice or homework assignments. Many of these questions have multiple parts, and odd-numbered questions include solutions in Appendix ??.

Probability tables for the normal, t , and chi-square distributions are in Appendix ??, and PDF copies of these tables are also available from **openintro.org** for anyone to download, print, share, or modify.

¹Full solutions are located down here in the footnote!

OpenIntro, online resources, and getting involved

OpenIntro is an organization focused on developing free and affordable education materials. *OpenIntro Statistics*, our first project, is intended for introductory statistics courses at the high school through university levels.

We encourage anyone learning or teaching statistics to visit **openintro.org** and get involved. We also provide many free online resources, including free course software. Data sets for this textbook are available on the website and through a companion R package.² All of these resources are free, and we want to be clear that anyone is welcome to use these online tools and resources with or without this textbook as a companion.

We value your feedback. If there is a particular component of the project you especially like or think needs improvement, we want to hear from you. You may find our contact information on the title page of this book or on the [About](#) section of **openintro.org**.

Acknowledgements

This project would not be possible without the dedication and volunteer hours of all those involved. No one has received any monetary compensation from this project, and we hope you will join us in extending a *thank you* to all those volunteers below.

The authors would like to thank Andrew Bray, Meenal Patel, Yongtao Guan, Filipp Brunshteyn, Rob Gould, and Chris Pope for their involvement and contributions. We are also very grateful to Dalene Stangl, Dave Harrington, Jan de Leeuw, Kevin Rader, and Philippe Rigollet for providing us with valuable feedback.

²Diez DM, Barr CD, Çetinkaya-Rundel M. 2012. `openintro`: OpenIntro data sets and supplement functions. <http://cran.r-project.org/web/packages/openintro>.

Chapter 1

Introduction to data

Scientists seek to answer questions using rigorous methods and careful observations. These observations – collected from the likes of field notes, surveys, and experiments – form the backbone of a statistical investigation and are called **data**. Statistics is the study of how to best collect, analyze, and draw conclusions from data. It is helpful to place statistics in the context of a general process of investigation:

1. Identify a question or problem.
2. Collect relevant data on the topic.
3. Analyze the data.
4. Form a conclusion.

Statistics as a subject focuses on making stages 2-4 objective, rigorous, and efficient. That is, statistics has three primary components: How can data best be collected? How should it be analyzed? What can be inferred from the analysis?

This chapter provides a brief introduction to the basic principles of data collection techniques and analytic tools, and illustrates the important role statistics plays in medicine and biology.

JV: Make reference to how this chapter can be used by either someone new to statistics or someone who has had the material before?

JV: Something to address later – where in the formatting does it specify that the first paragraph in a section does not get indented, but the others do...

1.1 Case study: preventing peanut allergies

Section 1.1 introduces an important problem in medicine: evaluating the effect of an intervention. *JV: Intervention should be defined, could work in a dependent clause.* Terms in this section, and indeed much of this chapter, will all be revisited later in more detail.

The proportion of young children in Western countries with peanut allergies has doubled in the last 10 years. Previous research suggests that exposing infants to peanut-based foods, rather than excluding such foods from their diets, may be an effective strategy for preventing the development of peanut allergies. This section describes an experiment (a clinical trial, in the terminology of medical research) designed to address the following research question: Does early exposure to peanut products reduce the probability that a child will develop peanut allergies?

The "Learning Early about Peanut Allergy" (LEAP) study was reported in the New England Journal of Medicine in 2015.¹ The study team enrolled children in the United Kingdom between 2006 and 2009, selecting 640 infants with eczema, egg allergy, or both. Each child was randomly assigned to the treatment group (peanut consumption) or the control group (peanut avoidance); children in the treatment group were fed at least 6 grams of peanut protein until 5 years of age, while children in the control group were to avoid consuming peanut protein until 5 years of age. *In this study, the control group provides a reference point for estimating the effect of peanut exposure in the treatment group.* *JV: This last sentence comes off as a bit technical/vague, perhaps due to "estimating the effect" – could be omitted, I think.*

At age 5, each child was tested for peanut allergy using an oral food challenge (OFC): 5 grams of peanut protein in a single dose. Children had been previously been tested for peanut allergy through a skin test, conducted at the time of study entry; the main analysis presented in the paper was based on the 530 children with a negative skin test

¹Du Toit, George, et al. Randomized trial of peanut consumption in infants at risk for peanut allergy. New England Journal of Medicine 372.9 (2015): 803-813.

result. Of these children, 263 were assigned to "Peanut Avoidance" and 267 to "Peanut Consumption." The outcome at 5 years was reported as either "Fail OFC" (allergic reaction) or "Pass OFC" (no allergic reaction).

Table 1.1 shows the participant study ID number, treatment assignment, and OFC outcome for 5 children. All five of these children passed the food challenge.

JV: The last two row labels should be corrected to 529 and 530; not sure how to fix that in the R code. We should be careful later on to clarify that in the file, the row.names column corresponds to the dataset with all 640 children. Alternatively, that column could be omitted and a separate dataset made with the children with a positive skin test, which could be analyzed in an end-of-chapter exercise.

JV: Just curious...so 267 for the peanut consumption group includes the one child with a positive skin test?

	participant.ID	treatment.group	overall.V60.outcome
1	LEAP_100522	Peanut Consumption	PASS OFC
2	LEAP_103358	Peanut Consumption	PASS OFC
3	LEAP_105069	Peanut Avoidance	PASS OFC
	⋮	⋮	⋮
639	LEAP_994047	Peanut Avoidance	PASS OFC
640	LEAP_997608	Peanut Consumption	PASS OFC

Table 1.1: Results for five children from the peanut study.

Summary tables are generally more helpful than individual participant listings when looking for patterns in data. Table 1.2 shows outcomes grouped by treatment group and the result of the OFC test. From this table, it is possible to compute some simple summary statistics.

	FAIL OFC	PASS OFC	Sum
Peanut Avoidance	36	227	263
Peanut Consumption	5	262	267
Sum	41	489	530

Table 1.2: LEAP Study Results

A **summary statistic** is a single number summarizing a large amount of data.² In

²Formally, a summary statistic is a value computed from the data. Some summary statistics are more useful than others.

the Peanut Avoidance group, the proportion of participants failing the food challenge at 5 years of age is $36/263 = 0.137$ (13.7%); in the Peanut Consumption intervention, the proportion failing is $5/267 = 0.019$ (1.9%). The difference between these two proportions, 11.8%, is a single summary statistic describing the extent to which these two proportions differ. A second summary statistic, the ratio of the two proportions, $0.137/0.019 = 7.31$, indicates that the proportion failing in the Avoidance group is more than 7 times that of the Consumption group. This ratio is called a **relative risk**.

JV: Interpretation of RR should be more clearly stated here, or else RR should be omitted entirely.

The summary statistics for the LEAP study highlight an important point – the results of a study can sometimes be surprising. A parent of a child already known to be allergic to eggs might be justifiably skeptical about feeding peanut butter to their child. The LEAP study suggests that, at least for children similar to those enrolled in the study, the benefits of early exposure might be substantial.

There are important aspects of the study to be cautious about. This study was conducted in the United Kingdom at a single site of pediatric care; it is not at all clear that results in children from that site can be generalized to other countries or cultures. Furthermore, the results also raise an important statistical issue: does the study provide definitive evidence that peanut consumption is beneficial? In other words, is the 11.8% difference between the two groups larger than one would expect by chance variation alone?

Suppose a coin is flipped 100 times. While the chance a coin lands heads in any given coin flip is 50%, observing exactly 50 heads is unlikely; instead, the coin may land heads 43 times, 51 times, 59 times, etc. This type of fluctuation is part of almost any experiment or study. It may well be possible that the 11.8% difference in the peanut allergy study is only due to this natural variation, and that the two interventions are actually equally effective. However, the larger the difference observed (for a particular study size), the less credible it is that the difference is due to chance alone. If out of 100 flips, a coin landed heads only 5 times, it would be reasonable to doubt that the outcome was due to chance; perhaps the coin is weighted so that tails are more likely to occur.

For the LEAP study, the 11.8% difference is indeed larger than that expected by

chance alone, suggesting that peanut consumption is the more effective intervention for preventing subsequent allergies. The material on hypothesis testing in later chapters will provide the statistical tools to examine this issue.

1.2 Data basics

Effective presentation and description of data is a first step in most analyses. This section introduces one structure for organizing data as well as some terminology that will be used throughout this book.

1.2.1 Observations, variables, and data matrices

This section describes data used in a study published in the *Journal of Evolutionary Biology* about maternal investment at differing altitudes, conducted in a frog species endemic to the Tibetan Plateau (*Rana kukunoris*).³ Reproduction is a costly process for females, necessitating a trade-off between individual egg size and total number of eggs produced. Researchers collected measurements on egg clutches found at breeding ponds across 11 study sites; for 5 sites, they also collected data on individual female frogs.

	altitude	latitude	egg.size	clutch.size	clutch.volume	body.size
1	3,462.00	34.82	1.95	181.97	177.83	3.63
2	3,462.00	34.82	1.95	269.15	257.04	3.63
3	3,462.00	34.82	1.95	158.49	151.36	3.72
150	2,597.00	34.05	2.24	537.03	776.25	NA

Table 1.3: Frog Study Data Matrix

Table 1.3 displays rows 1, 2, 3, and 150 of the data from the 431 clutches. The complete set of observations will be referred to as the frog dataset. Each row in the table corresponds to a single clutch, indicating where the clutch was collected (`altitude` and `latitude`), `egg.size`, `clutch.size`, `clutch.volume`, and `body.size` of the mother when available. "NA" corresponds to a missing value; information on individual females was not collected for that particular site. The columns represent characteristics, called **variables**,

³ Chen, W., et al. Maternal investment increases with altitude in a frog on the Tibetan Plateau. *Journal of evolutionary biology* 26.12 (2013): 2710-2715.

variable	description
altitude	Altitude of the study site in meters above sea level
latitude	Latitude of the study site measured in degrees
egg.size	Average diameter of an individual egg to the 0.01 mm
clutch.size	Estimated number of eggs in clutch
clutch.volume	Volume of egg clutch in mm ³
body.size	Length of mother frog in cm

Table 1.4: Variables and their descriptions for the frog dataset.

for each clutch.

For example, the first row represents a clutch collected at altitude 3,462 meters above sea level, latitude 34.82 degrees; the clutch contained an estimated 182 eggs, with individual eggs averaging 1.95 mm in diameter, for a total volume of 177.8 mm³. The eggs were laid by a female measuring 3.63 cm long. It is important to understand the definitions of variables, as they are not always obvious. For example, why has `clutch.size` not been recorded as whole numbers? This has to do with how the observations were collected. In a given clutch, researchers counted approximately 5 grams' worth of eggs and then estimated the total number of eggs based on the mass of the entire clutch. Definitions of the variables are given in Table 1.4.⁴

The data in Table 1.3 are organized as a **data matrix**. Each row of a data matrix corresponds to a unique observational unit, and each column corresponds to a variable. A data matrix for the LEAP study introduced in Section 1.1 is shown in Table 1.1 on page 10, in which the cases were patients and three variables were recorded for each patient. Data matrices are a convenient way to record and store data. If the data are collected for another individual, another row can easily be added; similarly, another column can be added for a new variable.

1.2.2 Types of variables

The Functional polymorphisms Associated with Human Muscle Size and Strength study (FAMuSS), funded by the National Institutes of Health (NIH), measured a variety of demographic, phenotypic, and genetic characteristics for about 1,300 participants.⁵ Data

⁴The data discussed here are in the original scale; in the published paper, values have been log-transformed.

⁵Thompson PD, Moyna M, Seip, R, et al., 2004. Functional Polymorphisms Associated with Human Muscle Size and Strength. *Medicine and Science in Sports and Exercise* 36:1132 - 1139

from the study has been used in many subsequent studies⁶, such as one examining the relationship between muscle strength and genotype at a location on the ACTN3 gene.⁷ Four rows of the famuss dataset are shown in Table 1.5, and the variables are summarized in Table 1.6.⁸

	sex	age	race	height	weight	actn3.r577x	ndrm.ch
1	Female	27	Caucasian	65.0	199.0	CC	40.0
2	Male	36	Caucasian	71.7	189.0	CT	25.0
3	Female	24	Caucasian	65.0	134.0	CT	40.0
	⋮	⋮	⋮	⋮	⋮	⋮	
595	Female	30	Caucasian	64.0	134.0	CC	43.8

Table 1.5: Four rows from the famuss data matrix.

The variables `age`, `height`, `weight`, and `ndrm.ch` are **numerical** variables. They can take on a wide range of numerical values, and it is possible to add, subtract, or take averages with these values. On the other hand, a variable reporting telephone numbers would not be classified as numerical, since averages, sums, and differences in this context would have no meaning. Age measured in years is said to be **discrete**, since it can only take numerical values with jumps. On the other hand, percent change in strength in the non-dominant arm (`ndrm.ch`) is said to be **continuous**.

The variables `sex`, `race`, and `actn3.r577x` are **categorical** variables, and the possible values are called the variable's **levels**.⁹ For example, the levels of `actn3.r577x` are the three possible genotypes at this particular locus: CC, CT, or TT. Categorical variables with levels that have a natural ordering can be more specifically referred to as **ordered categorical** variables. There are no ordered categorical variables in the famuss data, but it would be easy to create one; age of the participants grouped into 5-year intervals (15-20, 21-25, 26-30, etc.) would be an ordered categorical variable. Statistical software such as R calls categorical variables **factors**, and the possible values of factors are called **levels**.

In the frog data, the variables `egg.size`, `clutch.size`, `clutch.volume`, and `body.size` are all continuous variables. *DH: JV, agree about latitude? JV: I don't think either altitude or*

⁶Pescatello L, et al. Highlights from the functional single nucleotide polymorphisms associated with human muscle size and strength or FAMuSS study, BioMed Research International 2013.

⁷Clarkson P, et al., Journal of Applied Physiology 99: 154-163, 2005.

⁸Data freely available at <http://people.umass.edu/foulkes/asg/data.html>

⁹Categorical variables are sometimes called **nominal** variables.

variable	description
sex	Sex of the participant
age	Age in years
race	Recorded as African Am (African American), Caucasian, Asian, Hispanic and Other
height	Height in inches
weight	Weight in lbs
actn3.r577x	Genotype at the location r577x in the ACTN3 gene.
ndrm.ch	Percent change in strength in the non-dominant arm, comparing strength after to before training

Table 1.6: Variables and their descriptions for the famuss data set.




Figure 1.7: Breakdown of variables into their respective types.

latitude qualifies, because they refer to 11 specific locations; both function more like categorical variables in this context.

● **Example 1.1** Suppose a research assistant collected data on the first 20 individuals to visit one of the new walk-in clinics being offered by major commercial pharmacies. In addition to other variables, the research assistant collected age (measured as less than 21, 21 - 65, and older than 65 years of age), sex, height, weight, and reason for the visit. Classify each of the variables as continuous numerical, discrete numerical, or categorical.

Height and weight are continuous numerical variables. Age as measured by the research assistant is ordered categorical. Sex and the reason for the visit are nominal categorical variables; sex has two categories, while reason for the visit will have many possible values.

⦿ **Guided Practice 1.2** Characterize the variables `participant.ID`, `treatment.group`, and `overall.V60.outcome` from the LEAP study (discussed in Section 1.1).¹⁰

¹⁰These variables measure non-numerical quantities, and thus are categorical variables. The variables `treatment.group` and `outcome.V60.overall` have two values or levels, while `participant.ID` has many possible values.

1.2.3 Relationships between variables

Many studies are motivated by a researcher examining a possible relationship between two or more variables. Statistical relationships between two variables occur when they tend to vary in a related way. A **response variable** measures an outcome of interest, while an **explanatory variable** may be useful in predicting or understanding the response variable. There may be several possible explanatory variables for a single response variable in a given study.

Researchers were interested in using the `famuss` data to answer several questions, including: is ACTN3 genotype associated with variation in muscle function? The ACTN3 gene codes for a protein involved in muscle function. A common mutation (polymorphism) at residue 577 in the ACTN3 gene changes C to T; TT individuals are unable to produce any ACTN3 protein in their muscle. Thus, researchers hypothesized that ACTN3 genotype might influence muscle function in humans. The response variable in this study is `ndrm.ch`, the change in non-dominant arm strength, with strength gain being used as a way to measure muscle function. The explanatory variable of interest is `actn3.r557x`, ACTN3 genotype at residue 577.

Both numerical and graphical ways to examine possible relationships between two variables will be covered later in the text.

- **Example 1.3** In the study conducted on Tibetan frogs, researchers collected measurements on egg clutches and female frogs at 11 study sites, located at differing altitudes. Identify the explanatory and response variables in the study.

The explanatory variable examined in the study is `altitude`. The variables `egg.size`, `clutch.size`, `clutch.volume`, and `body.size` are response variables measuring the level of maternal investment.

- **Guided Practice 1.4** Refer to the variables from the `famuss` data set described in Table 1.6 to formulate two questions about the relationships between these variables that differ from the one addressed by the research team.¹¹

¹¹Two sample questions: (1) Do participants appear respond differently to training according to race? (2) Do male participants appear to respond differently to training than females?

1.3 Data collection principles

The first step in conducting research is to identify questions to investigate. A clearly articulated research question is essential for selecting subjects to be studied, identifying relevant variables, and determining how data should be measured. In order to obtain reliable data, it is also important to consider *how* data are collected.

1.3.1 Populations and samples

1. What is the average mercury content in swordfish in the Atlantic Ocean?
2. If an infant seems predisposed to a peanut allergy, is it better to introduce or to avoid peanut products during the first 6 months of the infant's life?
3. What proportion of female college students experience sexual victimization?

Each of these questions refers to a target **population**. In the first question, the target population is all swordfish in the Atlantic ocean, and each fish represents a case. Almost always, it is either too expensive or logistically impossible to collect data for every case in a population, so nearly all research is based on samples from populations. A **sample** represents a subset of the cases and is often a small fraction of the population. For instance, 60 swordfish (or some other number) in the population might be selected, and this sample data may be used (with some assumptions) to provide an estimate of the population average and answer the research question.

DH: Removed the exercise question on identifying samples for three reasons: the stent example is gone; it refers to the stent example as if it was a drug (it isn't); and I want us to be both realistic and clear about samples. They are almost never random samples from the ideal target population. If asked, I think almost any student, even at the high school level would say that drawing a random sample from the population of people with a disease is fundamentally impossible. We can replace the stent example by LEAP, but as in other trials, the validity comes from the randomization among the recruited subjects, not the assumption of it being a RS.

1.3.2 Anecdotal evidence

Anecdotal evidence is typically composed of unusual observations that are easily recalled based on their striking characteristics. Physicians are sometimes more likely to remember the characteristics of a single patient with an unusually good response to a drug as opposed to the many patients who did not respond. The dangers of drawing general conclusions from anecdotal information are obvious; no single observation can be used to draw conclusions about a population. Often, the anecdotal case may not have been remembered correctly or may have involved errors in measurements. To learn about the characteristics of a population, it is necessary to examine a sample of many cases drawn randomly from the population.

Thomas Jefferson, the second president of the United States, recognized the pitfall of drawing conclusions from a single observation. In a letter to his nephew, he wrote "The patient, treated on the fashionable theory, sometimes gets well in spite of the medicine. The medicine therefore restored him, and the young doctor receives new courage to proceed in his bold experiments on the lives of his fellow creatures."¹²

JV: I find the reference to TJ rather funny, but I suspect the story only appeals to people with a very specific sense of humor...Would it break your heart to omit this? Doing so may also improve the flow here.

While it is incorrect to generalize from individual observations, scientists know that unusual observations can sometimes be valuable; such observations may be a reason to question previously held assumptions or to design a study to examine an unconventional perspective. Cures for certain diseases have been discovered through research inspired by a patient with a disease thought to be incurable responding to a new drug. *DH: Insert references sent by D Longo and D. Spriggs here. JV: This last sentence is too vague/unwieldy, adjust once references are included.*

An anecdotal observation can never be the basis for a conclusion, but it may well lead to the design of a more systematic study that could be definitive.

¹²Jefferson, T.(1985). Letters, 1760–1826. Ed. Merrill D. Peterson. New York: Viking.

1.3.3 Sampling from a population

Sampling from a population is a useful tool in population-based research in the health sciences. When done carefully, it provides reliable information about the health characteristics of a large population without having to directly measure those characteristics for each member, which is often an impossible task. The US Centers for Disease Control (US CDC) conducts many such surveys, including the Behavioral Risk Factors Surveillance System (BRFSS)¹³. The BRFSS conducts approximately 400,000 telephone interviews annually to ask U.S. residents questions regarding their health-related risk behaviors, chronic health conditions, and use of preventive services. The CDC conducts similar surveys for diabetes, health care access, and immunization. Likewise, the World Health Organization (WHO) conducts the World Health Survey in partnership with approximately 70 countries to learn about the health of adult populations and the health systems in those countries.¹⁴ In 2000, the US Department of Justice released the *The Sexual Victimization of College Women*, based on a survey conducted in 1996 of 4,446 undergraduate women.¹⁵

DH: we should check to see if the data are available at DoJ website. JV: The page for the sexual victimization survey only references NCVS data, which is freely available online (scroll down to "Codebooks and Datasets" under Documentation section), but I couldn't find data for questions specific to that survey. Perhaps better to switch to a survey for which data are available online? For example, Gender and Violent Victimization, see <http://doi.org/10.3886/ICPSR27082.v1>

DH: placeholder for the Harvard survey, despite its flaws

Sampling from a population is easier when the population is relatively small and members of the population are easy to identify and contact. For instance, the quality care team at an integrated health care system, such as Kaiser Permanente or Harvard Pilgrim Health Care, might like to learn about how members of the system perceive quality of care. Since health plans have contact information for each of their members, a selected subset can be contacted (with their consent) for participation in an interview or mailed survey. More complex methods are required for other surveys, such as the study on sexual

¹³<http://www.cdc.gov/brfss/>

¹⁴<http://www.who.int/healthinfo/survey/en/>

¹⁵<https://www.ncjrs.gov/pdffiles1/nij/182369.pdf>




Figure 1.8: Instead of sampling from all members equally, approaching members visiting a clinic during a particular week would disproportionately select members who typically use outpatient services.

victimization of college women.

One common downfall in conducting a sample is to use a **convenience sample**, in which individuals who are easily accessible are more likely to be included in the sample. For instance, the quality control team in the healthcare plan might ask interviewers to approach plan members visiting an outpatient clinic during a particular week. The sample would fail to enroll generally healthy members who typically do not use outpatient services or schedule routine physical examinations. Similarly, the Department of Justice could have only sampled women in colleges or universities in or near the District of Columbia.

The general principle of sampling is straightforward; a sample from a population is useful for learning about a population only when the sample matches, on average, the characteristics of the population. Random sampling is the best way to ensure that a sample reflects a population, because random samples do not reflect the conscious or unconscious bias of the team gathering the sample. However, even a well-defined sampling strategy can lead to an unrepresentative sample if there are substantial barriers to subject participation, such as questions that assume participants are fluent in English or calls to potential participants that do not account for working hours or time-zone differences.

The easiest random samples to analyze are those in which each member of a population has the same chance of being sampled. In a **simple random sample**, each member of the population is directly chosen at random for the sample, with probability the size of the sample divided by the size of the population. Simple random samples are essentially equivalent to how raffles are conducted. For example, if there are 5 prizes available and

100 people each have a single ticket, each person has a 5% chance ($5/100$) of being called. In the health plan example, a subset of members might be chosen randomly from the plan membership roster for an interview. *OpenIntro*, third edition, Section 1.4.2 describes the four most commonly used sampling strategies.




Figure 1.9: In this graphic, five members are randomly selected from the population to be interviewed.

Sometimes a simple random sample is difficult to implement and an alternative method is helpful. One such substitute is a **systematic sample**, in which one case is sampled after letting a fixed number of other cases pass by. Since this approach uses a mechanism that is not easily subject to personal biases, it often yields a reasonably representative sample. This book will focus on random samples since the use of systematic samples is uncommon and requires additional considerations of the context. *JV: It may be possible to omit this paragraph. If not, needs an example.*

The act of taking a simple random sample helps minimize bias, but bias can crop up in other ways. Even when people are picked at random, caution must be exercised if the **non-response** is high. For instance, if only 30% of the people randomly sampled for a survey actually respond, then it is unclear whether the results are truly **representative** of the entire population. Such **non-response bias** can skew results; it is important to minimize barriers that might discourage subject participation in order to collect reliable data.

• Guided Practice 1.5

DH: replace this with a better example We can easily access ratings for products, sellers, and companies through websites. These ratings are based only on those people who go out of their way to provide a rating. If 50% of online reviews for a product




Figure 1.10: Due to the possibility of non-response, surveys studies may only reach a certain group within the population. For example, a survey written in English may only result in responses from health plan members fluent in English.

are negative, do you think this means that 50% of buyers are dissatisfied with the product?¹⁶

1.3.4 Introducing experiments and observational studies

Experiments and observational studies are the two primary types of study designs used to collect data.

When researchers want to investigate the possibility of a causal connection, they conduct an **experiment**. For instance, it might be hypothesized that administering a certain drug will reduce mortality in heart attack patients. To find evidence for a causal connection between the explanatory and response variables, researchers will collect a sample of individuals and randomly assign them into one of two groups. The first group, called a control group, may receive either a **placebo** (an inert substance with the appearance of the study drug) or a commonly used drug known to have some effect; the second group (the experimental group) receives the new drug.

Researchers perform an **observational study** when they collect data in a way that does not directly interfere with how the data arise. For instance, to study why certain diseases develop, researchers may collect information through conducting surveys, reviewing medical or company records, or following a **cohort** of many similar individu-

¹⁶Answers will vary. From our own anecdotal experiences, we believe people tend to rant more about products that fell below expectations than rave about those that perform as expected. For this reason, we suspect there is a negative bias in product ratings on sites like Amazon. However, since our experiences may not be representative, we also keep an open mind.

als. In each of these situations, researchers merely observe the data that arise. Observational studies can provide evidence of an association between variables, but they cannot by themselves show a causal connection. In general, causation can only be inferred from a randomized experiment.

1.3.5 Experiments

Studies in which researchers assign treatments to cases are called **experiments**. Randomized experiments are generally built on three principles.

Controlling. Researchers assign treatments to cases, doing their best to **control** for any other differences in the groups. For example, all infants enrolled in the LEAP study were required to be between 4 and 11 months of age, with severe eczema and/or allergies to eggs.

Randomization. Researchers randomize patients into treatment groups to account for variables that cannot be controlled. For example, some infants may have been more susceptible to peanut allergies because of an unmeasured genetic condition. Randomly assigning patients to the treatment or control group helps even out such differences. In situations where researchers suspect that variables other than the treatment may influence the response, they may first group individuals into **blocks** and then, within each block, randomize cases to treatment groups; this technique is referred to as **blocking** or **stratification**. In the LEAP study, infants were stratified into two cohorts based on whether or not the child developed a red, swollen mark (a wheal) after a skin test at the time of enrollment. The main analysis of the study analyzed data collected for infants without a wheal after the skin test. Figure 1.11 illustrates the blocking scheme used in the study. General methods for analyzing blocked data are relatively complicated and will not be covered in this book.

Replication. The more cases researchers observe, the more accurately they can estimate the effect of the explanatory variable on the response. In a single study, **replication** is accomplished by collecting a sufficiently large sample. The LEAP study randomized a total of 640 infants; 542 infants were in the block without the wheal response.

It is important to incorporate the three experimental design principles into any study; this book describes applicable methods for analyzing data from such experiments. Blocking is a slightly more advanced technique, and statistical methods in this book may be extended to analyze data collected using blocking.

DH: Update the following figure for the LEAP data. JV: I changed the labels, but I'm getting lost in how to change the number in individuals in each group so that they are not equal.

1.3.6 Reducing bias in human experiments

Randomized experiments are the gold standard for data collection, but they do not automatically ensure an unbiased perspective in all cases. Human studies are perfect examples where bias can arise unintentionally.

In 1980, researchers reported the results of a study assessing the efficacy of a new drug used to treat heart attack patients.¹⁷ Researchers wanted to know whether the drug reduced deaths in patients; in order to draw causal conclusions about the effect of the drug, they designed a randomized experiment in which study volunteers were randomly assigned to one of two study groups.¹⁸ The **treatment group** received the drug; the other group, called the **control group**, did not receive any drug treatment.

Typically, researchers do not want patients to know which group they are in. The emotional response of a patient who knows they are either receiving or not receiving a potentially helpful new drug may cause different behavior between the two groups. In order to eliminate this source of bias from the study, researchers conduct a **blinded** study in which patients are kept uninformed about their treatment. Patients in the control group, instead of being given a drug, are given an inert substance called a **placebo**. An effective placebo is the key to making a study truly blind. A placebo may often result in a slight but real improvement in patients; this effect is referred to as the **placebo effect**.¹⁹

The patients are not the only ones who should be blinded: doctors and researchers can accidentally bias a study. For example, out of concern for potential side effects of

¹⁷Anturane Reinfarction Trial Research Group. 1980. Sulfipyrazone in the prevention of sudden death after myocardial infarction. *New England Journal of Medicine* 302(5):250-256.

¹⁸Human subjects are often called **patients**, **volunteers**, or **study participants**.

¹⁹Kapchuk, TJ and Miller, FG. 2015. Placebo effects in medicine, *New England Journal of Medicine*, 373(1):8-9.




Figure 1.11: A simplified schematic of the blocking scheme used in the LEAP study, depicting 54 patients that underwent randomization. Patients are first divided into groups based on response to the initial skin test, then each block is evenly separated into the treatment groups using randomization. This strategy ensures an even representation of patients in each treatment group from both the skin test positive and skin test negative groups.

a new drug, a doctor might inadvertently give a patient in the treatment group more attention and care than they would to a patient known to be taking a placebo. To guard against this bias, which has also been found to have a measurable effect in some instances, most modern studies employ a **double-blind** setup in which doctors who interact with patients are, just like the patients, unaware of who is or is not receiving the experimental treatment.²⁰

- **Guided Practice 1.6** Look back to the study in Section 1.1 in which researchers were testing whether peanut product consumption was effective at reducing the likelihood of peanut allergies in children at-risk for these allergies. Is this an experiment? Was the study blinded? Was it double-blinded? ²¹

1.3.7 Observational studies

Generally, data in observational studies are collected only by monitoring what occurs, while experiments require researchers to assign the primary explanatory variable in a study for each subject *JV*: *This sentence should be reworded for clarity, not sure how.* Making causal conclusions based on experiments is often reasonable; however, making the same causal conclusions solely based on observational data should be avoided.


DH: This is another instance where absolute statements are risky. The data on smoking and lung cancer are all observational, or were until a short time ago. I wonder if we should mention that. JV: I have adjusted the language in the first paragraph. Scroll down past the sunscreen example to see where I think the mention of smoking and lung cancer can come in.

Suppose an observational study tracked sunscreen use and skin cancer, finding that the more sunscreen a person uses, the more likely they are to have skin cancer. However, this does not mean that sunscreen *causes* skin cancer. One important piece of missing information is sun exposure – if someone is out in the sun all day, they are both more likely to use sunscreen and to get skin cancer. Sun exposure is a **confounding variable**:

²⁰There are always some researchers involved in the study who do know which patients are receiving which treatment. However, they do not directly interact with patients and do not tell the blinded health care professionals who is receiving which treatment.

²¹The researchers assigned the patients into their treatment groups, so this study was an experiment. However, the patients (and their parents) could distinguish which treatment they received, so this study was not blind. The study could not be double-blind since it was not blind.

a variable correlated with both the explanatory and response variables.²² There is no guarantee that all confounding variables can be examined or measured; as a result, it is difficult to justify making causal conclusions from observational studies.



Observational studies are useful in that they can reveal interesting patterns or associations, providing researchers with the information necessary to design follow-up experiments. For example...

JV: Smoking and lung cancer can come in here, or anything else that illustrates the point, which I think is an important one. There might also be an example here that asks for a description of a reasonable follow-up experiment from some observational data – frog and famuss both seem too difficult, though.

Observational studies come in two forms: prospective and retrospective studies. A **prospective study** identifies individuals and collects information as events unfold. For instance, medical researchers may identify and follow a group of similar individuals over many years to assess the possible influences of behavior on cancer risk. One example of such a study is The Nurses' Health Study, started in 1976 and expanded in 1989.²³ This prospective study recruits registered nurses and then collects data from them using questionnaires. **Retrospective studies** collect data after events have taken place, e.g. researchers may review past events in medical records. Some datasets may contain both prospectively- and retrospectively-collected variables. *DH: need an example of this. famuss does not qualify, I think. The flanders dental study would qualify but has not been introduced at this point. JV: The dental study can be introduced here in the same way as the Nurses' Health study, then.*

DH: I have commented out the section on sampling methods, though I like it. It will take some work to find example to make the sampling methods concrete. I have left the OpenIntro

²²Also called a **lurking variable**, **confounding factor**, or a **confounder**.

²³www.channing.harvard.edu/nhs

source for this topic in our source.

1.4 Examining numerical data

This section introduces techniques for exploring and summarizing numerical variables, using the frog data from Section 1.2.

1.4.1 Measures of center: mean and median

The **mean**, sometimes called the average, is a common way to measure the center of a **distribution** of data. To find the average clutch volume for the observed egg clutches, we add up all the clutch volumes and divide by the total number of clutches. For computational convenience, the volumes are rounded to the first decimal.

$$\bar{x} = \frac{177.8 + 257.0 + \cdots + 933.3}{431} = 882.5 \text{ mm}^3 \quad (1.7)$$

\bar{x}
sample
mean

The sample mean is often labeled \bar{x} . The letter x is being used as a generic placeholder for the variable of interest, `clutch.volume`, and the bar over the x communicates that the average volume of the 431 clutches is 882.5 mm^3 . It is useful to think of the mean as the balancing point of the distribution. *JV: This last sentence needs some further clarification if is to be included – I do not think the meaning is obvious to someone new to statistics.*

Mean

The sample mean of a numerical variable is computed as the sum of all of the observations divided by the number of observations:

$$\bar{x} = \frac{x_1 + x_2 + \cdots + x_n}{n} \quad (1.8)$$

where x_1, x_2, \dots, x_n represent the n observed values.

Another measure of center is the **median**, which is the middle number in a distri-

bution after the values have been ordered from smallest to largest. If the distribution contains an even number of observations, the median is the average of the middle two observations. There are 431 clutches in the dataset, so the median is the clutch volume of the 216th observation in the sorted values of `clutch.volume`: 831.8 mm³.

DH: We have removed the concept of a weighted mean here; we do not have a context for it. But it is an important idea that can be profitably used later, perhaps with the `brfss` data. Perhaps we can re-insert it later. JV: Recommend introducing weighted mean in probability with calculating expected value.

1.4.2 Measures of spread: standard deviation and interquartile range

DH: Don't like the verbal description of the sd here, but have not replaced it yet. Note also that I have changed some bar to overline. Let me know which you think is better in the pdf; I like overline. JV: I also like overline, since I prefer reading at relatively small magnification. I will change the rest to overline.

The standard deviation measures approximately the distance between a typical observation and the mean. The distance of an observation from its mean is its **deviation**. Below are the deviations for the 1st, 2nd, 3rd, and 431th observations in the `clutch.volume` variable. For computational convenience, clutch volume is rounded to the first decimal.

$$\begin{aligned}x_1 - \bar{x} &= 177.8 - 882.5 = -704.7 \\x_2 - \bar{x} &= 257.0 - 882.5 = -625.5 \\x_3 - \bar{x} &= 151.4 - 882.5 = -731.1 \\&\vdots \\x_{431} - \bar{x} &= 933.2 - 882.5 = 50.7\end{aligned}$$

If we square these deviations and then take an average, the result is the sample **variance**, denoted by s^2 :

s^2
sample
variance

$$\begin{aligned}
 s^2 &= \frac{(-704.7)^2 + (-625.5)^2 + (-731.1)^2 + \cdots + (50.7)^2}{431 - 1} \\
 &= \frac{496,602.09 + 391,250.25 + 534,507.21 + \cdots + 2570.49}{430} \\
 &= 143,680.9
 \end{aligned}$$

The denominator is $n - 1$ rather than n when computing the variance; this mathematical nuance comes from statistical theory and the reason for doing so is not covered in this text.

The **standard deviation** is the square root of the variance:

$$s = \sqrt{143,680.9} = 379.05$$

s

sample
standard
deviation

The standard deviation of clutch volume for the egg clutches observed is about 380 mm^3 .
DH: excellent place to give an interpretation of sd, referring back to verbal definition, or perhaps to use it to mention the empirical rule, which is in the caption to one of the Open Intro plots. JV: Return to this once verbal definition of SD is refined. The empirical rule is best introduced with a picture – I considered introducing it in an example after histograms, but that would interrupt the flow. May work in the transforming data subsection.

Formulas and methods used to compute the variance and standard deviation for a population are similar to those used for a sample.²⁴ However, like the mean, the population values have special symbols: σ^2 for the variance and σ for the standard deviation. The symbol σ is the Greek letter *sigma*.

σ^2

population
variance

σ

population
standard
deviation

²⁴The only difference is that the population variance has a division by n instead of $n - 1$.

Standard Deviation

The sample standard deviation of a numerical variable is computed as the square root of the variance, which is the sum of squared deviations divided by the number of observations minus 1.

$$s = \sqrt{\frac{(x_1 - \bar{x})^2 + (x_2 - \bar{x})^2 + \cdots + (x_n - \bar{x})^2}{n - 1}} \quad (1.9)$$

where x_1, x_2, \dots, x_n represent the n observed values.

Variability can also be measured using the **interquartile range** (IQR). To calculate the IQR, find the **first quartile** (the 25th percentile, i.e. 25% of the data fall below this value) and the **third quartile** (the 75th percentile). These are often labeled Q_1 and Q_3 , respectively. The IQR is the difference: $Q_3 - Q_1$.

The IQR for `clutch.volume` is $1096.0 - 609.6 = 486.4 \text{ mm}^3$. The middle 50% of the values for `clutch.volume` lie between 609.6 mm^3 and 1096.0 mm^3 .

1.4.3 Robust statistics

In the `frog` data, there are four observed clutch volumes larger than $2,000 \text{ mm}^3$ (2138.0, 2630.3, 2454.7, 2511.9). These values can be clearly identified by plotting the data as points on a single axis, as shown in Figure 1.12; this basic graphical display is known as a **dot plot**. How do these extreme values affect the summary statistics for the clutch volume variable in the `frog` data?




Figure 1.12: Dot plot of the clutch volume variable in the `frog` data.

The sample statistics are computed under each of two scenarios in Table 1.13, one

with and one without the four largest observations. The median and IQR are referred to as **robust estimates** because extreme observations have little effect on their values. For these data, the median does not change, while the IQR differs by only about 6 mm³. In contrast, the mean and standard deviation are much more affected, particularly the standard deviation; since standard deviation depends on the squared distances from the mean, its change in the presence of large observations is more noticeable. Typically, extreme observations have a greater effect on the standard deviation than on the mean.

scenario	robust		not robust	
	median	IQR	\bar{x}	s
original frog data	831.8	486.9	882.5	379.1
drop four largest observations	831.8	493.92	867.9	349.2

Table 1.13: A comparison of how the median, IQR, mean (\bar{x}), and standard deviation (s) change when extreme observations are present.

JV: Agree that famuss data can be used in end-of-chapter exercise.

1.4.4 Visualizing distributions of data: histograms and boxplots

Visualizing how data are distributed can reveal characteristics of the data that are not obvious from summary statistics. Graphical summaries, such as histograms and boxplots, complement the information provided by numerical summaries.

Dot plots show the exact value of each observation; while this is useful for small datasets, dot plots are not ideal for larger samples. Instead, observations can be grouped into bins and plotted as bars to form a **histogram**. Table 1.14 shows the number of clutches with volume between 0 and 200 mm³, 200 and 400 mm³, etc. up until 2,600 and 2,800 mm³. These binned counts are plotted in Figure 1.15.

Clutch volumes	0-200	200-400	400-600	600-800	...	2400-2600	2600-2800
Count	4	29	69	99	...	2	1

Table 1.14: The counts for the binned `clutch.volume` data.

Histograms provide a view of the **data density**. Higher bars indicate more common observations, while lower bars represent relatively rare observations. For instance, there




Figure 1.15: A histogram of `clutch.volume`.

are many more egg clutches with volumes smaller than 1,500 mm³ than clutches with larger volumes. The bars make it easy to see how the density of the data changes relative to clutch volume.

Histograms are especially convenient for describing the shape of the data distribution. Figure 1.15 shows that most clutches have a relatively small volume, while fewer clutches are very large. When data trail off to the right, with a long right tail, the data are said to be **right skewed**.²⁵ Data with the reverse characteristic – a long, thin tail to the left – are said to be **left skewed**. The term **symmetric** is used to describe data that show roughly equal trailing off in both directions.

A **mode** is represented by a prominent peak in the distribution.²⁶ Figure 1.16 shows histograms that have one, two, or three prominent peaks. Such distributions are called **unimodal**, **bimodal**, and **multimodal**, respectively. Any distribution with more than two prominent peaks is called multimodal. Notice that there was one prominent peak in the unimodal distribution with a second less prominent peak that was not counted since it only differs from its neighboring bins by a few observations. *JV: Last sentence should be refined – thoughts on how to communicate that "prominent" is a subjective term?*

²⁵Other ways to describe data that are skewed to the right: **skewed to the right**, **skewed to the high end**, or **skewed to the positive end**.

²⁶Another definition of mode, which is not typically used in statistics, is the value with the most occurrences. It is common to have *no* observations with the same value in a data set, which makes this other definition useless for many real datasets.




Figure 1.16: From left to right: unimodal, bimodal, and multimodal distributions.

• **Guided Practice 1.10** Describe the distribution of `clutch.volume` using the histogram in Figure 1.15. Are the data skewed? Is it a unimodal, bimodal, or multimodal distribution? ²⁷

A **boxplot** summarizes a dataset using five statistics while also plotting unusual observations.²⁸ Figure 1.17 provides a vertical dot plot alongside a boxplot of the `clutch.volume` variable from the `frog` dataset.

In a boxplot, a rectangle extending from the first quartile to the third quartile represents the middle 50% of the data (the IQR); the rectangle is split in half by the **median**. Extending outwards from the box, the **whiskers** capture the data that fall between $Q_1 - 1.5 \times IQR$ and $Q_3 + 1.5 \times IQR$.²⁹ Note that the whiskers must end at data points; the values given by adding or subtracting $1.5 \times IQR$ define the maximum reach of the whiskers. For example, for the `clutch.volume` variable: $Q_3 + 1.5 \times IQR = 1,096.5 - 1.5 \times 486.4 = 1,826.1 \text{ mm}^3$. However, there was no clutch with volume $1,826.1 \text{ mm}^3$; thus, the upper whisker extends to $1,819.7 \text{ mm}^3$, the largest observation that is smaller than $Q_3 + 1.5 \times IQR$.

Any observation that lies beyond the whiskers is labeled with a dot; these observations are called outliers. An **outlier** is a value that appears extreme relative to the rest of the data. For the `clutch.volume` variable, there are several large outliers and no small

²⁷The data is strongly skewed to the right; while many counts fall in the 600-1,000 mm^3 range, there are a few clutches with volume greater than 1,500 mm^3 . The distribution is unimodal, with only one prominent peak.

²⁸Boxplots are also known as box-and-whisker plots.

²⁹While the choice of exactly 1.5 is arbitrary, it is a commonly used value for drawing boxplots.




Figure 1.17: A vertical dot plot next to a labeled boxplot for the volumes of 431 egg clutches. The median (831.8 mm^3), splits the data into the bottom 50% and the top 50%, marked in the dot plot by horizontal dashes and open circles, respectively.

outliers, indicating the presence of some unusually large egg clutches. Outliers can potentially provide insight into interesting properties of the data.

1.4.5 Scatterplots

A **scatterplot** provides a case-by-case view of data for two numerical variables. In the frog data, `clutch.volume` and `body.size` are two numerical variables of interest; previous research has reported that larger body size allows females to produce larger egg clutches. The relationship between clutch volume and female body size is examined via scatterplot in Figure 1.18. In any scatterplot, each point represents a single case. Since body size was measured for 129 frogs, there are 129 points in Figure 1.18.

The variables `clutch.volume` and `body.size` are said to be **associated** because the plot shows a discernible pattern. Since the points tend to lie in a straight line, the two variables are **linearly associated**. Two variables are **positively associated** if increasing values of one tend to occur with increasing values of the other; similarly, variables are **negatively associated** if increasing values of one variable occurs with decreasing values of the other. Figure 1.18 shows an upward trend – as expected, larger frogs tend to pro-




Figure 1.18: A scatterplot showing `clutch.volume` (horizontal axis) vs. `body.size` (vertical axis).

duce egg clutches with larger volumes. Frog embryos are surrounded by a gelatinous matrix that may protect developing embryos from temperature fluctuation or ultraviolet radiation; these observations suggest that larger females are indeed capable of producing greater quantities of this material.

Figure 1.19 shows the relationship between height and weight for participants in the FAMuSS study. Each point on the plot represents a participant. As expected, taller participants tend to be heavier, so the variables height and weight are positively associated.

Taller people naturally tend to be heavier; as a consequence, weight itself is not a good measure of whether someone is overweight. Body mass index (BMI) is a measure of weight that is less affected by a person's height. A BMI of 30 or above is considered overweight.³⁰ In the metric system, BMI is calculated as weight in kilograms (kg) divided by height squared (m^2). If height and weight are measured in inches and pounds, as in the famuss data, then BMI is weight in pounds (lb) divided by height squared (in^2), then multiplied by 703. The famuss data includes the variable `bmi` for each participant, and Figure 1.20 shows the relationship between height and `bmi`. The strong upward trend in Figure 1.19 is no longer evident, indicating that height and `bmi` have a much weaker association. For this reason, health agencies such as the US NIH and the World Health

³⁰http://www.nhlbi.nih.gov/health/educational/lose_wt/risk.htm




Figure 1.19: A scatterplot showing height (horizontal axis) vs. weight (vertical axis). One participant 70.5 inches tall and weighing 308 pounds is highlighted.

Organization (WHO) use BMI as a measure of obesity.

If two variables are not associated, then they are said to be **independent**. That is, two variables are independent if there is no evident relationship between the two. Generally, it is not easy to determine definitively whether two variables are independent from looking at a scatterplot, even in Figure 1.20.

JV: Need an example here showing a nonlinear relationship (along the lines of the car price vs weight example). Anything in famuss or LEAP? Might even work to quickly introduce caries data.

1.4.6 Transforming data (special topic)

DH: We should include a section on transforming data, esp since the frog data has been transformed. JV: Agree. I replaced the MLB histograms with ones for the clutch volume data; left the scatterplots as a placeholder, as I'm not sure the clutch volume vs body size plot is a good candidate for that example. This would be a good place to introduce the empirical rule, since the transformation on frogs makes the data look more normal; empirical rule will be discussed in more detail once normal distribution is introduced, anyways (so seems fine that it would first be mentioned as part of a special topic).

Scientists may choose to transform strongly skewed data in order to make them easier




Figure 1.20: A scatterplot showing height (horizontal axis) vs. bmi (vertical axis). The same individual highlighted in Figure 1.19 is marked here, with BMI 43.56.

to model and analyze. A **transformation** is a rescaling of the data using a function. Consider the histogram of egg clutch volumes from the `frog` data, shown in Figure 1.21(a). In the published paper, researchers used a \log_{10} transformation on the data before conducting analyses. Figure 1.21(b) shows a plot of the \log_{10} of clutch volumes.




Figure 1.21: (a) Histogram of egg clutch volumes. (b) Histogram of the log-transformed egg clutch volumes.

JV: Text after this point (in this section) needs re-writing.

There are some standard transformations that are often applied when much of the data cluster near zero (relative to the larger values in the data set) and all observations




Figure 1.22: (a) Scatterplot of `line_breaks` against `num_char` for 50 emails. (b) A scatterplot of the same data but where each variable has been log-transformed.

are positive. A **transformation** is a rescaling of the data using a function. For instance, a plot of the natural logarithm³¹ of player salaries results in a new histogram in Figure ?. Transformed data are sometimes easier to work with when applying statistical models because the transformed data are much less skewed and outliers are usually less extreme.

Transformations can also be applied to one or both variables in a scatterplot. A scatterplot of the `line_breaks` and `num_char` variables is shown in Figure 1.22(a), which was earlier shown in Figure ?. We can see a positive association between the variables and that many observations are clustered near zero. In Chapter ?, we might want to use a straight line to model the data. However, we'll find that the data in their current state cannot be modeled very well. Figure 1.22(b) shows a scatterplot where both the `line_breaks` and `num_char` variables have been transformed using a log (base e) transformation. While there is a positive association in each plot, the transformed data show a steadier trend, which is easier to model than the untransformed data.

Transformations other than the logarithm can be useful, too. For instance, the square root ($\sqrt{\text{original observation}}$) and inverse ($\frac{1}{\text{original observation}}$) are used by statisticians. Common goals in transforming data are to see the data structure differently, reduce skew, assist in modeling, or straighten a nonlinear relationship in a scatterplot.

³¹Statisticians often write the natural logarithm as `log`. You might be more familiar with it being written as `ln`.

1.5 Considering categorical data

Like numerical data, categorical data can also be organized and analyzed; however, numerical calculations cannot be done with categorical data. In this section, we will introduce tables and other basic tools for categorical data, using the famuss dataset introduced in Section 1.2.2.

1.5.1 Contingency tables

A table for a single variable is called a **frequency table**. Table 1.23 is a frequency table for the `actn3.r577x` variable. Recall that `actn3.r577x` is a categorical variable describing genotype at a location on the ACTN3 gene: CC, CT, or TT. If we replaced the counts with percentages or proportions, the table would be called a **relative frequency table**.

	CC	CT	TT	Sum
Counts	173	261	161	595

Table 1.23: A frequency table for the `actn3.r577x` variable.

Table 1.24 summarizes two variables: `race` and `actn3.r577x`. A table that summarizes data for two categorical variables in this way is called a **contingency table**.³² Each value in the table represents the number of times a particular combination of variable outcomes occurred. For example, the first row of the table shows that of the African-American individuals, 16 are CC, 6 are CT, and 5 are TT.

Row and column totals, known collectively as **marginal totals**, are also included. The **row totals** provide the total counts across each row; **column totals** are the total counts down each column.

JV: Not sure how they added the variable labels to their table. I have left the OI tables and code in the source.

Table 1.25 shows the row proportions for Table 1.24. The **row proportions** are computed as the counts divided by their row totals. The value 16 at the intersection of African American and CC is replaced by $16/27 = 0.593$; i.e., 16 divided by the row total, 27. The

³²Contingency tables are also known as **two-way tables**.

	CC	CT	TT	Sum
African Am	16	6	5	27
Asian	21	18	16	55
Caucasian	125	216	126	467
Hispanic	4	10	9	23
Other	7	11	5	23
Sum	173	261	161	595

Table 1.24: A contingency table for race and actn3.r577x.

value 0.593 corresponds to the proportion of African-Americans in the study with genotype CC.

	CC	CT	TT	Sum
African Am	$16/27 = 0.593$	$6/27 = 0.222$	$5/27 = 0.185$	$27/27 = 1.00$
Asian	$21/55 = 0.382$	$18/55 = 0.327$	$16/55 = 0.291$	$55/55 = 1.00$
Caucasian	$125/467 = 0.267$	$216/467 = 0.463$	$126/467 = 0.270$	$467/467 = 1.00$
Hispanic	$4/23 = 0.174$	$10/23 = 0.435$	$9/23 = 0.391$	$23/23 = 1.00$
Other	$7/23 = 0.304$	$11/23 = 0.478$	$5/23 = 0.217$	$23/23 = 1.00$
Sum	$173/595 = 0.291$	$261/595 = 0.438$	$161/595 = 0.271$	$595/595 = 1.00$

Table 1.25: A contingency table with row proportions for the race and actn3.r577x variables.

● **Example 1.11** What does Table 1.25 highlight about the distribution of genotypes between different populations?

Genotype distributions vary between populations. For the Caucasian individuals sampled in the study, CT is the most common genotype at 46.3%. In contrast, over half (59.3%) of African Americans sampled are CC. CC is also the most common genotype for Asians, but in this population, genotypes are more evenly distributed: 38.2% of Asians sampled are CC, 32.7% are CT, and 29.1% are TT.

A contingency table of the column proportions is computed in a similar way, in which each **column proportion** is computed as the count divided by the corresponding column total. Table 1.26 shows such a table, and here the value 0.092 indicates that 9.2% of CC individuals in the study are African-American.

JV: Not sure if the following exercise is very clear, but there should be something here to warn against misinterpretation of the column proportions in this context. Important to point

	CC	CT	TT	Sum
African Am	16/173 = 0.092	6/261 = 0.037	5/161 = 0.191	27/595 = 0.045
Asian	21/173 = 0.080	18/261 = 0.104	16/161 = 0.993	55/595 = 0.092
Caucasian	125/173 = 0.776	216/261 = 0.828	126/161 = 0.728	467/595 = 0.785
Hispanic	4/173 = 0.023	10/261 = 0.062	9/161 = 0.034	23/595 = 0.038
Other	7/173 = 0.027	11/261 = 0.063	5/161 = 0.031	23/595 = 0.038
Sum	173/173 = 1.000	261/261 = 1.000	161/161 = 1.000	595/595 = 1.000

Table 1.26: A contingency table with column proportions for the race and `actn3.r577x` variables.

out that one limitation of the data is uneven representation between groups.

- ⊙ **Guided Practice 1.12** As computed in Table 1.26, 77.6% of CC individuals in the study are Caucasian. Does this data suggest that in the general population, people of CC genotype are highly likely to be Caucasian?³³

1.5.2 Bar plots

A bar plot is a common way to display a single categorical variable. The left panel of Figure 1.27 shows a **bar plot** for the `actn3.r577x` variable. In the right panel, the counts are converted into proportions (e.g. $173/595 = 0.291$ for the CC genotype), showing the proportion of observations that are in each level (i.e. in each category).




Figure 1.27: Two bar plots of `actn3.r577x`. The left panel shows the counts, and the right panel shows the proportions for each genotype.

³³No, this is not a reasonable conclusion to draw from the data. The high proportion of Caucasians among CC individuals primarily reflects the large number of Caucasians sampled in the study – 78.5% of the people sampled are Caucasian. The uneven representation of different races is one limitation of the famuss data.

JV: After poking around in the R code for the segmented plots, I found out that they are made by overlaying a second graph on top of the other – a method that only works well for one category mapped onto the total; I tested out this method to make Figure 1.28, showing Caucasians as a part of the total, but it’s not a particularly interesting/relevant plot.

JV: With the help of Google, I figured out how to make segmented bar plots using a different method – plots follow, one using genotype for the bars and one using race for the bars. The following paragraph needs to be re-written based on which plots we ultimately decide to include, and introduce the context for examining race and genotype together (mutant allele known to exist at different frequencies in various human populations). I did not find a way to make the standardized segmented plots.

Segmented bar plots provide a way to visualize the information in contingency tables. A **segmented bar plot** is a graphical display of contingency table information. For example, a segmented bar plot using data from Table 1.24 is shown in Figure 1.28, where a bar plot was created using the `actn3.r577x` variable, with each group divided by the levels of race. The column proportions of Table 1.26 have been translated into a standardized segmented bar plot in Figure 1.28(b), which is a helpful visualization of the races represented in each level of `actn3.r577x`.




Figure 1.28: (a) Segmented bar plot for individuals by genotype, in which the counts have been further broken down by Caucasian (blue) and non-Caucasian (yellow). (b) Standardized version of Figure (a).

JV: I don’t particularly like the mosaic plots, and they may not be the best choice for dis-




Figure 1.29: Segmented bar plot for individuals by race, where the counts have been further broken down by genotype.




Figure 1.30: Segmented bar plot for individuals by genotype, where the counts have been further broken down by race.

playing the famuss data, anyways. I also think the pie chart section doesn't necessarily need to be included.

1.5.3 Comparing numerical data across groups

In this section, two convenient methods for examining numerical data across groups are introduced: side-by-side boxplots and hollow histograms. The **side-by-side boxplot** is a traditional tool for comparing across categories. The **hollow histogram** method plots the outlines of histograms for each group onto the same axes.

Recall the question introduced in Section 1.2.3: is ACTN3 genotype associated with variation in muscle function? To explore this question, genotype and variation in muscle function (measured by `ndrm.ch`) can be compared using side-by-side boxplots and hollow histograms, as shown in Figure 1.31. The histograms are useful for seeing distribution shape, skew, and groups of anomalies, while the side-by-side boxplots are especially useful for comparing centers and spreads. Comparison of median change in non-dominant arm strength between the two groups reveals that the TT genotype is associated with a greater increase in strength than CC or CT. In other words, the T allele appears to be associated with greater muscle function.




Figure 1.31: Side-by-side boxplot (left panel) and hollow histograms (right panel) for `ndrm.ch`, split by ACTN3 genotype.




Figure 1.32: Side-by-side boxplot comparing the distribution of `clutch.volume` for different altitudes.

Not all data will show such apparent trends. For example, consider the question of interest in the `frog` dataset: how does maternal investment vary with altitude? Researchers collected data at 11 altitudes from 2,035 to 3,495 m above sea level, measuring attributes of egg clutches such as clutch volume. A side-by-side boxplot comparing clutch volume across altitudes is shown in Figure 1.32. It seems that as a general rule, clutches found at higher altitudes have greater volume. However, more advanced statistical methods, such as those used in the published study, are required to thoroughly investigate the potential association between altitude and clutch size.

1.6 Exercises

1.6.1 Case study: using stents to prevent strokes

1.1 Migraine and acupuncture. A migraine is a particularly painful type of headache, which patients sometimes wish to treat with acupuncture. To determine whether acupuncture relieves migraine pain, researchers conducted a randomized controlled study where 89 females diagnosed with migraine headaches were randomly assigned to one of two groups: treatment or control. 43 patients in the treatment group received acupuncture that is specifically designed to treat migraines. 46 patients in the control group received placebo acupuncture (needle insertion at non-acupoint locations). 24 hours after patients received acupuncture, they were asked if they were pain free. Results are summarized in the contingency table below.³⁴

		<i>Pain free</i>		Total
		Yes	No	
<i>Group</i>	Treatment	10	33	43
	Control	2	44	46
	Total	12	77	89




Figure from the original paper displaying the appropriate area (M) versus the inappropriate area (S) used in the treatment of migraine attacks.

- What percent of patients in the treatment group were pain free 24 hours after receiving acupuncture? What percent in the control group?
- At first glance, does acupuncture appear to be an effective treatment for migraines? Explain your reasoning.
- Do the data provide convincing evidence that there is a real pain reduction for those patients in the treatment group? Or do you think that the observed difference might just be due to chance?

1.2 Sinusitis and antibiotics. Researchers studying the effect of antibiotic treatment for acute sinusitis compared to symptomatic treatments randomly assigned 166 adults diagnosed with acute sinusitis to one of two groups: treatment or control. Study participants received either a 10-day course of amoxicillin (an antibiotic) or a placebo similar in appearance and taste. The placebo consisted of symptomatic treatments such as acetaminophen, nasal decongestants, etc. At the end of the 10-day period patients were asked if they experienced significant improvement in symptoms. The distribution of responses are summarized below.³⁵

		<i>Self-reported significant improvement in symptoms</i>		Total
		Yes	No	
<i>Group</i>	Treatment	66	19	85
	Control	65	16	81
	Total	131	35	166

- What percent of patients in the treatment group experienced a significant improvement in symptoms? What percent in the control group?
- Based on your findings in part (a), which treatment appears to be more effective for sinusitis?
- Do the data provide convincing evidence that there is a difference in the improvement rates of sinusitis symptoms? Or do you think that the observed difference might just be due to chance?

³⁴G. Allais et al. "Ear acupuncture in the treatment of migraine attacks: a randomized trial on the efficacy of appropriate versus inappropriate acupoints". In: *Neurological Sci.* 32.1 (2011), pp. 173–175.

³⁵J.M. Garbutt et al. "Amoxicillin for Acute Rhinosinusitis: A Randomized Controlled Trial". In: *JAMA: The Journal of the American Medical Association* 307.7 (2012), pp. 685–692.

1.6.2 Data basics

1.3 Air pollution and birth outcomes, study components. Researchers collected data to examine the relationship between air pollutants and preterm births in Southern California. During the study air pollution levels were measured by air quality monitoring stations. Specifically, levels of carbon monoxide were recorded in parts per million, nitrogen dioxide and ozone in parts per hundred million, and coarse particulate matter (PM₁₀) in $\mu\text{g}/\text{m}^3$. Length of gestation data were collected on 143,196 births between the years 1989 and 1993, and air pollution exposure during gestation was calculated for each birth. The analysis suggested that increased ambient PM₁₀ and, to a lesser degree, CO concentrations may be associated with the occurrence of preterm births.³⁶ Identify

- (a) the cases,
- (b) the variables and their types, and
- (c) the main research question

in this study.

1.4 Buteyko method, study components. The Buteyko method is a shallow breathing technique developed by Konstantin Buteyko, a Russian doctor, in 1952. Anecdotal evidence suggests that the Buteyko method can reduce asthma symptoms and improve quality of life. In a scientific study to determine the effectiveness of this method, researchers recruited 600 asthma patients aged 18-69 who relied on medication for asthma treatment. These patients were split into two research groups: one practiced the Buteyko method and the other did not. Patients were scored on quality of life, activity, asthma symptoms, and medication reduction on a scale from 0 to 10. On average, the participants in the Buteyko group experienced a significant reduction in asthma symptoms and an improvement in quality of life.³⁷ Identify

- (a) the cases,
- (b) the variables and their types, and
- (c) the main research question

in this study.

1.5 Cheaters, study components. Researchers studying the relationship between honesty, age and self-control conducted an experiment on 160 children between the ages of 5 and 15. Participants reported their age, sex, and whether they were an only child or not. The researchers asked each child to toss a fair coin in private and to record the outcome (white or black) on a paper sheet, and said they would only reward children who report white. Half the students were explicitly told not to cheat and the others were not given any explicit instructions. In the no instruction group probability of cheating was found to be uniform across groups based on child's characteristics. In the group that was explicitly told to not cheat, girls were less likely to cheat, and while rate of cheating didn't vary by age for boys, it decreased with age for girls.³⁸ Identify

- (a) the cases,
- (b) the variables and their types, and
- (c) the main research question

in this study.

³⁶B. Ritz et al. "Effect of air pollution on preterm birth among children born in Southern California between 1989 and 1993". In: *Epidemiology* 11.5 (2000), pp. 502–511.

³⁷J. McGowan. "Health Education: Does the Buteyko Institute Method make a difference?" In: *Thorax* 58 (2003).

³⁸Alessandro Bucciol and Marco Piovesan. "Luck or cheating? A field experiment on honesty with children". In: *Journal of Economic Psychology* 32.1 (2011), pp. 73–78.

1.6 Stealers, study components. In a study of the relationship between socio-economic class and unethical behavior, 129 University of California undergraduates at Berkeley were asked to identify themselves as having low or high social-class by comparing themselves to others with the most (least) money, most (least) education, and most (least) respected jobs. They were also presented with a jar of individually wrapped candies and informed that the candies were for children in a nearby laboratory, but that they could take some if they wanted. After completing some unrelated tasks, participants reported the number of candies they had taken. It was found that those who were identified as upper-class took more candy than others.³⁹ Identify

- (a) the cases,
- (b) the variables and their types, and
- (c) the main research question

in this study.

1.7 Fisher's irises. Sir Ronald Aylmer Fisher was an English statistician, evolutionary biologist, and geneticist who worked on a data set that contained sepal length and width, and petal length and width from three species of iris flowers (*setosa*, *versicolor* and *virginica*). There were 50 flowers from each species in the data set.⁴⁰

- (a) How many cases were included in the data?
- (b) How many numerical variables are included in the data? Indicate what they are, and if they are continuous or discrete.
- (c) How many categorical variables are included in the data, and what are they? List the corresponding levels (categories).



Photo by Ryan Claussen
(<http://flic.kr/p/6QTcuX>)
CC BY-SA 2.0 license

1.8 Smoking habits of UK residents. A survey was conducted to study the smoking habits of UK residents. Below is a data matrix displaying a portion of the data collected in this survey. Note that “£” stands for British Pounds Sterling, “cig” stands for cigarettes, and “N/A” refers to a missing component of the data.⁴¹

	sex	age	marital	grossIncome	smoke	amtWeekends	amtWeekdays
1	Female	42	Single	Under £2,600	Yes	12 cig/day	12 cig/day
2	Male	44	Single	£10,400 to £15,600	No	N/A	N/A
3	Male	53	Married	Above £36,400	Yes	6 cig/day	6 cig/day
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
1691	Male	40	Single	£2,600 to £5,200	Yes	8 cig/day	8 cig/day

- (a) What does each row of the data matrix represent?
- (b) How many participants were included in the survey?
- (c) Indicate whether each variable in the study is numerical or categorical. If numerical, identify as continuous or discrete. If categorical, indicate if the variable is ordinal.

³⁹P.K. Piff et al. “Higher social class predicts increased unethical behavior”. In: *Proceedings of the National Academy of Sciences* (2012).

⁴⁰R.A Fisher. “The Use of Multiple Measurements in Taxonomic Problems”. In: *Annals of Eugenics* 7 (1936), pp. 179–188.

⁴¹National STEM Centre, Large Datasets from stats4schools.

1.6.3 Overview of data collection principles

1.9 Air pollution and birth outcomes, scope of inference. Exercise 1.3 introduces a study where researchers collected data to examine the relationship between air pollutants and preterm births in Southern California. During the study air pollution levels were measured by air quality monitoring stations. Length of gestation data were collected on 143,196 births between the years 1989 and 1993, and air pollution exposure during gestation was calculated for each birth.

- (a) Identify the population of interest and the sample in this study.
- (b) Comment on whether or not the results of the study can be generalized to the population, and if the findings of the study can be used to establish causal relationships.

1.10 Cheaters, scope of inference. Exercise 1.5 introduces a study where researchers studying the relationship between honesty, age, and self-control conducted an experiment on 160 children between the ages of 5 and 15. The researchers asked each child to toss a fair coin in private and to record the outcome (white or black) on a paper sheet, and said they would only reward children who report white. Half the students were explicitly told not to cheat and the others were not given any explicit instructions. Differences were observed in the cheating rates in the instruction and no instruction groups, as well as some differences across children's characteristics within each group.

- (a) Identify the population of interest and the sample in this study.
- (b) Comment on whether or not the results of the study can be generalized to the population, and if the findings of the study can be used to establish causal relationships.

1.11 Buteyko method, scope of inference. Exercise 1.4 introduces a study on using the Buteyko shallow breathing technique to reduce asthma symptoms and improve quality of life. As part of this study 600 asthma patients aged 18-69 who relied on medication for asthma treatment were recruited and randomly assigned to two groups: one practiced the Buteyko method and the other did not. Those in the Buteyko group experienced, on average, a significant reduction in asthma symptoms and an improvement in quality of life.

- (a) Identify the population of interest and the sample in this study.
- (b) Comment on whether or not the results of the study can be generalized to the population, and if the findings of the study can be used to establish causal relationships.

1.12 Stealers, scope of inference. Exercise 1.6 introduces a study on the relationship between socio-economic class and unethical behavior. As part of this study 129 University of California Berkeley undergraduates were asked to identify themselves as having low or high social-class by comparing themselves to others with the most (least) money, most (least) education, and most (least) respected jobs. They were also presented with a jar of individually wrapped candies and informed that the candies were for children in a nearby laboratory, but that they could take some if they wanted. After completing some unrelated tasks, participants reported the number of candies they had taken. It was found that those who were identified as upper-class took more candy than others.

- (a) Identify the population of interest and the sample in this study.
- (b) Comment on whether or not the results of the study can be generalized to the population, and if the findings of the study can be used to establish causal relationships.

1.13 Relaxing after work. The 2010 General Social Survey asked the question, “After an average work day, about how many hours do you have to relax or pursue activities that you enjoy?” to a random sample of 1,155 Americans. The average relaxing time was found to be 1.65 hours. Determine which of the following is an observation, a variable, a sample statistic, or a population parameter.


- (a) An American in the sample.
- (b) Number of hours spent relaxing after an average work day.
- (c) 1.65.
- (d) Average number of hours all Americans spend relaxing after an average work day.

1.14 Cats on YouTube. Suppose you want to estimate the percentage of videos on YouTube that are cat videos. It is impossible for you to watch all videos on YouTube so you use a random video picker to select 1000 videos for you. You find that 2% of these videos are cat videos. Determine which of the following is an observation, a variable, a sample statistic, or a population parameter.

- (a) Percentage of all videos on YouTube that are cat videos.
- (b) 2%.
- (c) A video in your sample.
- (d) Whether or not a video is a cat video.


1.15 GPA and study hours. A survey was conducted on 193 Duke University undergraduates who took an introductory statistics course in 2012. Among many other questions, this survey asked them about their GPA, which can range between 0 and 4 points, and the number of hours they spent studying per week. The scatterplot below displays the relationship between these two variables.

- (a) What is the explanatory variable and what is the response variable?
- (b) Describe the relationship between the two variables. Make sure to discuss unusual observations, if any.
- (c) Is this an experiment or an observational study?
- (d) Can we conclude that studying longer hours leads to higher GPAs?



1.16 Income and education in US counties. The scatterplot below shows the relationship between per capita income (in thousands of dollars) and percent of population with a bachelor's degree in 3,143 counties in the US in 2010.

- (a) What are the explanatory and response variables?
- (b) Describe the relationship between the two variables. Make sure to discuss unusual observations, if any.
- (c) Can we conclude that having a bachelor's degree increases one's income?



1.6.4 Observational studies and sampling strategies

1.17 Course satisfaction across sections. A large college class has 160 students. All 160 students attend the lectures together, but the students are divided into 4 groups, each of 40 students, for lab sections administered by different teaching assistants. The professor wants to conduct a survey about how satisfied the students are with the course, and he believes that the lab section a student is in might affect the student's overall satisfaction with the course.


- What type of study is this?
- Suggest a sampling strategy for carrying out this study.

1.18 Housing proposal across dorms. On a large college campus first-year students and sophomores live in dorms located on the eastern part of the campus and juniors and seniors live in dorms located on the western part of the campus. Suppose you want to collect student opinions on a new housing structure the college administration is proposing and you want to make sure your survey equally represents opinions from students from all years.

- What type of study is this?
- Suggest a sampling strategy for carrying out this study.

1.19 Internet use and life expectancy. The following scatterplot was created as part of a study evaluating the relationship between estimated life expectancy at birth (as of 2014) and percentage of internet users (as of 2009) in 208 countries for which such data were available.⁴²

- Describe the relationship between life expectancy and percentage of internet users.
- What type of study is this?
- State a possible confounding variable that might explain this relationship and describe its potential effect.



1.20 Stressed out, Part I. A study that surveyed a random sample of otherwise healthy high school students found that they are more likely to get muscle cramps when they are stressed. The study also noted that students drink more coffee and sleep less when they are stressed.

- What type of study is this?
- Can this study be used to conclude a causal relationship between increased stress and muscle cramps?
- State possible confounding variables that might explain the observed relationship between increased stress and muscle cramps.

1.21 Evaluate sampling methods. A university wants to determine what fraction of its undergraduate student body support a new \$25 annual fee to improve the student union. For each proposed method below, indicate whether the method is reasonable or not.

- Survey a simple random sample of 500 students.
- Stratify students by their field of study, then sample 10% of students from each stratum.
- Cluster students by their ages (e.g. 18 years old in one cluster, 19 years old in one cluster, etc.), then randomly sample three clusters and survey all students in those clusters.

⁴²CIA Factbook, Country Comparisons, 2014.

1.22 Random digit dialing. The Gallup Poll uses a procedure called random digit dialing, which creates phone numbers based on a list of all area codes in America in conjunction with the associated number of residential households in each area code. Give a possible reason the Gallup Poll chooses to use random digit dialing instead of picking phone numbers from the phone book.

1.23 Haters are gonna hate, study confirms. A study published in the *Journal of Personality and Social Psychology* asked a group of 200 randomly sampled men and women to evaluate how they felt about various subjects, such as camping, health care, architecture, taxidermy, crossword puzzles, and Japan in order to measure their dispositional attitude towards mostly independent stimuli. Then, they presented the participants with information about a new product: a microwave oven. This microwave oven does not exist, but the participants didn't know this, and were given three positive and three negative fake reviews. People who reacted positively to the subjects on the dispositional attitude measurement also tended to react positively to the microwave oven, and those who reacted negatively also tended to react negatively to it. Researcher concluded that "some people tend to like things, whereas others tend to dislike things, and a more thorough understanding of this tendency will lead to a more thorough understanding of the psychology of attitudes."⁴³

- (a) What are the cases?
- (b) What is (are) the response variable(s) in this study?
- (c) What is (are) the explanatory variable(s) in this study?
- (d) Does the study employ random sampling?
- (e) Is this an observational study or an experiment? Explain your reasoning.
- (f) Can we establish a causal link between the explanatory and response variables?
- (g) Can the results of the study be generalized to the population at large?

1.24 Family size. Suppose we want to estimate household size, where a "household" is defined as people living together in the same dwelling, and sharing living accommodations. If we select students at random at an elementary school and ask them what their family size is, will this be a good measure of household size? Or will our average be biased? If so, will it overestimate or underestimate the true value?

1.25 Flawed reasoning. Identify the flaw(s) in reasoning in the following scenarios. Explain what the individuals in the study should have done differently if they wanted to make such strong conclusions.

- (a) Students at an elementary school are given a questionnaire that they are asked to return after their parents have completed it. One of the questions asked is, "Do you find that your work schedule makes it difficult for you to spend time with your kids after school?" Of the parents who replied, 85% said "no". Based on these results, the school officials conclude that a great majority of the parents have no difficulty spending time with their kids after school.
- (b) A survey is conducted on a simple random sample of 1,000 women who recently gave birth, asking them about whether or not they smoked during pregnancy. A follow-up survey asking if the children have respiratory problems is conducted 3 years later, however, only 567 of these women are reached at the same address. The researcher reports that these 567 women are representative of all mothers.
- (c) A orthopedist administers a questionnaire to 30 of his patients who do not have any joint problems and finds that 20 of them regularly go running. He concludes that running decreases the risk of joint problems.

⁴³Justin Hepler and Dolores Albarracín. "Attitudes without objects - Evidence for a dispositional attitude, its measurement, and its consequences". In: *Journal of personality and social psychology* 104.6 (2013), p. 1060.

1.26 City council survey. A city council has requested a household survey be conducted in a suburban area of their city. The area is broken into many distinct and unique neighborhoods, some including large homes, some with only apartments, and others a diverse mixture of housing structures. Identify the sampling methods described below, and comment on whether or not you think they would be effective in this setting.

- (a) Randomly sample 50 households from the city.
- (b) Divide the city into neighborhoods, and sample 20 households from each neighborhood.
- (c) Divide the city into neighborhoods, randomly sample 10 neighborhoods, and sample all households from those neighborhoods.
- (d) Divide the city into neighborhoods, randomly sample 10 neighborhoods, and then randomly sample 20 households from those neighborhoods.
- (e) Sample the 200 households closest to the city council offices.

1.27 Sampling strategies. A statistics student who is curious about the relationship between the amount of time students spend on social networking sites and their performance at school decides to conduct a survey. Various research strategies for collecting data are described below. In each, name the sampling method proposed and any bias you might expect.

- (a) He randomly samples 40 students from the study's population, gives them the survey, asks them to fill it out and bring it back the next day.
- (b) He gives out the survey only to his friends, making sure each one of them fills out the survey.
- (c) He posts a link to an online survey on Facebook and asks his friends to fill out the survey.
- (d) He randomly samples 5 classes and asks a random sample of students from those classes to fill out the survey.

1.28 Reading the paper. Below are excerpts from two articles published in the *NY Times*:

- (a) An article titled *Risks: Smokers Found More Prone to Dementia* states the following:⁴⁴

"Researchers analyzed data from 23,123 health plan members who participated in a voluntary exam and health behavior survey from 1978 to 1985, when they were 50-60 years old. 23 years later, about 25% of the group had dementia, including 1,136 with Alzheimer's disease and 416 with vascular dementia. After adjusting for other factors, the researchers concluded that pack-a-day smokers were 37% more likely than nonsmokers to develop dementia, and the risks went up with increased smoking; 44% for one to two packs a day; and twice the risk for more than two packs."

Based on this study, can we conclude that smoking causes dementia later in life? Explain your reasoning.

- (b) Another article titled *The School Bully Is Sleepy* states the following:⁴⁵

"The University of Michigan study, collected survey data from parents on each child's sleep habits and asked both parents and teachers to assess behavioral concerns. About a third of the students studied were identified by parents or teachers as having problems with disruptive behavior or bullying. The researchers found that children who had behavioral issues and those who were identified as bullies were twice as likely to have shown symptoms of sleep disorders."

A friend of yours who read the article says, "The study shows that sleep disorders lead to bullying in school children." Is this statement justified? If not, how best can you describe the conclusion that can be drawn from this study?

⁴⁴R.C. Rabin. "Risks: Smokers Found More Prone to Dementia". In: *New York Times* (2010).

⁴⁵T. Parker-Pope. "The School Bully Is Sleepy". In: *New York Times* (2011).

1.29 Shyness on Facebook. Given the anonymity afforded to individuals in online interactions, researchers hypothesized that shy individuals might have more favorable attitudes toward Facebook, and that shyness might be positively correlated with time spent on Facebook. They also hypothesized that shy individuals might have fewer Facebook “friends” as they tend to have fewer friends than non-shy individuals have in the offline world. 103 undergraduate students at an Ontario university were surveyed via online questionnaires. The study states “Participants were recruited through the university’s psychology participation pool. After indicating an interest in the study, participants were sent an e-mail containing the study’s URL.” Are the results of this study generalizable to the population of all Facebook users?⁴⁶

1.6.5 Experiments

1.30 Stressed out, Part II. In a study evaluating the relationship between stress and muscle cramps half the subjects are randomly assigned to be exposed to increased stress by being placed into an elevator that falls rapidly and stops abruptly and the other half are left at no or baseline stress.

- (a) What type of study is this?
- (b) Can this study be used to conclude a causal relationship between increased stress and muscle cramps?

1.31 Light and exam performance. A study is designed to test the effect of light level on exam performance of students. The researcher believes that light levels might have different effects on males and females, so wants to make sure both are equally represented in each treatment. The treatments are fluorescent overhead lighting, yellow overhead lighting, no overhead lighting (only desk lamps).

- (a) What is the response variable?
- (b) What is the explanatory variable? What are its levels?
- (c) What is the blocking variable? What are its levels?

1.32 Vitamin supplements. In order to assess the effectiveness of taking large doses of vitamin C in reducing the duration of the common cold, researchers recruited 400 healthy volunteers from staff and students at a university. A quarter of the patients were assigned a placebo, and the rest were evenly divided between 1g Vitamin C, 3g Vitamin C, or 3g Vitamin C plus additives to be taken at onset of a cold for the following two days. All tablets had identical appearance and packaging. The nurses who handed the prescribed pills to the patients knew which patient received which treatment, but the researchers assessing the patients when they were sick did not. No significant differences were observed in any measure of cold duration or severity between the four medication groups, and the placebo group had the shortest duration of symptoms.⁴⁷

- (a) Was this an experiment or an observational study? Why?
- (b) What are the explanatory and response variables in this study?
- (c) Were the patients blinded to their treatment?
- (d) Was this study double-blind?
- (e) Participants are ultimately able to choose whether or not to use the pills prescribed to them. We might expect that not all of them will adhere and take their pills. Does this introduce a confounding variable to the study? Explain your reasoning.

⁴⁶E.S. Orr et al. “The influence of shyness on the use of Facebook in an undergraduate sample”. In: *CyberPsychology & Behavior* 12.3 (2009), pp. 337–340.

⁴⁷C. Audera et al. “Mega-dose vitamin C in treatment of the common cold: a randomised controlled trial”. In: *Medical Journal of Australia* 175.7 (2001), pp. 359–362.

1.33 Light, noise, and exam performance. A study is designed to test the effect of light level and noise level on exam performance of students. The researcher believes that light and noise levels might have different effects on males and females, so wants to make sure both are equally represented in each treatment. The light treatments considered are fluorescent overhead lighting, yellow overhead lighting, no overhead lighting (only desk lamps). The noise treatments considered are no noise, construction noise, and human chatter noise.

- (a) What type of study is this?
- (b) How many factors are considered in this study? Identify them, and describe their levels.
- (c) What is the role of the sex variable in this study?

1.34 Music and learning. You would like to conduct an experiment in class to see if students learn better if they study without any music, with music that has no lyrics (instrumental), or with music that has lyrics. Briefly outline a design for this study.

1.35 Soda preference. You would like to conduct an experiment in class to see if your classmates prefer the taste of regular Coke or Diet Coke. Briefly outline a design for this study.

1.36 Exercise and mental health. A researcher is interested in the effects of exercise on mental health and he proposes the following study: Use stratified random sampling to ensure representative proportions of 18-30, 31-40 and 41- 55 year olds from the population. Next, randomly assign half the subjects from each age group to exercise twice a week, and instruct the rest not to exercise. Conduct a mental health exam at the beginning and at the end of the study, and compare the results.

- (a) What type of study is this?
- (b) What are the treatment and control groups in this study?
- (c) Does this study make use of blocking? If so, what is the blocking variable?
- (d) Does this study make use of blinding?
- (e) Comment on whether or not the results of the study can be used to establish a causal relationship between exercise and mental health, and indicate whether or not the conclusions can be generalized to the population at large.
- (f) Suppose you are given the task of determining if this proposed study should get funding. Would you have any reservations about the study proposal?

1.37 Chia seeds and weight loss. Chia Pets – those terra-cotta figurines that sprout fuzzy green hair – made the chia plant a household name. But chia has gained an entirely new reputation as a diet supplement. In one 2009 study, a team of researchers recruited 38 men and divided them randomly into two groups: treatment or control. They also recruited 38 women, and they randomly placed half of these participants into the treatment group and the other half into the control group. One group was given 25 grams of chia seeds twice a day, and the other was given a placebo. The subjects volunteered to be a part of the study. After 12 weeks, the scientists found no significant difference between the groups in appetite or weight loss.⁴⁸


- (a) What type of study is this?
- (b) What are the experimental and control treatments in this study?
- (c) Has blocking been used in this study? If so, what is the blocking variable?
- (d) Has blinding been used in this study?
- (e) Comment on whether or not we can make a causal statement, and indicate whether or not we can generalize the conclusion to the population at large.

⁴⁸D.C. Nieman et al. “Chia seed does not promote weight loss or alter disease risk factors in overweight adults”. In: *Nutrition Research* 29.6 (2009), pp. 414–418.

1.6.6 Examining numerical data

1.38 Mammal life spans. Data were collected on life spans (in years) and gestation lengths (in days) for 62 mammals. A scatterplot of life span versus length of gestation is shown below.⁴⁹


- What type of an association is apparent between life span and length of gestation?
- What type of an association would you expect to see if the axes of the plot were reversed, i.e. if we plotted length of gestation versus life span?
- Are life span and length of gestation independent? Explain your reasoning.



1.39 Associations. Indicate which of the plots show a

- positive association
- negative association
- no association

Also determine if the positive and negative associations are linear or nonlinear. Each part may refer to more than one plot.



1.40 Office productivity. Office productivity is relatively low when the employees feel no stress about their work or job security. However, high levels of stress can also lead to reduced employee productivity. Sketch a plot to represent the relationship between stress and productivity.

1.41 Reproducing bacteria. Suppose that there is only sufficient space and nutrients to support one million bacterial cells in a petri dish. You place a few bacterial cells in this petri dish, allow them to reproduce freely, and record the number of bacterial cells in the dish over time. Sketch a plot representing the relationship between number of bacterial cells and time.

1.42 Sleeping in college. A recent article in a college newspaper stated that college students get an average of 5.5 hrs of sleep each night. A student who was skeptical about this value decided to conduct a survey by randomly sampling 25 students. On average, the sampled students slept 6.25 hours per night. Identify which value represents the sample mean and which value represents the claimed population mean.

1.43 Parameters and statistics. Identify which value represents the sample mean and which value

⁴⁹T. Allison and D.V. Cicchetti. "Sleep in mammals: ecological and constitutional correlates". In: *Arch. Hydrobiol* 75 (1975), p. 442.

represents the claimed population mean.

- (a) American households spent an average of about \$52 in 2007 on Halloween merchandise such as costumes, decorations and candy. To see if this number had changed, researchers conducted a new survey in 2008 before industry numbers were reported. The survey included 1,500 households and found that average Halloween spending was \$58 per household.
- (b) The average GPA of students in 2001 at a private university was 3.37. A survey on a sample of 203 students from this university yielded an average GPA of 3.59 in Spring semester of 2012.

1.44 Make-up exam. In a class of 25 students, 24 of them took an exam in class and 1 student took a make-up exam the following day. The professor graded the first batch of 24 exams and found an average score of 74 points with a standard deviation of 8.9 points. The student who took the make-up the following day scored 64 points on the exam.

- (a) Does the new student's score increase or decrease the average score?
- (b) What is the new average?
- (c) Does the new student's score increase or decrease the standard deviation of the scores?

1.45 Days off at a mining plant. Workers at a particular mining site receive an average of 35 days paid vacation, which is lower than the national average. The manager of this plant is under pressure from a local union to increase the amount of paid time off. However, he does not want to give more days off to the workers because that would be costly. Instead he decides he should fire 10 employees in such a way as to raise the average number of days off that are reported by his employees. In order to achieve this goal, should he fire employees who have the most number of days off, least number of days off, or those who have about the average number of days off?

1.46 Medians and IQRs. For each part, compare distributions (1) and (2) based on their medians and IQRs. You do not need to calculate these statistics; simply state how the medians and IQRs compare. Make sure to explain your reasoning.

- (a) (1) 3, 5, 6, 7, 9
(2) 3, 5, 6, 7, 20
- (b) (1) 3, 5, 6, 7, 9
(2) 3, 5, 8, 7, 9
- (c) (1) 1, 2, 3, 4, 5
(2) 6, 7, 8, 9, 10
- (d) (1) 0, 10, 50, 60, 100
(2) 0, 100, 500, 600, 1000

1.47 Means and SDs. For each part, compare distributions (1) and (2) based on their means and standard deviations. You do not need to calculate these statistics; simply state how the means and the standard deviations compare. Make sure to explain your reasoning. *Hint:* It may be useful to sketch dot plots of the distributions.

- (a) (1) 3, 5, 5, 5, 8, 11, 11, 11, 13
(2) 3, 5, 5, 5, 8, 11, 11, 11, 20
- (b) (1) -20, 0, 0, 0, 15, 25, 30, 30
(2) -40, 0, 0, 0, 15, 25, 30, 30
- (c) (1) 0, 2, 4, 6, 8, 10
(2) 20, 22, 24, 26, 28, 30
- (d) (1) 100, 200, 300, 400, 500
(2) 0, 50, 300, 550, 600

1.48 Stats scores. Below are the final exam scores of twenty introductory statistics students.


57, 66, 69, 71, 72, 73, 74, 77, 78, 78, 79, 79, 81, 81, 82, 83, 83, 88, 89, 94

Create a box plot of the distribution of these scores. The five number summary provided below may be useful.


Min	Q1	Q2 (Median)	Q3	Max
57	72.5	78.5	82.5	94

1.49 Infant mortality. The infant mortality rate is defined as the number of infant deaths per 1,000 live births. This rate is often used as an indicator of the level of health in a country. The relative frequency histogram below shows the distribution of estimated infant death rates for 224 countries for which such data were available in 2014.⁵⁰


- Estimate Q1, the median, and Q3 from the histogram.
- Would you expect the mean of this data set to be smaller or larger than the median? Explain your reasoning.




1.50 Mix-and-match. Describe the distribution in the histograms below and match them to the box plots.




(a)




(b)




(c)



(1)



(2)



(3)


1.51 Air quality. Daily air quality is measured by the air quality index (AQI) reported by the Environmental Protection Agency. This index reports the pollution level and what associated health effects might be a concern. The index is calculated for five major air pollutants regulated by the Clean Air Act and takes values from 0 to 300, where a higher value indicates lower air quality. AQI was reported for a sample of 91 days in 2011 in Durham, NC. The relative frequency histogram below

shows the distribution of the AQI values on these days.⁵¹


- Estimate the median AQI value of this sample.
- Would you expect the mean AQI value of this sample to be higher or lower than the median? Explain your reasoning.
- Estimate Q1, Q3, and IQR for the distribution.
- Would any of the days in this sample be considered to have an unusually low or high AQI? Explain your reasoning.

⁵⁰CIA Factbook, *Country Comparisons*, 2014.


⁵¹US Environmental Protection Agency, *AirData*, 2011.




1.52 Median vs. mean. Estimate the median for the 400 observations shown in the histogram, and note whether you expect the mean to be higher or lower than the median.




1.53 Histograms vs. box plots. Compare the two plots below. What characteristics of the distribution are apparent in the histogram and not in the box plot? What characteristics are apparent in the box plot but not in the histogram?




1.54 Marathon winners. The histogram and box plots below show the distribution of finishing times for male and female winners of the New York Marathon between 1970 and 1999.



- What features of the distribution are apparent in the histogram and not the box plot? What features are apparent in the box plot but not in the histogram?
- What may be the reason for the bimodal distribution? Explain.
- Compare the distribution of marathon times for men and women based on the box plot shown below.



- The time series plot shown below is another way to look at these data. Describe what is visible in this plot but not in the others.



1.55 Distributions and appropriate statistics, Part I. For each of the following, state whether you expect the distribution to be symmetric, right skewed, or left skewed. Also specify whether the mean or median would best represent a typical observation in the data, and whether the variability of observations would be best represented using the standard deviation or IQR. Explain your reasoning.

- (a) Number of pets per household.
- (b) Distance to work, i.e. number of miles between work and home.
- (c) Heights of adult males.

1.56 Distributions and appropriate statistics, Part II. For each of the following, state whether you expect the distribution to be symmetric, right skewed, or left skewed. Also specify whether the mean or median would best represent a typical observation in the data, and whether the variability of observations would be best represented using the standard deviation or IQR. Explain your reasoning.

- (a) Housing prices in a country where 25% of the houses cost below \$350,000, 50% of the houses cost below \$450,000, 75% of the houses cost below \$1,000,000 and there are a meaningful number of houses that cost more than \$6,000,000.
- (b) Housing prices in a country where 25% of the houses cost below \$300,000, 50% of the houses cost below \$600,000, 75% of the houses cost below \$900,000 and very few houses that cost more than \$1,200,000.
- (c) Number of alcoholic drinks consumed by college students in a given week. Assume that most of these students don't drink since they are under 21 years old, and only a few drink excessively.
- (d) Annual salaries of the employees at a Fortune 500 company where only a few high level executives earn much higher salaries than the all other employees.

1.57 TV watchers. Students in an AP Statistics class were asked how many hours of television they watch per week (including online streaming). This sample yielded an average of 4.71 hours, with a standard deviation of 4.18 hours. Is the distribution of number of hours students watch television weekly symmetric? If not, what shape would you expect this distribution to have? Explain your reasoning.

1.58 Exam scores. The average on a history exam (scored out of 100 points) was 85, with a standard deviation of 15. Is the distribution of the scores on this exam symmetric? If not, what shape would you expect this distribution to have? Explain your reasoning.



1.59 Facebook friends. Facebook data indicate that 50% of Facebook users have 100 or more friends, and that the average friend count of users is 190. What do these findings suggest about the shape of the distribution of number of friends of Facebook users?⁵²

1.60 A new statistic. The statistic $\frac{\bar{x}}{\text{median}}$ can be used as a measure of skewness. Suppose we have a distribution where all observations are greater than 0, $x_i > 0$. What is the expected shape of the distribution under the following conditions? Explain your reasoning.

- (a) $\frac{\bar{x}}{\text{median}} = 1$
- (b) $\frac{\bar{x}}{\text{median}} < 1$
- (c) $\frac{\bar{x}}{\text{median}} > 1$

⁵²Lars Backstrom. "Anatomy of Facebook". In: *Facebook Data Team's Notes* (2011).

1.61 Income at the coffee shop. The first histogram below shows the distribution of the yearly incomes of 40 patrons at a college coffee shop. Suppose two new people walk into the coffee shop: one making \$225,000 and the other \$250,000. The second histogram shows the new income distribution. Summary statistics are also provided.





	(1)	(2)
n	40	42
Min.	60,680	60,680
1st Qu.	63,620	63,710
Median	65,240	65,350
Mean	65,090	73,300
3rd Qu.	66,160	66,540
Max.	69,890	250,000
SD	2,122	37,321

- (a) Would the mean or the median best represent what we might think of as a typical income for the 42 patrons at this coffee shop? What does this say about the robustness of the two measures?
- (b) Would the standard deviation or the IQR best represent the amount of variability in the incomes of the 42 patrons at this coffee shop? What does this say about the robustness of the two measures?


1.62 Midrange. The *midrange* of a distribution is defined as the average of the maximum and the minimum of that distribution. Is this statistic robust to outliers and extreme skew? Explain your reasoning

1.63 Commute times. The US census collects data on time it takes Americans to commute to work, among many other variables. The histogram below shows the distribution of average commute times in 3,143 US counties in 2010. Also shown below is a spatial intensity map of the same data.



- (a) Describe the numerical distribution and comment on whether or not a log transformation may be advisable for these data.
- (b) Describe the spatial distribution of commuting times using the map below.


1.64 Hispanic population. The US census collects data on race and ethnicity of Americans, among many other variables. The histogram below shows the distribution of the percentage of the population that is Hispanic in 3,143 counties in the US in 2010. Also shown is a histogram of logs of these values.



- Describe the numerical distribution and comment on why we might want to use log-transformed values in analyzing or modeling these data.
- What features of the distribution of the Hispanic population in US counties are apparent in the map but not in the histogram? What features are apparent in the histogram but not the map?
- Is one visualization more appropriate or helpful than the other? Explain your reasoning.

1.6.7 Considering categorical data

1.65 Antibiotic use in children. The bar plot and the pie chart below show the distribution of pre-existing medical conditions of children involved in a study on the optimal duration of antibiotic use in treatment of tracheitis, which is an upper respiratory infection.




- What features are apparent in the bar plot but not in the pie chart?
- What features are apparent in the pie chart but not in the bar plot?
- Which graph would you prefer to use for displaying these categorical data?

1.66 Views on immigration. 910 randomly sampled registered voters from Tampa, FL were asked if they thought workers who have illegally entered the US should be (i) allowed to keep their jobs and apply for US citizenship, (ii) allowed to keep their jobs as temporary guest workers but not allowed to apply for US citizenship, or (iii) lose their jobs and have to leave the country. The results of the survey by political ideology are shown below.⁵³

	<i>Political ideology</i>			Total
	Conservative	Moderate	Liberal	
<i>Response</i> (i) Apply for citizenship	57	120	101	278
(ii) Guest worker	121	113	28	262
(iii) Leave the country	179	126	45	350
(iv) Not sure	15	4	1	20
Total	372	363	175	910

- What percent of these Tampa, FL voters identify themselves as conservatives?
- What percent of these Tampa, FL voters are in favor of the citizenship option?
- What percent of these Tampa, FL voters identify themselves as conservatives and are in favor of the citizenship option?
- What percent of these Tampa, FL voters who identify themselves as conservatives are also in favor of the citizenship option? What percent of moderates share this view? What percent of liberals share this view?
- Do political ideology and views on immigration appear to be independent? Explain your reasoning.


1.67 Views on the DREAM Act. A random sample of registered voters from Tampa, FL were asked if they support the DREAM Act, a proposed law which would provide a path to citizenship for people brought illegally to the US as children. The survey also collected information on the political ideology of the respondents. Based on the mosaic plot shown below, do views on the DREAM Act and political ideology appear to be independent? Explain your reasoning.⁵⁴



⁵³SurveyUSA, News Poll #18927, data collected Jan 27-29, 2012.

⁵⁴SurveyUSA, News Poll #18927, data collected Jan 27-29, 2012.

1.68 Raise taxes. A random sample of registered voters nationally were asked whether they think it's better to raise taxes on the rich or raise taxes on the poor. The survey also collected information on the political party affiliation of the respondents. Based on the mosaic plot shown below, do views on raising taxes and political affiliation appear to be independent? Explain your reasoning.⁵⁵



1.6.8 Case study: gender discrimination

1.69 Side effects of Avandia. Rosiglitazone is the active ingredient in the controversial type 2 diabetes medicine Avandia and has been linked to an increased risk of serious cardiovascular problems such as stroke, heart failure, and death. A common alternative treatment is pioglitazone, the active ingredient in a diabetes medicine called Actos. In a nationwide retrospective observational study of 227,571 Medicare beneficiaries aged 65 years or older, it was found that 2,593 of the 67,593 patients using rosiglitazone and 5,386 of the 159,978 using pioglitazone had serious cardiovascular problems. These data are summarized in the contingency table below.⁵⁶

		Cardiovascular problems		Total
		Yes	No	
Treatment	Rosiglitazone	2,593	65,000	67,593
	Pioglitazone	5,386	154,592	159,978
	Total	7,979	219,592	227,571

- (a) Determine if each of the following statements is true or false. If false, explain why. *Be careful:* The reasoning may be wrong even if the statement's conclusion is correct. In such cases, the statement should be considered false.
- Since more patients on pioglitazone had cardiovascular problems (5,386 vs. 2,593), we can conclude that the rate of cardiovascular problems for those on a pioglitazone treatment is higher.
 - The data suggest that diabetic patients who are taking rosiglitazone are more likely to have cardiovascular problems since the rate of incidence was $(2,593 / 67,593 = 0.038)$ 3.8% for patients on this treatment, while it was only $(5,386 / 159,978 = 0.034)$ 3.4% for patients on pioglitazone.
 - The fact that the rate of incidence is higher for the rosiglitazone group proves that rosiglitazone causes serious cardiovascular problems.
 - Based on the information provided so far, we cannot tell if the difference between the rates of incidences is due to a relationship between the two variables or due to chance.

(See the next page for additional parts to this question.)


- (b) What proportion of all patients had cardiovascular problems?

⁵⁵Public Policy Polling, *Americans on College Degrees, Classic Literature, the Seasons, and More*, data collected Feb 20-22, 2015.


⁵⁶D.J. Graham et al. "Risk of acute myocardial infarction, stroke, heart failure, and death in elderly Medicare patients treated with rosiglitazone or pioglitazone". In: *JAMA* 304.4 (2010), p. 411. issn: 0098-7484.

- (c) If the type of treatment and having cardiovascular problems were independent, about how many patients in the rosiglitazone group would we expect to have had cardiovascular problems?
- (d) We can investigate the relationship between outcome and treatment in this study using a randomization technique. While in reality we would carry out the simulations required for randomization using statistical software, suppose we actually simulate using index cards. In order to simulate from the independence model, which states that the outcomes were independent of the treatment, we write whether or not each patient had a cardiovascular problem on cards, shuffled all the cards together, then deal them into two groups of size 67,593 and 159,978. We repeat this simulation 1,000 times and each time record the number of people in the rosiglitazone group who had cardiovascular problems. Use the relative frequency histogram of these counts to answer (i)-(iii).

- i. What are the claims being tested?
- ii. Compared to the number calculated in part (b), which would provide more support for the alternative hypothesis, *more* or *fewer* patients with cardiovascular problems in the rosiglitazone group?
- iii. What do the simulation results suggest about the relationship between taking rosiglitazone and having cardiovascular problems in diabetic patients?



1.70 Heart transplants. The Stanford University Heart Transplant Study was conducted to determine whether an experimental heart transplant program increased lifespan. Each patient entering the program was designated an official heart transplant candidate, meaning that he was gravely ill and would most likely benefit from a new heart. Some patients got a transplant and some did not. The variable *transplant* indicates which group the patients were in; patients in the treatment group got a transplant and those in the control group did not. Another variable called *survived* was used to indicate whether or not the patient was alive at the end of the study.⁵⁷



- (a) Based on the mosaic plot, is survival independent of whether or not the patient got a transplant? Explain your reasoning.
(See the next page for additional parts to this question.)


⁵⁷B. Turnbull et al. "Survivorship of Heart Transplant Data". In: *Journal of the American Statistical Association* 69 (1974), pp. 74–80.

- (b) What do the box plots below suggest about the efficacy (effectiveness) of the heart transplant treatment.
- (c) What proportion of patients in the treatment group and what proportion of patients in the control group died?
- (d) One approach for investigating whether or not the treatment is effective is to use a randomization technique.

- i. What are the claims being tested?
- ii. The paragraph below describes the set up for such approach, if we were to do it without using statistical software. Fill in the blanks with a number or phrase, whichever is appropriate.

We write *alive* on _____ cards representing patients who were alive at the end of the study, and *dead* on _____ cards representing patients who were not. Then, we shuffle these cards and split them into two groups: one group of size _____ representing treatment, and another group of size _____ representing control. We calculate the difference between the proportion of *dead* cards in the treatment and control groups (treatment - control) and record this value. We repeat this 100 times to build a distribution centered at _____. Lastly, we calculate the fraction of simulations where the simulated differences in proportions are _____. If this fraction is low, we conclude that it is unlikely to have observed such an outcome by chance and that the null hypothesis should be rejected in favor of the alternative.

- iii. What do the simulation results shown below suggest about the effectiveness of the transplant program?



Index

- associated, 35
- bar plot, 42
 - segmented bar plot, 43
- blinded, 24
- blocking, 23
- blocks, 23
- boxplot, 34
 - side-by-side boxplot, 45
- categorical, 14
- cohort, 22
- column totals, 40
- confounder, 27
- confounding factor, 27
- confounding variable, 27
- contingency table, 40
 - column proportion, 41
 - row proportions, 40
- continuous, 14
- control, 23
- control group, 24
- convenience sample, 20
- data, 8
 - famuss, 13–14, 36–37, 40–46
 - frog, 12–13, 28–36, 38–39, 46
 - LEAP, 9–12, 23–24
- data density, 32
- data matrix, 13
- deviation, 29
- df, *see* degrees of freedom (df)
- discrete, 14
- distribution, 28
- dot plot, 31
- double-blind, 26
- experiment, 22, 23
- explanatory variable, 16
- factors, 14
- first quartile, 31
- frequency table, 40
- Greek
 - sigma (σ), 30
- histogram, 32
- hollow histogram, 45
- independent, 37
- interquartile range, 31
- levels, 14
- linearly associated, 35
- lurking variable, 27
- marginal totals, 40
- mean, 28
 - average, 28
- median, 29, 34
- modality
 - bimodal, 33
 - multimodal, 33
 - unimodal, 33
- mode, 33
- negatively associated, 35
- nominal, 14
- non-response, 21
- non-response bias, 21
- numerical, 14
- observational study, 22
- ordered categorical, 14
- outlier, 34
- patients, 24
- percentile, 31
- placebo, 22, 24
- placebo effect, 24
- population, 17, 17–22
- positively associated, 35
- probability sample, *see* sample
- prospective study, 27
- quartile
 - first quartile (Q_1), 31

- third quartile (Q_3), 31
- relative frequency table, 40
- relative risk, 11
- replication, 23
- representative, 21
- response variable, 16
- retrospective studies, 27
- robust estimates, 32
- row totals, 40
-
- s, 30
- sample, 17, 17–22
 - convenience sample, 20
 - non-response, 21
 - non-response bias, 21
 - random sample, 20–22
- scatterplot, 35
- side-by-side boxplot, 45
- simple random sample, 20
- skew
 - left skewed, 33
 - right skewed, 33
 - tail, 33
- standard deviation, 30
- stratification, 23
- study participants, 24
- summary statistic, 10
- symmetric, 33
- systematic sample, 21
-
- tail, 33
- third quartile, 31
- transformation, 38
- treatment group, 24
- two-way tables, 40
-
- variables, 12
- variance, 29
- volunteers, 24
-
- whiskers, 34