

AN OVERVIEW OF THE $3x + 1$ PROBLEM

MORITZ M. KONARSKI

ABSTRACT. This paper gives an overview of the Collatz function, a function that takes a positive integer as input and divides it by 2 if it is even or multiplies it by 3 and adds 1. The Collatz conjecture states that this function, when repeatedly applied to itself, will eventually yield 1. This conjecture has been computationally verified to be correct for very large numbers, but the proof is still outstanding. The background and reasons to study the conjecture will be discussed. Furthermore, details of the $3x + 1$ problem will be illustrated. Those details are trajectories, cycles, stopping time, and stochastic approximations.

CONTENTS

1. Introduction	2
1.1. The Collatz Conjecture	3
1.2. Background Information	3
1.3. Reasons to Study the $3x + 1$ Problem	3
2. The $3x + 1$ Problem in Detail	5
2.1. Trajectories	5
2.2. Cycles	5
2.3. Stopping Time	6
2.4. Stochastic Approximations	7
3. Examples	7
3.1. $3x + 1$ Function for $x = 39$	7
3.2. $3x + 1$ Function for $x = 27$	9
3.3. $3x + 1$ Function for $x = 489$	9
4. Conclusion	10
References	10

1. INTRODUCTION

The $3x + 1$ Problem is a number theoretical problem that has been studied by mathematicians since the 1950s, but still today it remains unsolved [3]. This problem is defined in terms of the *Collatz function*, which, for any positive integer x , returns $x/2$ if x is even and $3x + 1$ if x is odd. Mathematicians study the behavior of this function when it is repeatedly iterated, meaning it is applied to itself. Following Lagarias [3], the Collatz function is defined as

$$C(x) = \begin{cases} 3x + 1 & \text{if } x \equiv 1 \pmod{2}, \\ x/2 & \text{if } x \equiv 0 \pmod{2}. \end{cases} \quad (1)$$

When the $3x + 1$ Problem is studied, the $3x + 1$ *function*

$$T(x) = \begin{cases} (3x + 1)/2 & \text{if } x \equiv 1 \pmod{2}, \\ x/2 & \text{if } x \equiv 0 \pmod{2}. \end{cases} \quad (2)$$

The $3x + 1$ function $T(x)$ takes integers as input and yields integers as outputs, making it a number theoretic function like $C(x)$. More specifically, its domain are all positive integers and its range are also the positive integers. As such, $T(x)$ can be seen as the mapping of $\mathbb{N} + 1 \rightarrow \mathbb{N} + 1$, where $\mathbb{N} + 1$ represents the positive integers [4]. The $3x + 1$ function is, like the Collatz function, repeatedly applied to itself and has a *stopping time*, *total stopping time*, and a *trajectory* for each $x \in \mathbb{N} + 1$.

The $3x + 1$ function is used instead of the Collatz function because it is more predictable [3]. If x is odd, $3x + 1$ yields an even number that number is then immediately divided by 2. This has the effect that an otherwise following iteration of $C(x)$ is not needed. Furthermore, the function cannot grow as quickly because the factor is now $3/2$ instead of 3. These two changes make the function easier to study [3]. Accordingly, $T(x)$ can alternatively be defined in terms of equation (1) as $C(x)$ if x is even and as $C(C(x))$ if x is odd:

$$T(x) = \begin{cases} C(C(x)) & \text{if } x \equiv 1 \pmod{2}, \\ C(x) & \text{if } x \equiv 0 \pmod{2}. \end{cases}$$

1.1. The Collatz Conjecture. *For all $x \in \mathbb{N} + 1$ there is a $k \in \mathbb{N} + 1$ such that $T^{(k)}(x) = 1$. For any positive integer x , k iterations of the $3x + 1$ function will yield the result 1.*

This conjecture has not yet been proven and remains of interest to mathematicians who continue to study it to this day [3]. According to Chamberland [2], because $T(x)$ is a number theoretical function, successive applications of $T(x)$ can have the following results:

- (1) reach 1, which is equivalent to entering the trivial cycle, $\{2, 1, 2, 1, \dots\}$
- (2) enter a non-trivial cycle that does not include 1, or
- (3) diverge to infinity and not enter any type of cycle.

The Collatz conjecture states that point (1) is the only possible result and happens regardless of the positive integer that the iteration starts with.

1.2. Background Information. The Collatz conjecture and function are named after the German mathematician Lothar Collatz. Collatz worked on problems similar to the $3x + 1$ problem in the early 1930s, according to Lagarias [3]. The problem is also known as Hasse's Algorithm, Kakutani's Problem, and Ulam's Problem after other scientists who studied it or similar problems. During the 1950s the problem was circulated in various mathematical circles and the first publication was written about it in 1971 [3].

Today, the conjecture has been verified for over 10^{20} numbers using the help of computers to check if iterations of the $3x + 1$ function eventually reach 1 [4]. A number that does not iterate to 1 has not yet been found.

One of the most recent papers on the $3x + 1$ problem, "*Almost All Orbits of the Collatz Map Attain Almost Bounded Values*" by Terence Tao [4], was published in September of 2019. He proved that almost all iterations of the Collatz function are bounded, bringing mathematics one step closer to proving the Collatz conjecture. Currently, the problem is being actively researched.

1.3. Reasons the Study the $3x + 1$ Problem. According to Lagarias [3], the fact that the problem is easy to state but hard to prove makes it an interesting challenge for mathematicians. The fact that the $3x + 1$ problem has remained unsolved after over 50 years despite much research being done about it underlines

the challenge.

One of the challenging aspects of this problem is the pseudo-random nature of the $T(x)$ function. The function is difficult to predict when it comes to how iterations of it will behave. Because mathematical proofs tend to rely on patterns, this pseudo-randomness makes a proof of the conjecture very challenging.

Another reason for studying the $3x + 1$ problem is that it can be studied as an iterative mapping of positive integers to positive integers. These kinds of mappings are currently a popular research topic, according to Lagarias [3].

Computer scientists also have an interest in the Collatz conjecture because the computational testing of its correctness is an important part of current research. Since the 1960s, computers have been used to verify if positive integers follow the Collatz conjecture. Because these calculations have verified the conjecture for over 10^{20} numbers [4], finding and optimizing computer algorithms is important. Furthermore, whether or not a given positive integer will iterate to 1 can be studied as a decision problem in computer science [3].

The study of prime numbers is also connected to the $3x + 1$ problem and the Collatz conjecture. If it were to be proven to be correct, the proof may further the understanding of prime factorization of integers. This is because the Collatz function, in a way, factors integers when dividing by 2. When odd numbers are multiplied by 3, another prime factor is introduced to the number. The addition of 1 then can drastically change the way an integer can be factored and accordingly this behavior is interesting to study [3].

Finally, when talking about this problem, mathematician Paul Erdős said that

Mathematics is not ready for such problems.

If he is right and mathematics is indeed not ready for problems such as the $3x + 1$ problem, that could mean that new areas of mathematics are required to prove or disprove it. A proof of the Collatz conjecture may thus involve new areas of mathematics or maybe require old areas to be applied in new and different ways. Whatever the case may be, the $3x + 1$ problem is an interesting and challenging problem in mathematics and has connections to multiple other areas of mathematics and computer science. If it were to be proven or disproven, the effects could have wide-ranging implications [3].

2. THE $3x + 1$ PROBLEM IN DETAIL

This section will explain five main attributes connected to the $3x + 1$ problem and the study of the Collatz conjecture. First, trajectories of $T(x)$ will be considered. Second, cycles of $T(x)$ will be considered. Then, the stopping time and total stopping time of the $3x + 1$ function are considered. Finally, stochastic approximations of this function are explained.

2.1. Trajectories. The trajectory of x under $T(x)$ is the set of the successive iterations of $T(x)$ that starts with x and ends when $T^{(k)}(x) = 1$. This means that the trivial cycle $\{2, 1, 2, \dots\}$ is not included. This makes sense because otherwise all trajectories would be infinite in length. Accordingly, the trajectory of $T(x)$ has the length $k + 1$ because the first element of the trajectory x has the iteration number $k = 0$. Trajectories are also called forward orbits $O^+(x)$ of x under $T(x)$. Following Chamberland [2], the trajectory of x under $T(x)$ is defined as

$$O^+(x) := \{x, T(x), T^{(2)}(x), T^{(3)}(x), \dots\}$$

These trajectories contain all the numbers that the iteration of $T(x)$ yields. They can be graphed to illustrate what $T(x)$ does when repeatedly iterated. Looking at graphs of trajectories illustrates why $T(x)$ is considered a pseudo-random function, its erratic behavior is evident [3].

2.2. Cycles. Cycles of $T(x)$ are sequences of iterations of $T(x)$ that periodically repeat. It is known that $T(x)$ has the trivial cycle $\{2, 1, 2, \dots\}$, which is equivalent to reaching 1. Because of this equivalence, the trivial cycle is not included in the trajectories because it never ends and the resulting values don't leave the cycle. The Collatz conjecture states that any initial positive integer x will, after k iterations of $T(x)$, enter this trivial cycle. This means that the trivial cycle is the only cycle that can exist if the Collatz conjecture is true. In case non-trivial cycles do exist, which would disprove the Collatz conjecture, it has been proven that those non-trivial cycles must be more than 10.4 billion numbers long.

The $3x + 1$ function can also be applied to all integers. When this is done, it results in three more cycles, which, together with the trivial cycle, are conjectured to be the only ones that exist for $x \in \mathbb{Z}$. These cycles start at -1, -5, and -17.

$$\begin{aligned}
O^+(-1) &= \{-1, -1, -1, -1\}, \\
O^+(-5) &= \{-5, -7, -10, -5\}, \\
O^+(-17) &= \{-17, -25, -37, -55, -82, -41, \\
&\quad -61, -91, -136, -68, -34, -17\}
\end{aligned}$$

2.3. Stopping Time. The stopping time of $T(x)$ is the number of iterations k of $T(x)$ that it takes for the result of an iteration to be less than x . This is called stopping time because as soon as $x > T^{(k)}(x)$ it is known that the iteration will reach 1 and *stop*. Additionally, reaching this stopping point means that we can stop to check if an iteration reaches one. To be able to apply this stopping time, all numbers up to and including $x - 1$ have to be verified to iterate to 1. Then, if $T^{(k)}(x) < x$, it is obvious that $T(x)$ will also iterate to 1. The reason for that is that the trajectory of $T(x)$ is always the same and it contains some trajectories of other numbers. If we know that one of the numbers in the trajectory of $T(x)$ iterates to one, we know that $T(x)$ does so too. The stopping time of x is called $\sigma(x)$ and defined as

$$\sigma(x) = \inf\{k : T^{(k)}(x) < x\}.$$

Defining it like this makes sense because if one checks integers for compliance, it makes sense to start small and work towards larger integers. Furthermore, because, on average, the value of $T(x)$ decreases, this definition makes a lot of sense. This abbreviation makes it simple to check if a particular number iterates to 1 and saves a lot of computations that have already been performed. If the Collatz conjecture is true, all $x \in \mathbb{N} + 1$ have a finite stopping time, meaning they iterate to 1 eventually.

Related to the stopping time $\sigma(x)$ is the total stopping time $\sigma_\infty(x)$. This is the total number of iterations needed for $T(x)$ to iterate to 1. It is defined as

$$\sigma_\infty(x) = \inf\{k : T^{(k)}(x) = 1\}.$$

The total stopping time of $T(x)$ is conjectured to be finite for all $x \in \mathbb{N} + 1$ as a direct result of the Collatz conjecture.

2.4. Stochastic Approximations. When looking at the values that individual iterations of $T(x)$ yield, mathematicians noticed that for the trajectories of large numbers, the number of odd and even numbers is approximately the same. Furthermore, they observed that $T(x)$ behaves pseudo-randomly, meaning that it is hard to predict which value the function will take on in one or two iterations. Mathematicians use this property of $T(x)$ to make probabilistic statements about groups of trajectories and study their behaviors in this way. For example, the upper bound of for the total stopping time has been proven to be $41.677647 \log x$ [3].

One useful application of these stochastic approximations is the approximation of the total stopping time of $T(x)$. The total stopping time for most trajectories is approximated to be $6.95212 \log x$ steps.

3. EXAMPLES

3.1. $3x + 1$ Function for $x = 39$.

- trajectory
- graph
- stopping time
- total stopping time
- total stopping time approximation
- even and odd numbers counted

With the trajectory of $T(39)$ being

$$O^+(39) := \{39, 59, 89, 134, 67, 101, 152, 76, \mathbf{38}, 19, 29, \\ 44, 22, 11, 17, 26, 13, 20, 10, 5, 8, 4, 2, 1\},$$

we see that 38 is the first number < 39 . If we have previously shown that $T(38)$ iterates to 1, we can infer that $T(39)$ does so too. Thus $\sigma(39) = 8$, as 38 is the result of the 8th iteration.

3.1.1. *The trajectory of $T(39)$.* The trajectory of $T(39)$ is

$$O^+(39) := \{39, 59, 89, 134, 67, 101, 152, 76, 38, 19, 29, \\ 44, 22, 11, 17, 26, 13, 20, 10, 5, 8, 4, 2, 1\}$$

When graphed, one can see that $T^{(k)}(39)$ increases and decreases in a seemingly random manner.




FIGURE 1. The trajectory of $T^{(k)}(39)$ by number of iterations

For $T(39)$ we have the approximation

$$6.95212 \log 39 \approx 25.4952$$

Compared to the known $\sigma_\infty(39) = 23$ this is not bad.


For $T(39)$, which we considered above,

$$O^+(39) := \{39, 59, 89, 134, 67, 101, 152, 76, 38, 19, 29, \\ 44, 22, 11, 17, 26, 13, 20, 10, 5, 8, 4, 2, 1\}$$

and we see that $T^{(23)}(39) = 1$, this means that it takes 23 iterations of $T(x)$ to iterate to one. Accordingly, $\sigma_\infty(39) = 23$.

3.2. $3x + 1$ **Function** for $x = 27$.

- trajectory
- graph
- stopping time
- total stopping time
- total stopping time approximation
- even and odd numbers counted

FIGURE 2. The trajectory of $T^{(k)}(27)$ by number of iterations3.3. $3x + 1$ **Function** for $x = 489$.

- trajectory
- graph
- stopping time
- total stopping time
- total stopping time approximation
- even and odd numbers counted




FIGURE 3. The trajectory of $T^{(k)}(489)$ by number of iterations

4. CONCLUSION

- the Collatz Conjecture states that for $x, k \in \mathbb{N} + 1$ $T^{(k)}(x) = 1$
- the conjecture has not been proven, but verified for 10^{20} numbers
- pseudo-random nature
- relation to other fields of maths
- all orbits of $T(x)$ should reach the trivial cycle
- $T(x)$ can be probabilistically described because of pseudo-randomness

REFERENCES

- [1] Marc Chamberland, *An Update on the $3x + 1$ Problem*, http://www.math.grinnell.edu/~chamber1/papers/3x_survey_eng.pdf, 2005.
- [2] R. E. Crandall, *On the " $3x + 1$ " Problem*, *Mathematics of Computation*, **32** (1978), no. 144, 1281-1292.
- [3] Jeffrey C. Lagarias, *The $3x + 1$ Problem: An Overview*, <https://pdfs.semanticscholar.org/100046dd8b4ee901bc71043da7d42f5d87ca0224.pdf>, 2010.
- [4] Terence Tao, *Almost All Orbits of the Collatz Map Attain Almost Bounded Values*, arXiv:1909.03562v2 [math.PR], 2019.