

## EXERCISES 8

In Problems 1-2, verify that the following probability experiments represent hypergeometric probability experiments. Then determine the values of  $N$ ,  $n$ ,  $k$  and list the possible values of the random variable  $X$ .

1. In Michigan's Winfall Lottery, a player must choose 6 numbers between 1 and 49, inclusive. Six balls numbered between 1 and 49 are then randomly selected from an urn. The random variable  $X$  represents the number of matching numbers.
2. A manufacturer received an order of 250 computer chips. Unfortunately, 12 of the chips are defective. To test the shipment, the quality-control engineer randomly selects 20 chips from the box of 250 and tests them. The random variable  $X$  represents the number of defective chips in the sample of 20.

In Problems 3-4, a hypergeometric probability experiment is conducted with the given parameters. Compute the probability of obtaining  $x$  successes.

3.  $N = 150, n = 20, k = 30, x = 5$
4.  $N = 230, n = 15, k = 200, x = 12$

In Problems 5-6, compute the mean and standard deviation of the hypergeometric random variable  $X$ .

5.  $N = 150, n = 20, k = 30$
6.  $N = 230, n = 15, k = 200$

7. In Michigan's Classic Lotto 47 Lottery, a player must choose 6 numbers between 1 and 47, inclusive. Six balls numbered from 1 and 47 are then randomly selected from an urn. The random variable  $X$  represents the number of matching numbers.
  - (a) What is the probability of matching 3 numbers?
  - (b) What is the probability of matching 4 numbers?
  - (c) What is the probability of matching 5 numbers?
  - (d) What is the probability of matching 6 numbers?
  - (e) A winning ticket is one in which the player matches 3, 4, 5, or 6 numbers. What is the probability of purchasing a winning ticket? Would it be unusual to purchase a winning ticket?
  - (f) What is the mean and standard deviation of the random variable  $X$ ? For a randomly selected ticket, how many numbers do you expect to match?
8. A manufacturer received an order of 250 computer chips. Unfortunately, 12 of the chips are defective. To test the shipment, the quality-control engineer randomly selects 20 chips from the box of 250 and tests them. The random variable  $X$  represents the number of defective chips in the sample.
  - (a) What is the probability of obtaining 4 defective chips?
  - (b) What is the probability of obtaining 3 defective chips?
  - (c) What is the probability that the quality-control engineer will not find any defective chips?
  - (d) What is the probability of obtaining 14 defective chips?
  - (e) How many defective chips would you expect to select?
9. A hung jury is one that is unable to come to a unanimous decision regarding the guilt of the defendant. Suppose that there is a pool of 30 potential jurors, but 2 of the 30 potential jurors would never be willing to convict, regardless of the evidence presented. What is the probability that the trial will result in a hung jury, regardless of the evidence, if the jury consists of 12 randomly selected jurors?
10. Suppose that a concrete manufacturer has made 200 concrete cylinders that are supposed to withstand 4,000 pounds per square inch of pressure. As the quality-control manager, you decide to randomly test 4 of the cylinders to be sure they are manufactured to specification. You will only accept the shipment if all 4 cylinders pass the inspection. What is the probability that the shipment is accepted:
  - (a) If 10% of the 200 cylinders are defective?
  - (b) If 20% of the 200 cylinders are defective?
  - (c) If 40% of the 200 cylinders are defective?
  - (d) If 60% of the 200 cylinders are defective?
  - (e) If 80% of the 200 cylinders are defective?
  - (f) Draw a horizontal axis and label it Percent Defective.

Draw a vertical axis and label it Probability Accept Shipment. Plot probability accept shipment against the percent defective and connect the points in a smooth curve. This curve is referred to as an **operating characteristic curve**.

## Answers

1.  $N = 49, n = 6, k = 6, X = 0, 1, 2, \dots, 6$
2.  $N = 250, n = 20, k = 12, X = 0, 1, 2, \dots, 12$
3. 0.1856
4. 0.1939
5.  $\mu_X = 4, \sigma_X = 1.67$
6.  $\mu_X = 13.0, \sigma_X = 1.26$
7. (a) 0.01986  
(b) 0.001146  
(c) 0.0000229  
(d) 0.000000093  
(e) 0.02103  
(f) 0.766; 0.772; 0.766
8. (a) 0.0087  
(b) 0.0507  
(c) 0.3590  
(d) 0  
(e) 0.96
9. 0.6483
10. (a) 0.6539  
(b) 0.4065  
(c) 0.1270  
(d) 0.0245  
(e) 0.0014  
(f)

