

# An Overview of the $3x + 1$ Problem

Moritz M. Konarski

Applied Mathematics Department,  
American University of Central Asia

March 3rd, 2020

## Introduction

The  $3x + 1$  Function

Background Information

## $3x + 1$ Problem in Detail

Trajectories

Cycles

Stopping Time

Stochastic Approximations

## Summary

### Introduction

Function

Background

### Details

Trajectories

Cycles

Stopping Time

Approximations

### Summary

Introduction

Function

Background

Details

Trajectories

Cycles

Stopping Time

Approximations

Summary

# Introduction

The  $3x + 1$  Problem is based on the **Collatz Function**

$$C(x) = \begin{cases} 3x + 1 & \text{if } x \equiv 1 \pmod{2}, \\ x/2 & \text{if } x \equiv 0 \pmod{2}. \end{cases}$$

When the  $3x + 1$  Problem is studied, the  $3x + 1$  **Function**

$$T(x) = \begin{cases} (3x + 1)/2 & \text{if } x \equiv 1 \pmod{2}, \\ x/2 & \text{if } x \equiv 0 \pmod{2}. \end{cases}$$

is used.

- ▶  $T(x)$  is a function in **number theory**
- ▶ domain of  $T(x)$  are positive integers, its range are positive integers
- ▶ mathematically,  $T(x)$  maps  $\mathbb{N} + 1 \rightarrow \mathbb{N} + 1$
- ▶  $T(x)$  has a **stopping time**, **total stopping time**, and **trajectory** for each  $x \in \mathbb{N} + 1$
- ▶  $T(x)$  is repeatedly applied to an initial  $x$

## Conjecture

For all  $x \in \mathbb{N} + 1$  there is a  $k \in \mathbb{N} + 1$  such that  $T^{(k)}(x) = 1$ .

- ▶ starting at any positive integer  $x$ ,  $k$  iterations of  $T(x)$  will give the result 1
- ▶ the Collatz Conjecture has **not been proven**

$T(x)$  can:

1. reach 1, which is equivalent to entering the **trivial cycle**  $\{2, 1, 2, 1, \dots\}$
2. enter a non-trivial cycle that does not include 1
3. diverge to infinity and not enter any type of cycle

The Collatz Conjecture states that **1. always happens**.

- ▶ named after German mathematician Lothar Collatz
- ▶ problem circulated since the 1950s
- ▶ academic publications started in the 1970s
- ▶ conjecture has been verified for over  $10^{20}$  numbers
- ▶ most recent progress was in September of 2019
- ▶ problem is still being actively researched



- ▶ problem is simple to state, but hard to prove
- ▶ remains unsolved after over 50 years of research
- ▶ iterative mappings are currently a popular research topic
- ▶ verifying large numbers is computationally interesting
- ▶ could yield results connected to prime factorization using 2 and 3

*Mathematics is not ready for such problems.*

— *Paul Erdős*

Introduction

Function

Background

Details

Trajectories

Cycles

Stopping Time

Approximations

Summary

# Details

## Introduction

Function

Background

## Details

**Trajectories**

Cycles

Stopping Time

Approximations

## Summary

- ▶ the trajectory of  $x$  under  $T(x)$  is the set of successive iterations of  $T(x)$
- ▶ it is also called the forward orbit  $O^+(x)$  of  $x$  under  $T(x)$
- ▶ trajectories can be graphed

$$O^+(x) := \{x, T(x), T^{(2)}(x), T^{(3)}(x), \dots\}$$

# Example: Trajectory of $T(39)$


$3x + 1$  Problem

M. Konarski

The trajectory of  $T(39)$  is

$$O^+(39) := \{39, 59, 89, 134, 67, 101, 152, 76, 38, 19, 29, \\ 44, 22, 11, 17, 26, 13, 20, 10, 5, 8, 4, 2, 1\}$$

and can be graphed like this for  $k$  and  $T^{(k)}(39)$



[Introduction](#)

[Function](#)

[Background](#)

[Details](#)

[Trajectories](#)

[Cycles](#)

[Stopping Time](#)

[Approximations](#)

[Summary](#)

- ▶  $T(x)$  has the **trivial cycle**  $\{2, 1, 2, \dots\}$ , which is equivalent to reaching 1
- ▶ the Collatz Conjecture states that **all orbits will eventually enter the trivial cycle** and thus that **it is the only cycle**
- ▶ if  $T(x)$  has non-trivial cycles, they have been proven to be over 10.4 billion numbers long

## Introduction

Function

Background

## Details

Trajectories

Cycles

**Stopping Time**

Approximations

## Summary

- ▶ the number of iterations of  $T(x)$  until the result is smaller than  $x$
- ▶ first it is checked that every positive integer up to  $x - 1$  iterates to 1
- ▶ then, if  $T^{(k)}(x) < x$ , we know it will iterate to 1
- ▶ if the Collatz Conjecture is true, all  $x \in \mathbb{N} + 1$  have a finite stopping time

$$\sigma(x) = \inf\{k : T^{(k)}(x) < x\}$$

## Example: Stopping time of $T(39)$

$3x + 1$  Problem

M. Konarski

Introduction

Function

Background

Details

Trajectories

Cycles

**Stopping Time**

Approximations

Summary

With the trajectory of  $T(39)$

$$O^+(39) := \{39, 59, 89, 134, 67, 101, 152, 76, \mathbf{38}, 19, 29, \\ 44, 22, 11, 17, 26, 13, 20, 10, 5, 8, 4, 2, 1\},$$

we see that 38 is the first number  $< 39$ .

Thus  $\sigma(39) = 8$ , as 38 is the result of the 8th iteration.

The total stopping time is the number of steps needed for  $T(x)$  to iterate to 1. It is defined as

$$\sigma_{\infty}(x) = \inf\{k : T^{(k)}(x) = 1\}$$

## Example for $\sigma_{\infty}(39)$

For  $T(39)$ ,

$$O^+(39) := \{39, 59, 89, 134, 67, 101, 152, 76, 38, 19, 29, \\ 44, 22, 11, 17, 26, 13, 20, 10, 5, 8, 4, 2, 1\}$$

and we see that  $T^{(23)}(39) = 1$ , so  $\sigma_{\infty}(39) = 23$ .



## Introduction

Function

Background

## Details

Trajectories

Cycles

Stopping Time

Approximations

## Summary

- ▶ each trajectory has approximately the same number of odd and even elements
- ▶ the behavior of  $T(x)$  is pseudorandom for large numbers
- ▶ thus, probabilistic models describe its behavior
- ▶ these models describe groups of trajectories
- ▶ e.g., the upper bound for  $\sigma_\infty$  is  $41.677647 \log x$

# Example: Stopping Time Approximations

$3x + 1$  Problem

M. Konarski

Introduction

Function

Background

Details

Trajectories

Cycles

Stopping Time

Approximations

Summary

The total stopping time for most trajectories is approximated to be about  $6.95212 \log x$  steps.

Example for  $T(39)$

For  $T(39)$  we have the approximation

$$6.95212 \log 39 \approx 25.4952$$

Compared to the known  $\sigma_{\infty}(39) = 23$  this is not bad.

Introduction

Function

Background

Details

Trajectories

Cycles

Stopping Time

Approximations

Summary

# Summary

# Summary

$3x + 1$  Problem

M. Konarski

## Introduction

Function

Background

## Details

Trajectories

Cycles

Stopping Time

Approximations

## Summary



Marc Chamberland, *An Update on the  $3x + 1$  Problem*,  
[http://www.math.grinnell.edu/~chamberl/papers/3x\\_survey\\_eng.pdf](http://www.math.grinnell.edu/~chamberl/papers/3x_survey_eng.pdf), 2005.



R. E. Crandall, *On the " $3x + 1$ " Problem*,  
Mathematics of Computation, **32** (1978), no. 144,  
1281-1292.



Jeffrey C. Lagarias, *The  $3x + 1$  Problem: An Overview*,  
<https://pdfs.semanticscholar.org/100046dd8b4ee901bc71043da7d42f5d87ca0224.pdf>, 2010.



Terence Tao, *Almost All Orbits of the Collatz Map Attain Almost Bounded Values*, arXiv:1909.03562v2 [math.PR], 2019.