Grouping strategies for assessing and managing

2 persistent and mobile substances

- 3 Parviel Chirsir*1, Emma H. Palm1, Sivani Baskaran2, Emma L. Schymanski*1, Zhanyun
- 4 Wang³, Raoul Wolf², Sarah E. Hale⁴ and Hans Peter H. Arp*^{2,5}
- ¹ Luxembourg Centre for Systems Biomedicine (LCSB), University of Luxembourg, 6 avenue du
- 6 Swing, 4367, Belvaux, Luxembourg
- 7 ² Department of Environmental Engineering, Norwegian Geotechnical Institute, NO-0806 Oslo,
- 8 Norway
- 9 ³ Technology and Society Laboratory, Empa Swiss Federal Laboratories for Materials Science
- and Technology, 9014 St. Gallen, Switzerland
- 11 ⁴ TZW: DVGW-Technologiezentrum Wasser (German Water Centre), Karlsruher Straße 84,
- 12 76139 Karlsruhe / Germany
- 13 ⁵ Department of Chemistry, Norwegian University of Science and Technology (NTNU), NO-7491
- 14 Trondheim, Norway
- 15 * Corresponding authors: P.C. parviel.chirsir@uni.lu, E.L.S. emma.schymanski@uni.lu and
- 16 H.P.H.A. Hans.Peter.Arp@ngi.no
- 17 Keywords: PMT, vPvM, transformation products, hazardous properties, regrettable substitution,
- substance grouping, chemical regulation, persistent, mobile, toxic
- 19 ORCIDs:
- 20 PC: 0000-0002-9932-8609, EHP: 0000-0001-7774-2510, SB: 0000-0001-6192-5072, ELS:0000-
- 21 0001-6868-8145, ZW:0000-0001-9914-7659, RW: 0000-0001-5971-8525, SEH: 0000-0002-
- 22 7743-9199, HPHA: 0000-0002-0747-8838

24 Abstract

Background: Persistent, Mobile and Toxic (PMT), or very Persistent and very Mobile (vPvM) substances are a wide class of chemicals that are recalcitrant to degradation, easily transported, and potentially harmful to humans and the environment. Due to their persistence and mobility, these substances are often widespread in the environment once emitted, particularly in water resources, causing increased challenges during water treatment processes. Some PMT/vPvM substances such as GenX and perfluorobutane sulfonic acid have been identified as substances of very high concern (SVHCs) under the European Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) regulation. With hundreds to thousands of potential PMT/vPvM substances yet to be assessed and managed, effective and efficient approaches that avoid a case-by-case assessment and prevent regrettable substitution are necessary to achieve the European Union's zero-pollution goal for a non-toxic environment by 2050.

Main: Substance grouping has helped global regulation of some highly hazardous chemicals, e.g., through the Montreal Protocol and the Stockholm Convention. This article explores the potential of grouping strategies for identifying, assessing and managing PMT/vPvM substances. The aim is to facilitate early identification of lesser-known or new substances that potentially meet PMT/vPvM criteria, prompt additional testing, avoid regrettable use or substitution, and integrate into existing risk management strategies. Thus, this article provides an overview of PMT/vPvM substances and reviews the definition of PMT/vPvM criteria and various lists of PMT/vPvM substances available. It covers the current definition of groups, compares the use of substance grouping for hazard assessment and regulation, and discusses the advantages and disadvantages of grouping substances for regulation. The article then explores strategies for grouping PMT/vPvM substances, including read-across, structural similarity and commonly retained moieties, as well as the potential application of these strategies using cheminformatics to predict P, M and T properties for selected examples.

Conclusion: Effective substance grouping can accelerate the assessment and management of PMT/vPvM substances, especially for substances that lack information. Advances to read-across methods and cheminformatics tools are needed to support efficient and effective chemical management, preventing broad entry of hazardous chemicals into the global market and favouring safer and more sustainable alternatives.

54 Background

In 2019 water pollution was estimated to cause 1.4 million premature deaths globally [1]. Improving water quality by reducing pollution is also defined as one of the tasks in the Sustainable Development Goals [2]. Meanwhile, the number of known chemicals that are in use is increasing dramatically. Over 350,000 chemicals and mixtures have been registered in the global market over the past 50 years [3], while the largest chemical databases contain over 100 million chemicals, with PubChem [4] and the Chemical Abstracts Service (CAS) registry containing 116 million [5] and 219 million [6] chemicals, respectively, as of January 2023. Chemical production and pollution are outpacing global assessment capacity, posing more risks to human health, wildlife, and the environment [7, 8]. In 2021, about 224.8 and 86.4 million tonnes of chemicals hazardous to human health and the environment, respectively, were consumed in the European Union (EU) [9]. Some of these chemicals are persistent, mobile and toxic (PMT) substances or very persistent and very mobile (vPvM) substances, collectively referred to as PMT/vPvM substances.

PMT/vPvM substances have been listed by the European Commission's Scientific Committee on Health, Environmental and Emerging Risks (SCHEER) as one of the 14 emerging problems that could impact human health or the environment [10]. Recently, the EU Chemicals, Labelling and Packaging (CLP) regulation (1272/2008) introduced the new hazard classes PMT and vPvM, as well as defined criteria for PMT/vPvM substances based on chemical properties [11], as presented below in Figure 1. These substances do not degrade in the environment over an appreciable timescale and are easily transported through water and aquatic ecosystems due to poor sorption to soil and sediments [12–14]. PMT/vPvM substances can be found in a wide range of applications and sources, including in industrial processes, fire-fighting foams, and consumer products such as food, cosmetics and furniture [15]. They can cause long-term harm to humans and the environment and are also costly and difficult to remove from drinking water [16-19]. PMT/vPvM substances have been suggested to have an equivalent level of concern as persistent, bioaccumulative and toxic (PBT) substances or very persistent and very bioaccumulative (vPvB) substances. GenX and perfluorobutane sulfonic acid (PFBS), both classified as PMT/vPvM substances by the German Environmental Agency (UBA) [20] (but not yet officially classified under the CLP Regulation), have been identified as substances of very high concern (SVHCs) under the European Registration, Evaluation, Authorisation and Restriction of Chemicals, REACH, regulation (EC 1907/2006) [12]. The commonality between PMT/vPvM and PBT/vPvB

substances is their elevated potential for long-lasting and long-range exposure when released to the environment in substantial quantities. Their difference is related to the exposure pathway, with PMT/vPvM substances likely to be first monitored in water resources where they are most likely to accumulate and PBT/vPvB substances being first monitored in the food supply, humans and living organisms. Though this was not stated within the CLP regulation [11], it may be assumed this was introduced due to concerns from enhanced exposure potential leading to unforeseen effects, be they odour, aesthetic, geophysical, economical or other than established toxicological criteria. As an example, REACH Annex 1, Section 0.10 refers to "particular effects, such as ozone depletion, photochemical ozone creation potential, strong odour and tainting, for which [...] the risks associated with such effects shall be assessed on a case-by-case basis". Considering these concerns, a key consideration for vP substances is to further check if they exhibit enhanced exposure-related properties (vB or vM), toxicological or other hazards, following the current regulatory approaches.

Furthermore, due to the high mobility of PMT/vPvM substances, many of them can break through artificial barriers in wastewater treatment plants, including granular activated carbon filtration and ultrafiltration systems, posing challenges for removal and remediation [12, 21]. These substances can be further transported through natural media such as soils, riverbanks, aquifers and groundwater, making them hard to contain and remove from the environment. As such, these substances are problematic in drinking water, and many are detected frequently in European surface waters [12]. For example, 1,4-dioxin was found in Bavarian surface waters [22] and melamine in the Netherlands, France and Belgium [23].

Approximately 2% of identifiable unique chemicals registered under REACH are considered to meet the PMT/vPvM criteria (259 out of 13405 unique, identifiable chemicals registered under REACH); however, the true number is likely higher due to data gaps related to persistent, mobility or toxicity, with a potential maximum of 28% of REACH registered substances (3677 out of 13405 substances) [24]. Developing and applying substance grouping strategies is one way to potentially identify and manage PMT/vPvM substances more effectively. There are two main motivations for a substance grouping approach. The first is to expedite hazard assessments related to the large number of substances being introduced to the global chemical market. The second is to avoid regrettable substitution caused by drop-in substitution [25], where one substance is replaced by another with similar hazardous properties and effects [26]. Thus, this paper aims to explore how grouping strategies can be generally developed and used to identify, assess and manage new, emerging and well-known PMT/vPvM substances. To provide context, the paper first provides an

overview of PMT/vPvM substances, including scoping the numbers of substances covered by existing definitions, reviewing previous successful grouping strategies, and determining the relevance of grouping strategies in the context of PMT/vPvM substances while exploring the future efforts required to achieve this effectively. The principle aim of this article is not specifically targeting the ECHA technical guidance for PMT/vPvM substance assessment, but rather it is on how grouping strategies can be used to screen, assess, regulate and manage groups of PMT/vPvM substances more efficiently.

Methods

Definition of key terms

The International Union of Pure and Applied Chemistry (IUPAC) defines a chemical substance as "matter of constant composition best characterized by the entities (molecules, formula, units, atoms) it is composed of. Physical properties such as density, refractive index, electric conductivity, melting point etc. characterize the chemical substance" [27]. The term "substance" in this article is used in this context. Databases also refer to chemical entities as compounds and substances, but the context may be different. PubChem, for instance, define these as "a substance is a chemical sample description provided by a single source and a compound is a normalized chemical structure representation found in one or more contributed substances" [28]. Since this article later refers to calculations performed on PubChem queries, "compound" in this article refers to a chemical that fulfils the definition of a compound according to PubChem with a unique PubChem Compound Identifier (CID). A mixture, according to IUPAC, is a "portion of matter consisting of two or more chemical substances called constituents" [29]. Mixtures can be simple (e.g., xylene is a mixture of three isomers, o-xylene, m-xylene and p-xylene) or complex (e.g., C₉-C₁₄ alcohols, or mineral oils). The latter are often referred to as "substances of Unknown or Variable composition, Complex reaction products or Biological origin" (UVCBs) [30].

A group of substances is defined by REACH as substances that have "(1) a common functional group; (2) the common precursors and/or the likelihood of common breakdown products via physical and biological processes, which result in structurally similar chemicals; or (3) a constant pattern in the changing of the potency of the properties across the category." (REACH Regulation EC No 1907/2006, Annex XI, Section 1.5) [31]. In other words, the term "group" and "category"

are used interchangeably in this article. As an example, two widely regulated groups of substances that fulfil all three aspects of grouping substances in the REACH regulation are dioxins (polychlorinated dibenzo-p-dioxins and polychlorinated dibenzo-furans) [32] and polychlorinated biphenyls (PCBs) [33], which are composed of about 210 [34] and 209 individual congeners, respectively [35].

Literature analysis

To obtain relevant literature to underpin the arguments in this paper, a comprehensive literature search spanning both policy-related documents (e.g., regulations, guidelines, analysis, dossiers) and scientific literature was conducted using keywords like "persistent, mobile and toxic", "very persistent very mobile", "substance grouping" and "chemical regulation" in electronic databases such as PubMed and Google Scholar, while filtering for relevance. As the study focus area was mainly the European Union, recent policy documents in the EU and United Nations were considered most relevant due to the greater activity on PMT/vPvM substances, while detailed policy analysis of areas outside the European Union was considered out of scope. Scientific studies focused on grouping strategies for the assessment and management of substances, including chemicals, pollutants, and contaminants that exhibit persistence and mobility in the environment were also considered relevant. Backward citation searching was also conducted from the list of relevant articles to identify additional studies that may have been overlooked during the electronic search. By systematically executing the search strategy across multiple databases and supplementing it with hand-searching reference lists, a comprehensive selection of relevant literature with findings focused on the topic was compiled and considered in this article.

Cheminformatics analysis

For cheminformatics analysis, the search functions available in the PubChem database [4] were used to obtain the structural and substructural data of compounds of interest. The OPERA ReadyBiodegradable model version 2.9 [36] was used to predict the persistence of compounds, along with the KOCWIN model version 2.00 [37] to predict the mobility (log K_{oc} values) of the compounds from the PubChemLite for Exposomics dataset, Version 1.27.0 [38] and other subsets of compounds extracted from PubChem. Further details of other datasets used can be found in the "availability of data and material" section below. The toxicity predictions were performed with MS2Tox version 0.3.2 [39]. Data processing was performed in R version 4.3.1 [40] using the

packages Tidyverse version 2.0.0 [41, 42] and ggplot2 version 3.4.3 [43, 44] for data transformation and plotting, and RChemMass 0.1.28 [45] for obtaining the exact mass before MS2Tox calculations. Additional information is given in the "Cheminformatics challenges in grouping PMT/vPvM substances" section below.

Overview of PMT/vPvM Substances

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

Assessment of the number of substances being introduced to the global chemical market that may be PMT/vPvM substances requires a clear set of criteria and the ability to scale the application of these criteria to large numbers. The EU Chemicals, Labelling and Packaging (CLP) regulation (EC 1272/2008) criteria for PMT/vPvM substances [11] are displayed in Figure 1. The mobility (M) aspect (see Figure 1A) is defined based on the organic-carbon-water partition coefficient (K_{OC} , with units of L/kg) of a chemical which is then log transformed to log K_{OC} values, this is used as a proxy to describe the sorption potential of a chemical to organic carbon. Since experimental log K_{OC} data are rare, it was suggested that the logarithmic octanol-water partition coefficient (log K_{OW} or log P) or the pH-adjusted log D_{OW} (or log D) could be used as a screening parameter for hydrophobicity where high-quality log K_{OC} data are not available [24, 46]. Though the two parameters have some variance, particularly for ionogenic substances, substances with very low log D_{OW} values tend to have very low log K_{OC} values, hence log D_{OW} values were seen as a good alternative screening parameter for mobility. As such they are often used in a complementary fashion (see [24, 47] for more details), where $\log K_{OW}$ and $\log D_{OW}$ data are used when log K_{OC} data are unavailable. **Persistence** (P) is defined based on the half-lives of a chemical in different environmental water, sediment and soil systems (see Figure 1B), and toxicity (T) is defined based on one or a combination of the following: carcinogenicity, mutagenicity, reproductive toxicity, endocrine disruption and specific target organ toxicity after repeated exposure as well as long-term no-observed effect concentration (NOEC) and effect concentration at a given percentage of the population (EC_x) shown in Figure 1C.

A compound is a considered a PMT substances when it fulfils the persistence, mobility and toxicity criteria (i.e. a combination of at least one persistency criterion, one mobility criterion and one toxicity criterion are met) or in the case of vPvM at least one of the criteria for very persistent and the very mobile criterion must be met (see Figure 1). Some examples of PMT/vPvM substances are shown in Figure 2.




Figure 1: EU CLP criteria [11] for PMT/vPvM substances. *A) Mobility criteria based on log K*_{OC} (blue). *B) Persistence criteria, with persistent in green and C) Toxicity criteria, including carcinogenicity, mutagenicity, endocrine disruption, specific target toxicity and reproductive toxicity (1A, 1B, 1 and 2 refer to the categories for that criterion) as well as concentrations, where long-term no observable effect concentration (NOEC), and effect concentration (EC_x) for marine and freshwater organisms refers to the concentration where a toxic effect was observed in a given percentage of the population e.g. 10% for EC₁₀ at less than 0.01 mg/L.*




Figure 2: Selected examples of potential PMT/vPvM substances from the UBAPMT list assessed by Arp and Hale (2022) as meeting the criteria [20, 24]. *A) PMT substances; B) vPvM substances; C) both PMT and vPvM substances.*

There are several published lists containing PMT/vPvM substances (e.g. the PMT list by Holmberg et al., UBAPMT, EAWAGPMT, UFZHSFPMT, ZEROPM and PMTPFAS) [20, 48–52]. The later five have been digitized and are hosted on the NORMAN Suspect List Exchange (NORMAN-SLE) [53, 54] and thus openly available with a CC-BY license. The UBAPMT suspect list contains substances currently registered under REACH that meet the proposed PMT/vPvM criteria set by UBA in 2019 [47]. The original list has 254 substances [55, 56], while the 2022 revised version has 340 substances [57, 58] due to an increase in chemicals registered in REACH and updated chemical information [20]. The EAWAGPMT list contains 1,156 compounds identified in groundwater by Kiefer et al. [49, 59]. This list contains a mix of compounds on other PMT lists as well as compounds on the Swedish Chemical Agency (KEMI) market list [60] that have a low log Dow and high water exposure index, while the UFZHSFPMT list includes 1,063 potential persistent mobile compounds detected in surface water as described by Neuwald et al. [50, 61, 62]. The ZEROPMBOX1 contains 38 compounds, including representative per- and polyfluoroalkyl substances (PFAS), triazines and triazoles used to start the H2020 ZeroPM project [51]. Finally, the PMTPFAS list contains 180 fluorinated compounds extracted from the UBAPMT, EAWAGPMT and UFZHSFPMT lists [52]. In combination, the five lists contain 2,081 unique compounds, but this does not represent all PMT/vPvM substances in the global market.

Grouping and Regulations

Assessing and regulating individual PMT/vPvM substances that fall within a PMT/vPvM substance group is inefficient and more time-consuming than assessing an entire group. Past strategies have led to drop-in replacements, subsequently referred to as regrettable substitution [25]. Assessing chemicals individually also ignores cumulative exposures and risks of groups of substances [63]. Considering primarily the structural similarity and similar properties of PMT/vPvM substances, it may be more feasible and prudent to regulate these substances as a group. The idea of substance grouping based on the relationship between hazard and structural similarity is not new, since many of the very first organic substances to be regulated were groups sharing a similar structure. The successes of regulation in managing substances as groups such as ozone-depleting substances (ODS) under the Montreal Protocol [64, 65] and specific groups of persistent organic pollutants (POPs) under the Stockholm Convention [66] are discussed below. Moreover, successfully grouping substances for regulation not only accelerates but also prevents and

reduces inconsistencies in the regulatory process [67]. Grouping substances can enhance chemical safety management, facilitating the identification of regulated substances and potential substitutes for harmful ones [67].

While there may be several legislations about grouping substances globally, this section focuses on the use of grouping in EU legislation. In the EU, the European Chemical Agency (ECHA) coordinates the REACH regulation for the restriction, evaluation and authorization of substances based on hazard classifications and the CLP regulation for labelling based on hazard classifications. REACH Annex XI section 1.5 specifies how groups or categories of substances can be defined and which regulatory actions can be applied [31]. It states that "substances whose physicochemical, toxicological and ecotoxicological properties are likely to be similar or follow a regular pattern as a result of structural similarity may be considered as a group, or 'category' of substances" [31]. The toxicological or hazard properties referred to are described in the CLP regulation shown in Figure 1.

According to Article 36 of the CLP regulation, certain substances, including those with respiratory sensitizing properties (category 1), germ cell mutagenicity, carcinogenicity, and reproductive toxicity (category 1A, 1B and 2), are considered hazardous and are subject to harmonized classification. This also applies to active substances listed in Regulation (EC) No 1107/2009 and Regulation (EU) No 528/2012 [68, 69]. Moreover, Article 57 of REACH includes substances that are hazardous to the environment and are also subject to harmonized classification. A list of these hazardous substances for which harmonized classification and labelling has been established at the EU level can be found in Part 3 of Annex VI of the CLP regulation.

Under REACH, UVCBs are regulated based on the structural similarity of the constituents identified and can also be regulated as part of a different group. This means that certain components of these substances may be identified and registered as part of a different group. Annex XI section 1.5 of REACH provides more information on the regulation of UVCBs [31, 70].

Existing grouping legislation and impacts

The Montreal Protocol on Ozone-Depleting Substances

The Montreal Protocol regulates the manufacturing and consumption of ODS [64, 71]. This is a group of over 100 substances including chlorofluorocarbons (CFCs), methyl chloroform, hydrochlorofluorocarbons (HCFCs) and hydrobromofluorocarbons (HBFCs) that release chlorine and bromine into the stratosphere, damaging the ozone layer [64, 72]. This can lead to increased global warming, skin cancer and damage to marine ecosystems [73]. The Montreal Protocol is considered to have drastically decreased relative consumption (total of production and imports total of exports and destroyed) of ODS in the EU (100% to 0.36%) and globally (100% to 1.35%) between 1986 and 2021 [71, 74], as shown in Figure 3. This is an example of a property-based regulatory effort that has been successful [71, 75]. However, the Montreal Protocol has also been criticized for shifting the burden by transitioning from ODS to greenhouse gases and PMT/vPvM precursors, as discussed further below.




Figure 3: EU and global relative consumption of ODS since 1986 [74] showing the reduction of the consumption of ODS due to the Montreal Protocol.

The Stockholm Convention on Persistent Organic Pollutants

The Stockholm Convention was signed in 2001 to protect humans and the environment from POPs [66]. An EU Legislation regulation (EC No 850/2004) was adopted later to implement the Convention in the EU [76, 77]. Initially, the Convention regulated 12 POPs or groups of POPs

such as aldrin, chlordane and dichlorodiphenyltrichloroethane (DDT). Over time, the list expanded to include 39 substances and substance groups as of January 2024, with the regulation covering the manufacturing, sales, use, and waste management of these chemicals, as well as unintentional releases in some cases [66, 78]. Structural similarity and the resulting similar compound properties played a major role in managing the grouping of substances under the Convention, for instance, polychlorinated biphenyls (PCBs), dioxins and DDT derivatives [79]. The Stockholm Convention has been successful in reducing the emission of regulated POPs since its enforcement. The emission levels of PCBs, for example, decreased by over 60% from 2001 to 2020, while dioxin levels also dropped by 50% from 2001 to 2020 (top of Figure 4). This reduction is also noticeable as the number of registered patents for 17 dioxins, 12 PCBs and hexachlorobenzene (HCB) dropped since the Convention (bottom of Figure 4).




Figure 4: Top: Emission tonnage of PCB, dioxins and HCB from regulation (EU) No 277/2012, between 1990 to 2020 [80]. Bottom: The chemical stripes drawn using the chemical stripes package in R [81] show a decrease in the number of registered patents of PCB, dioxins and HCB in PubChem; the date range was chosen to match the emission data.

Substitution of regulated substances with PMT/vPvM substances

316

317

318

319

320

321

322

323

324

325

326

327

328 329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

The concept of regrettable substitution first appeared in the literature in 2011 [82]. There are many examples of regrettable substitution, which refers to the practice of replacing a hazardous substance with a structurally similar substance for a specific use or function, that is also hazardous or potentially more hazardous, or that has not been tested broadly enough for different hazards and other than those of the substance being replaced [83]. One well-known example is Bisphenol A, an endocrine-disrupting chemical (EDC) that was first discovered in 1891 [84]. Due to its harmful effects, it has since been replaced by various other EDCs with similar structures such as Bisphenol S and Bisphenol P. Currently, more than 30 bisphenols, including the ones mentioned above, are recommended for restriction by ECHA [85].

One of the reasons for the "drop-in" solution by companies after substance regulation (both as individual or as a group) is that structurally similar substances often have similar functions in the product being applied, and can be manufactured generally without requiring substantial changes in processes, infrastructure, or product testing [63]. Unfortunately, similar toxicological effects are likely to occur as well [26], and at other times "burden shifting" towards different hazards. A key example is the series of restrictions related to the Montreal Protocol that began with a global phase-out of CFCs [86, 87]. Later, hydrobromofluorocarbons (HBFCs) were introduced as replacements but were only briefly commercialized [88]. They were then replaced with hydrochlorofluorocarbons (HCFCs), which exhibited substantially less ozone-depleting potential, and exhibited the same commercial properties as CFCs [89]. HCFCs were eventually also included under the Montreal Protocol and were again replaced by chemicals with similar chemistry, hydrofluorocarbons (HFCs) [64]. Due to the removal of chlorine atoms from the molecules, HFCs no longer pose a threat to the stratospheric ozone layer [90], but they are still highly persistent and have high global warming potential. Therefore, in a subsequent amendment to the Montreal Protocol, the Kigali Amendment, HFCs themselves were listed [90, 91]. As a consequence of the Kigali amendment, the industry that manufactures fluorinated gases then sought to substitute HFCs with hydrofluoroolefins (HFOs), which have a short atmospheric lifetime of 6 days [92, 93]. However, these HFOs (almost) exclusively degrade to form trifluoroacetic acid (TFA), a vPvM substance that is ubiquitous in the environment, especially in drinking water systems, but the current understanding of its risk to human health and the environment is limited. While some research has argued that TFA has low health risks [94, 95], other studies have shown

mild liver hypertrophy in rats, and eye and skin irritation [94, 96]. As the levels of TFA are rapidly increasing, some EU countries have proposed to ban HFOs as part of the broad PFAS restriction under REACH. In summary, although the Montreal Protocol is often considered the most successful multilateral environmental agreement, this has led to burden shifting over time from ODS to the production of persistent, mobile substances such as TFA.

Similar to the Montreal Protocol, increased regulatory pressure on PBT/vPvB substances in Europe and the USA was seen as a driver for the chemical industry to produce more hydrophilic, and hence mobile, substances [97]. Individual fluoropolymer manufacturers have developed their own structurally similar per- and polyfluoroalkylether carboxylic acids (PFECAs) as replacements for PFOA, which was used as a processing aid in fluoropolymer production. However, as mentioned above, some of these replacements, including GenX, are similarly problematic despite being less bioaccumulative [98]. To date, many monitoring programs have seen an increase in these replacements in water resources and fish due to increasing use and emissions [99]. GenX and PFBS were identified as SVHCs under the EU REACH regulation in 2019 and were some of the first SVHCs that were considered an equivalent level of concern to PBT/vPvB substances [12]. Thus, the phasing out of some PBT/vPvB substances such as the long-chain PFASs has led to burden shifting towards PMT/vPvM substances such as PFECAs [100].

Existing regulation of substances as groups based on structural similarity and/or intrinsic properties has been successful, as demonstrated in the Montreal Protocol of ODSs and the Stockholm Convention of POPs. Meanwhile, regrettable substitution with increasing production of PMT/vPvM substances such as TFA and GenX has been noted as a side-effect. This shows a limitation of the grouping approach under the Montreal Protocol and the Stockholm Convention: limiting the scope of intrinsic properties during grouping may lead to burden shifting to other hazards. This calls for a grouping strategy that can prevent regrettable substitution and improve the efficiency and effectiveness of chemical regulatory processes.

Grouping strategies

Grouping based on structural similarity

Read-across in substance grouping is a strategy that relies on the use of important information obtained from tests conducted on a reference substance known as the "source" substance within a group to predict the properties of another substance in the group known as the "target" substance [101]. This method relies on justified similarities in structure, toxicokinetic, physicochemical and molecular properties, transformation process/endpoints and other similar data for interpolation and extrapolation of relevant information [102] and is recommended in Section 1.5 of Annex XI of the REACH regulation for groups of substances [70]. It can be applied from a single source substance to a single target substance (analogue approach) or from multiple source substances to multiple target substances within a group (category approach) [101, 103]. This method is commonly used to fill data gaps for chemical safety assessment in the regulatory process but has also been used to build grouping hypotheses for categories of chemicals [101, 102]. The Read-Across Assessment Framework (RAAF) by ECHA and the OECD guidance on the grouping of chemicals have been developed to guide systematic and consistent applications of read-across [104, 105].

Read-across is advantageous because it reduces the number of experimental tests needed during an assessment by using existing experimental data for the source substance(s) to predict the properties of the untested substance(s) [106]. If a clear hypothesis and justification are provided, read-across can be used efficiently to predict the hazard properties of target substances and fill data gaps in the regulatory process, thereby facilitating and speeding up assessment and regulatory decisions [107]. However, read-across requires an adequate justification for use, and appropriate documentation covering all assumptions and conclusions [36, 107]. This can be complex and may require a certain level of expertise for interpretation [108]. Despite its limitations, read-across can be applied to the grouping of PMT/vPvM substances.

For persistence in the environment, an important property is the readiness of the bonds to be broken down under ambient conditions, whether it be from radical reactions or through metabolic processes. For instance, the C-F bonds in PFAS (bond dissociation energy of 513.8 \pm 10.0 kJ/mol), and aromatic-Cl bonds in PCBs or dioxins (394.9 \pm 13.4 kJ/mol), are difficult to break down in the environment, making groups rich in such bonds likely to be persistent [109].

Compounds with molecules containing high bond dissociation energies can be flagged for persistence assessment and subsequent inclusion as PMT/vPvM if each of the M and/or T criteria are met. Biodegradability models can be used to quickly provide relevant information about the persistence of many substances or the presence of many persistent substructures, pending further investigation and confirmation by experimental studies [110]. However, the applicability domain of available models is dependent on their training data, resulting in unreliable predictions for compound classes that may not be covered yet [110].

For mobility, the chemical substructures that are associated with low K_{OC} values are those that are highly polar or ionic, as this favours their water solubility over sorption to soil organic carbon [24, 111]. A 2022 review found that most ionic compounds with measured log K_{OC} values have log K_{OC} values <4.0 [24]. Thus, the presence of many hydrophilic substituents may be a predictor of mobility, and conversely, chemicals with largely hydrophobic substructures are unlikely to be mobile in the environment. However, an important limitation of mobility is size. Extremely large molecules that are highly polar and ionic may not be mobile if they have sufficient hydrophobic substructures to decrease their solubility, or (as in the case of water-soluble polymers) can aggregate for charge neutrality and therefore lose/reduce their mobility as an aggregate [112]. Quantitative structure-activity relationship (QSAR) models are available to predict the mobility of certain substances and can be applied effectively (within their respective applicability domains) to classify the mobility of substances.

The toxicity of a substance is related to its chemical structure and determines the type of health effects it induces in the biological systems [113]. The interaction of a chemical with a biological system is determined by the functional groups, stereochemistry and other molecular features [114]. A change in the structure can lead to changes in toxicity, depending on how this change affects the toxicophore – the structural portion associated with the toxicity of the chemical [115]. The descriptors used for toxicity predictions vary between models, some models consider only 2D descriptors, while others use a mix of 2D and 3D descriptors. In recent years some models have also attempted to predict toxicity through graph neural networks which take the position of the different functional groups into consideration [116, 117]. Toxicity is also related to the absorption, distribution, metabolism and excretion ability of the body, which depends on the chemical structure of the substance. Increased hydrophobicity leads to increased absorption and increased potential toxic effects (due to potential bioaccumulation). A concern with PMT/vPvM substances is chronic exposure via water consumption, which can lead to elevated concentrations in humans and diverse biota [12, 118].

Considering the chemical features that make these substances P, M and T, and the provided criteria for classification, PMT/vPvM substances can be identified and grouped accordingly. Readacross can be applied to identify PMT substances on the basis that they have similar (sub)structures, similar properties and available toxicological data. This can be done through a fragment-based approach, which relies on the identification of small similar fragments or functional groups with similar properties [119]. Substance grouping approaches take the fragment models one step further. Substances can have similar properties due to the similarity in structures - this includes functional groups, common precursors or reaction products, and a constant pattern in the changing of the potency of the properties across the group [31, 104]. The substance grouping approach is used in read-across techniques and alternative assessments [104]. This has been supported by the use of various models that predict physical-chemical property information such as the KOCWIN model [120]. This is a fragment-based model that relates substructures and mobility (K_{OC}) based on appropriate training data [120]. The mobility (K_{OC}) of new chemicals can be predicted using the established relationship (when they are within the applicability domain). Similarly, the ReadyBiodegradable model in OPERA [36] and CERAPP and CoMPARA models [116, 121] can be used to predict the biodegradability and toxicity, respectively, of many new PMT/vPvM substances.

Grouping based on retained moieties from transformation reactions

Both biotic and abiotic transformations generally result in transformation products (TPs) with significantly higher mobility (lower $\log K_{\rm OW}$) than their parent compounds or precursors, making them more mobile in the environment. This seems intuitive as one of the "goals" of metabolism and wastewater treatment processes is to increase the polarity of the compounds. For metabolic processes, this allows the compounds to be expelled from the body with the urine. However, some persistent substructures may be retained during the transformation resulting in similar toxic properties between the parent compound and TPs. For example, decabromodiphenyl ethane forms 6 metabolites that were found to be carcinogenic like the parent compound [122]. In other cases, even small structural changes can result in large differences in toxicity such as for bis(pentabromophenyl) ether, which has low thyroid binding affinity itself but can form the TP 2,3,5,6-tetrabromo-4-(2,3,4,5,6-pentabromophenoxy)phenol with a strong thyroid binding affinity [122]. Therefore, it is of interest not only to identify parent compounds that may be PMT/vPvM,

but also compounds that can form PMT/vPvM TPs, along with identifying the common persistent 472 and mobile moieties that were retained and not readily metabolized. Such an approach would be 473 consistent with the approach to substance grouping in REACH of identifying "common breakdown 474 products via physical and biological processes, which result in structurally similar chemicals" 475 (REACH Annex XI, Section 1.5).

471

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

One way to do this identification is by using known reactions, such as those contained within the "transformations" section in PubChem, which links parent compounds to TPs with basic reaction information [123]. Of all the substances on the five suspect lists mentioned above, perfluorooctanoic acid (PFOA) has the highest number of recorded parent compounds (22) in the PubChem transformations section (from version 0.1.6 of the dataset archived on Zenodo) [124]. This is also one of several PMT/vPvM TPs where parent compounds are included in the legislation as part of the substance group. Specifically, the Stockholm Convention restricts the use of PFOArelated compounds, which include compounds that form PFOA after degradation [78]. TFA is also a known TP of several different compounds containing a CF₃- moiety. Since there are over 5 million compounds containing a CF₃-moiety included in PubChem [125], many of which are also industrially relevant, there are myriad potential sources of TFA.

Triazoles are another example of moieties preserved from parent compounds to TPs. Out of the triazole compounds in PubChem, 62 have recorded TPs (12 Jan 2024) [124]. This corresponds to 233 unique reactions, out of which the triazole moiety is retained in 88% of the cases, showing its high stability. While usually not acutely toxic in low doses, triazoles may cause several severe chronic toxic effects such as endocrine disruption and neurotoxicity [126]. Triazole substances such as benzotriazoles are present in the environment at high levels [126]. Thus, the retention of this functional group and other PMT/vPvM moieties may be concerning from an exposure point of view, supporting the idea that grouping based on retained moieties or TPs is relevant to obtain comprehensive and inclusive groups for PMT/vPvM substances.

Another example of how substances can be regulated based on their precursors are the aromatic amines registered under REACH some of which can also be found on PMT/vPvM suspect lists, such as 4,4'-methylenedianiline, 4,4'-oxydianiline, 4-chloroaniline, 3,3'-dichlorobenzidine, 4,4'methylenebis(2-chloroaniline), 4,4'-methylenedianiline, and 2-methoxy-5-methylaniline [20, 31, 50]. However, there are several other aromatic amines that have multiple known precursors listed in the PubChem transformations library (e.g. 4-aminophenol and aniline) that are also listed as

PMT by the UBAPMT suspect list. As such, regulating aromatic amines and precursor substances as a group based on PMT/vPvM substances warrants further investigation and consideration.

Cheminformatics challenges in grouping PMT/vPvM substances

The strategies mentioned above regarding grouping can be matched to substance regulation in various ways. Firstly, substances falling within a PMT/vPvM group with little known data could be flagged for follow-up to see if persistence and mobility measurements have been conducted in literature, and if read-across methods (based on patterns with other substances) could be used to fill data gaps. This could help prioritise filling data gaps and would help develop read-across approaches for further substances within the group. Secondly, if any other substance with a similar structure exists in the group and is hazardous (source substance), this could be seen as a reason to investigate for similar toxicological hazards for this substance (target substance). If several members within a group are shown to be PMT/vPvM substances, then a precautionary approach would be to assume all group members with no assessments are similarly hazardous, until there is sufficient scientific data to show otherwise. An overview of an assessment procedure to identify PMT/vPvM substances has also been presented by Neumann and Schliebner (2019), suggesting to first assess compounds for persistence followed by mobility and toxicity for both precursors and transformation products [47].

Estimating the number of individuals and groups of chemicals that fit the PMT/vPvM classification is a challenging task. Most substances lack readily available persistence, mobility and toxicity data. Due to the limited data availability, the prediction models of these properties are also limited in their accuracy and applicability domains. However, some data and models are available as a starting point, especially for $\log K_{OW}$. For example, both XlogP and the newer XlogP3 [127, 128] and the KOWWIN [129] module in EPI Suite [130] use a multivariate regression approach to K_{OW} prediction. While K_{OC} data is more limited than K_{OW} data, there are also several models which attempt to predict K_{OC} , including KOCWIN from EPI Suite [37] and the K_{OC} module of OPERA [36], which both utilise the training data from the PHYSPROP database [36, 37]. For persistence, OPERA also contains three modules for predicting biodegradability (BiodegHL,

ReadyBiodegradable and Km) as well as one module predicting rate constants for gas-phase reactions with hydroxyl-radicals, though with more limited applicability domains compared to the K_{OC} module [36]. The prediction of toxicity is even more challenging due to the limited availability of training data. However, models already exist for several toxicological endpoint predictions such as endocrine disruption, mutagenicity and developmental toxicity for many possible structures [131, 132]. For example, the CERAPP and CoMPARA models in OPERA can be used to predict estrogen and androgen receptor interactions [116, 121].

An example of how a combination of experimental and predicted property data has been used to classify compounds as PMT/vPvM is the study by Arp and Hale (2022), which focused on the compounds in the REACH inventory [24]. Their study concluded that about 1.9% of substances registered in REACH and 24% of REACH-registered substances that were detected in drinking water sources would be classified as PMT. However, over 40% of the inventory could not be assessed due to data gaps. This illustrates the importance of acquiring experimental data for a wider range of compounds.

To further illustrate how predicted properties can be used for assessing PMT compounds, the PubChemLite for Exposomics [133] dataset was used to give a preliminary estimate of the number of potential environmental contaminants that are predicted to meet the CLP definition of mobility. PubChemLite is a subset of ~350,000 compounds from PubChem with environmentally relevant annotation content [133]. In total, $K_{\rm OC}$ values were predicted for 346,133 compounds (96.0% of PubChemLite) using the molecular connectivity index (MCI) model from KOCWIN [37, 120]. The results of the log $K_{\rm OC}$ distribution are shown in Figure 5. Based on these predicted values log $K_{\rm OC}$, 147,930 compounds (41%) would be considered mobile and 78,107 compounds (21%) would be considered very mobile. The MCI model assigns values of 1 and 0 for chemicals when $K_{\rm OC}$ predictions have been overcorrected. The large peak at log $K_{\rm OC}$ 10 is an upper bound for these predictions within the MCI model output. Note that the results of this analysis will differ if the $K_{\rm OW}$ -based model in KOWWIN is used.




Figure 5: Predicted log K_{OC} via KOCWIN values for PubChemLite (346,133 compounds total) [38, 133]. Some K_{OC} predictions fall outside the applicability domain of the model. The "count" is the number of compounds inside each bin. Bin widths are 0.2.

To investigate this concept in more detail on slightly smaller subsets, four compound classes: triazines, aromatic amines, PFAS and triazoles were selected based on the five suspect lists mentioned previously (UBAPMT, EAWAGPMT, UFZHSFPMT, PMTPFAS and ZEROPMBOX1) [20, 49–52]. The substructure search function of PubChem was then used to estimate the number of compounds that belong to these compound classes. An overview of these search queries can be found in Table 1. As can be seen in the first three rows, relatively unspecific substructures can generate very large query results. For the 1,2,3-triazole substructure over 1 million substances were found, which corresponded to a search of only 87% of the database as the search results are capped at 1 million substances. Similarly, the aromatic amines (with aniline as the searched substructure) resulted in over 1 million substances with only 2% of the database searched. The 1,3,5-triazine query resulted in 718,119 substances. The compounds classified within the OECD definition of PFAS in the PubChem PFAS tree were used to source PFAS, which contained 6,604,017 compounds at the time the queries were performed (Dec. 17, 2023) [125]. It is unlikely that all compounds containing these substructures would meet the PMT/vPvM criteria outlined in the CLP. For example, only 250,873 of the 718,199 triazines meet the CLP criteria for mobility

based on the predicted log K_{OW} from XlogP3. As such the substructures used when discussing PMT/vPvM compound grouping for regulation purposes should be more specific.

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

More specific substructure queries were used to interrogate these results further according to the PMT/vPvM criteria, shown in the remaining rows of Table 1 and Figure 6, using the structures of melamine, benzotriazole and benzidine as well as the "larger PFAS parts" definition (contains -CF₂CF₂-) from the PubChem PFAS tree [125]. The search was then progressively restricted to exclude the searched substructures from being part of larger ring systems and substances with molecular weights greater than 300 g/mol. The results of this search were also used to perform persistence and toxicity prediction as discussed above. The top two rows of Figure 6 show these substructure query results. Like their less specific triazine counterpart, they seem too broad to capture only PMT/vPvM substances. However, when restricting the search to only compounds below 300 g/mol, between 87 and 99% of the compounds would meet the CLP definition of mobility. The most restricted PubChem search results were also used for biodegradability and LC₅₀ predictions using the ReadyBiodegradable model from OPERA and MS2Tox (compounds which fell outside the applicability domain of the ReadyBiodegradable model were excluded in the biodegradability results; this information is not given by MS2Tox). As can be seen in Figure 6, almost all compounds inside the applicability domain were classified as non-biodegradable. This indicates that in addition to their mobility, most compounds within these classes are potentially persistent. Based on the EU CLP definition of toxicity, an EC₁₀ less than 0.01 mg/L is toxic for marine or freshwater organisms. Some of the predicted LC₅₀ toxicity values (the lethal concentration at 50% of the population) shown in Figure 6 go below 0.1 mg/L. As toxicity is usually observed before lethality, this indicates that some of the compounds may be toxic below 0.01 mg/L. In addition, it is important to note that this is not the only toxicity criterion used in the CLP, for example, triazole compounds have been shown to have endocrine-disrupting properties by e.g., Wang et al. (2022) [134] and thus it is likely that at least a fraction of these compounds would be classified as PMT.

Table 1: Overview of the PubChem substructure searches (queries performed on 19/12/2023). Default settings for all queries were (1) single or double bonds match aromatic bonds (2) chain bonds in the query may match rings in hits and (3) remove any explicit hydrogens before searching. Extra settings "Ring" (rings may not be embedded in a larger system) and the filter "MW<300" (Molecular weight < 300 g/mol) are indicated in the respective column. Query URLs are embedded into the substructure column. The first three rows in italics are generic queries, the latter rows are the queries used to generate Figure 6 and are coloured accordingly. The OECD PFAS results were obtained via the PubChem classification browser. DB = database, values in brackets indicate the % of the database searched (queries are capped at 1 M results).

Substructure	Substructure SMARTS	Extra Settings	Number of Structures
1,3,5-Triazine	<u>C1=NC=NC=N1</u>	None	718,134
Aromatic amine	<u>C1=CC=C(C=C1)N</u>	None	>1 M (2 % of DB)
1,2,3-Triazole	<u>C1=NNN=C1</u>	None	>1 M (87 % of DB)
Melamine	C1(=NC(=NC(=N1)N)N)N	None	96,660
Melamine	C1(=NC(=NC(=N1)N)N)N	Ring	47,519
Melamine	C1(=NC(=NC(=N1)N)N)N	Ring, MW<300	10,528
Benzotriazole	C1=CC2=NNN=C2C=C1	None	165,361
Benzotriazole	C1=CC2=NNN=C2C=C1	Ring	158,766
Benzotriazole	<u>C1=CC2=NNN=C2C=C1</u>	Ring, MW<300	30,945
Benzidine	C1=CC(=CC=C1C2=CC=C(C=C2)N)N	None	208,934
Benzidine	<u>C1=CC(=CC=C1C2=CC=C(C=C2)N)N</u>	Ring	48,123
Benzidine	C1=CC(=CC=C1C2=CC=C(C=C2)N)N	Ring, MW<300	1,618
OECD PFAS	PFAS Tree - OECD PFAS definition	None	6,604,017
OECD PFAS OECD PFAS	PFAS Tree - OECD PFAS definition PFAS Tree - OECD PFAS definition	Larger PFAS Parts Larger PFAS Parts, MW<300	222,174 29,521




Figure 6: Distribution of PubChem XlogP3 values for four searches with varying restrictions (top three rows), plus predicted biodegradability from the OPERA ReadyBiodegradable model and LC₅₀ values (in mg/L) predicted from MS2Tox for the most restricted search queries (last two rows). Search settings and total numbers are given in Table 1. It should be noted that MS2Tox is used for illustration purposes only, and does not reflect the CLP toxicity criteria in Figure 1.

Perspectives for grouping strategies

Challenges for assessing and managing PMT/vPvM substances

Based on the criteria for persistence, mobility and toxicity, many chemicals (as shown from the prediction results above) may be PMT/vPvM substances, warranting further assessments and management (see Table 1). One of the major challenges of dealing with PMT/vPvM substances is the lack of high-quality data. As shown above, there is a growing need for computational modelling to complement experimental approaches as the number of substances to be tested increases, yet efficient computational methods rely on high quality and sufficient experimental data availability, which is a major limiting factor. Structurally diverse substances with multiple functional groups and stereochemistry can lead to unexpected behaviour, posing a critical challenge [135]. However, developing high-throughput screening (HTS) methods for toxicity testing may be difficult for these diverse substances and toxicological endpoints, which produce large amounts of data, and require expert knowledge to interpret and manage [136].

Identification, monitoring and removal are other challenges to managing PMT/vPvM substances. Targeted identification and monitoring approaches will not be sufficient to detect all PMT/vPvM substances, hence it is crucial to ensure wider availability of analytical methods and reference standards to properly identify, quantify and assess potential PMT/vPvM substances. Targeted analysis could be complemented with non-targeted analytical approaches. Non-target analysis can be a first step in identifying unknown and emerging PMT/vPvM substances and their TPs in environmental and biological matrices. It can also be used for the monitoring of known PMT/vPvM substances, yet regulatory acceptance of non-targeted monitoring is still lacking [137]. Moreover, the lack of data on persistence, mobility, toxicity, TPs, complex mixtures and appropriate suspect lists for broader suspect screening poses identification and monitoring challenges [22].

Prioritization strategies for testing and assessment

Considering many potential PMT/vPvM substances, and the possible challenges outlined, it is necessary to develop other strategies that could be used in combination with the proposed

grouping strategies above to prioritize chemicals for assessing their PMT/vPvM properties. Exposure and emission information could serve in the prioritization of chemicals for testing. Utilising exposure information such as mode of exposure (chemicals in food or drinking water) [138, 139] or Occupational Exposure Banding strategy for categorization of airborne substances lacking defined limits [140], could help identify chemicals with high exposure potential. Additionally, the hazard level of the toxicity can be used as a criterion, where specific persistent and mobile substances with high toxicity can be targeted for prioritization.

The use of *in vitro*, *in vivo*, and *in silico* approaches can aid the grouping and prioritization of chemicals for testing or assessments by regulatory bodies. This can be done by generating grouping hypotheses and justification for inclusion or exclusion criteria for substances in groups [140]. In particular, novel approaches may generate huge amounts of toxicological and high-throughput "omics" data including metabolomics, transcriptomics, and exposomics to support the validation and establishment of grouping hypotheses needed by regulatory authorities [141]. An example of a grouping hypothesis is the mode of action (MOA) hypothesis, which states that all chemicals that share a common mode of action are candidates for grouping [142]. Additionally, concepts such as adverse outcome pathways and toxicity pathways can be translated into prioritization hypotheses that can target specific substances, hence advancing prioritization for hazard assessment [143, 144]. These can facilitate regulator efforts to restrict hazardous substances.

Measurable biological and physiological effects of chemical exposure, known as effect biomarkers, can be used to group PMT/vPvM substances according to their toxicological profiles. If two substances induce similar biomarker responses indicative of for instance genotoxicity, or endocrine disruption, they can be grouped. This approach can also help to identify common pathways of toxicity, adverse outcome pathways (AOPs), or mechanisms of toxicity. By grouping substances with similar toxic profiles as PMT/vPvM substances, effect biomarkers can aid in characterizing cumulative risks posed by mixtures and complex chemical substances. This proactive approach focuses on biological effects rather than relying solely on the structural and chemical properties of PMT/vPvM substances [145, 146]

It is crucial to recognize that the effectiveness of strategies used for prioritizing substances largely depends on the availability of relevant data, including hazard and exposure data. To improve the availability and quality of such data, it is necessary to foster collective efforts such as high-quality data generation, community-level data collection, Open Science, FAIR data, and enhanced data

sharing policies. When potential vPvM substances are identified by read-across and experimental evidence as substances that meet the PMT/vPvM criteria, then a hazard classification would be required. The data collected through these efforts can then be used to enhance the prioritization, identification, and regulation of PMT/vPvM substances. To manage PMT/vPvM substances effectively, manufacturers can be encouraged to submit a plan to remove these substances from wastewater. This approach can help improve the overall management of PMT/vPvM substances. As a precaution, the principle of "as low as reasonably achievable" (ALARA) can be applied to minimize exposure levels and reduce the risk of harm to PMT/vPvM substances [147].

Conclusion

To achieve the EU's zero pollution ambition of a non-toxic environment by 2050, regulating the production, use and disposal of PMT/vPvM substances is necessary. As shown in Table 1, scaling PMT/vPvM criteria to big substance collections reveals that there are potentially thousands of PMT/vPvM substances that could cause harm to human health and the environment, especially concerning water quality and drinking water treatment.

The Montreal Protocol and the Stockholm Convention have demonstrated that grouping substances can be an effective strategy to expedite the elimination of the production, use, and emission of toxic substances such as ozone-depleting substances (ODSs) and persistent organic pollutants (POPs). This approach can also accelerate the identification and regulatory processes for substances that lack hazard information. Grouping can prevent the introduction of new hazardous substances into the global market. However, it is important to ensure that grouping is done in a way that is feasible and promotes the use of safer and more sustainable alternatives. Otherwise, it could result in the production of regrettable substitutes, as was the case with ODSs and POPs, which led to the creation of some PMT/vPvM substances.

Read-across based on structural or substructural similarity is one of the strategies that could be used to group PMT/vPvM substances, which relies on the idea that substances with similar structures have similar properties. Commonly retained moieties from transformation reactions could also be a grouping strategy for PMT/vPvM substances. Substances that are structurally

similar to PMT/vPvM according to read-across, or form persistent, mobile TPs could be flagged for subsequent assessment and/or regulatory actions.

Cheminformatics may be used for substance grouping based on predictive models for properties such as biodegradability, mobility, and toxicity. PubChemLite predictions suggest that 41% of potentially environmentally relevant compounds would be considered mobile (147,930 compounds) and 21% would be considered very mobile (78,107 compounds). Certain compound classes, such as triazines, aromatic amines, triazoles and PFAS, are likely to be persistent, non-biodegradable, and toxic. As shown in Table 1, the numbers are high and restricting these compounds as a group would be challenging; however, prioritizing members of these large groups for property testing is warranted as they contain a substructure associated with a PMT/vPvM substance group. Additional strategies are needed to prioritize some substances for regulation, such as mapping the uses of these chemicals and exposure which requires more data availability following FAIR principles.

Some strategies proposed for the prioritization of substances for testing or assessment of PMT/vPvM include (i) better understanding and the use of exposure/ emissions information such as Occupational Exposure Banding and environmental exposure to the PMT/vPvM substances, (ii) the use of *in vitro*, *in vivo* and *in silico* techniques to generate relevant toxicological data that will support prioritization, identification and regulation of PMT/vPvM substances. These strategies in combination with substance grouping could result in substituting PMT/vPvM substances with safer alternatives.

List of abbreviations

Abbreviations	Full Meaning
CAS	Chemical Abstracts Service
CFCs	Chlorofluorocarbons
CID	PubChem Compound Identifier
CLP	Chemicals, Labelling and Packaging
DDT	Dichlorodiphenyltrichloroethane
D_{OW}	pH-dependent octanol-water partition coefficient
EC	European Commission
EC _x	Effect concentration at X percentage
ECHA	European Chemical Agency
EDC	Endocrine-disrupting chemical
EU	European Union
HBFCs	Hydrobromofluorocarbons

Abbreviations	Full Meaning	
HCB	Hexachlorobenzene	
HCFCs	Hydrochlorofluorocarbons	
HFCs	Hydrofluorocarbons	
HFOs	Hydrofluoroolefins	
HTS	High-Throughput Screening	
IUPAC	International Union of Pure and Applied Chemistry	
KEMI	Swedish Chemical Agency	
Koc	Organic-carbon-water partition coefficient	
Kow	Octanol-water partition coefficient at pH = 7	
log P	Logarithm of Kow	
M	Mobility	
MCI	Molecular Connectivity Index	
MOA	Mode Of Action	
NOEC	No Observable Effect Concentration	
NORMAN-SLE	NORMAN Suspect List Exchange	
ODS	Ozone-Depleting Substances	
Р	Persistence	
PBT	Persistent, Bioaccumulative and Toxic	
PCBs	Polychlorinated biphenyls	
PFAS	Per- and poly-fluoroalkyl substances	
PFBS	Perfluorobutane sulfonic acid	
PFECAs	Per- and polyfluoroalkylether carboxylic acids	
PFOA	Perfluorooctanoic acid	
PMT	Persistent, Mobile and Toxic	
PMT/vPvM	Persistent, Mobile and Toxic or very Persistent and very Mobile	
POPs	Persistent Organic Pollutants	
QSAR	Quantitative Structure-Activity Relationship	
RAAF	Read-Across Assessment Framework	
REACH	Registration, Evaluation, Authorisation and Restriction of Chemicals	
SCHEER	Scientific Committee on Health, Environmental and Emerging Risks	
SVHCs	Substances of Very High Concern	
T	Toxicity	
TFA	Trifluoroacetic acid	
TPs	Transformation products	
UBA	Umweltbundesamt (German Environmental Agency)	
UVCBs	Substances of Unknown or Variable composition, Complex reaction	
	products or Biological origin	
νM	very Mobile	
vP	very Persistent	
vPvB	very Persistent and very Bioaccumulative	
vPvM	very Persistent and very Mobile	

728	Declarations
729	Ethics approval and consent to participate
730	Not applicable.
731	Consent for publication
732	Not applicable.
733	Availability of data and material
734 735 736 737 738 739 740 741 742 743	The datasets analysed during the current study are available in PubChem, the NORMAN Suspect List Exchange (NORMAN-SLE - https://www.norman-network.com/nds/SLE/) and on Zenodo under the following URLs: S36 UBAPMT (DOI: 10.5281/zenodo.6482414), S82 EAWAGPMT (DOI: 10.5281/zenodo.6482414), S82 EAWAGPMT (DOI: 10.5281/zenodo.6482414), S82 EAWAGPMT (DOI: 10.5281/zenodo.5500132), S84 UFZHSFPMT (DOI: 10.5281/zenodo.5535288), S90 ZeroPMBox1 (DOI: 10.5281/zenodo.5500132), S84 UFZHSFPMT (DOI: 10.5281/zenodo.5535288), S90 ZeroPMBox1 (DOI: 10.5281/zenodo.5535288), PubChem Transformations Dataset (Version 0.1.6, Jul. 5th, 2023, DOI: 10.5281/zenodo.8117741) and the PubChem PFAS Tree (

- 751 Research Fund (FNR) for project A18/BM/12341006. SB acknowledges support from Natural
- 752 Sciences and Engineering Research Council of Canada via a Postdoctoral Fellowship.

753 Author contributions:

- 754 PC: Investigation (lead), Visualization, Writing original draft (lead), Writing review and editing,
- 755 EHP: Investigation, Formal analysis (lead), Visualization, Writing original draft (lead), Writing
- review and editing, SB: Writing the original draft, Writing review and editing, ELS: Investigation,
- 757 Writing the original draft, Writing review and editing, Conceptualization, Supervision, Funding
- acquisition, ZW: Writing the original draft, Writing review and editing, Funding acquisition, RW:
- 759 Writing review and editing, SH: Writing original draft, Writing review and editing, Funding
- acquisition, Conceptualization, HPA: Writing original draft, Writing review and editing, Funding
- 761 acquisition, Conceptualization

762

763

Acknowledgements

- The authors would like to acknowledge discussions with other ZeroPM partners and PubChem
- during the preparation of this article. All authors acknowledge funding from the European Union's
- Horizon 2020 research and innovation programme under grant agreement No 101036756, project
- 767 ZeroPM: Zero pollution of persistent, mobile substances. ELS acknowledges funding support from
- the Luxembourg National Research Fund (FNR) for project A18/BM/12341006. SB acknowledges
- support from Natural Sciences and Engineering Research Council of Canada via a Postdoctoral
- 770 Fellowship.

771 References

- 772 1. Fuller R, Landrigan PJ, Balakrishnan K, et al (2022) Pollution and health: a progress update. The Lancet Planetary Health 6:e535–e547. https://doi.org/10.1016/S2542-5196(22)00090-0
- United Nations (2015) Resolution adopted by the General Assembly on 25 September
 2015. Transforming our world: the 2030 Agenda for Sustainable Development.
 https://undocs.org/Home/Mobile?FinalSymbol=A%2FRES%2F70%2F1&Language=E&DeviceType=Desktop&LangRequested=False
- 779 3. Wang Z, Walker GW, Muir DCG, Nagatani-Yoshida K (2020) Toward a Global Understanding of Chemical Pollution: A First Comprehensive Analysis of National and

- 781 Regional Chemical Inventories. Environ Sci Technol 54:2575–2584. 782 https://doi.org/10.1021/acs.est.9b06379
- 783 4. Kim S, Chen J, Cheng T, et al (2023) PubChem 2023 update. Nucleic Acids Research 51:D1373–D1380. https://doi.org/10.1093/nar/gkac956
- 785 5. PubChem (2023) PubChem-Explore Chemistry. https://pubchem.ncbi.nlm.nih.gov/.
 786 Accessed 4 Dec 2023
- 787 6. CAS CAS REGISTRY | CAS. https://www.cas.org/cas-data/cas-registry. Accessed 30 Jan 2024
- 789 7. Resnik DB, Portier CJ (2022) Environment, Ethics, and Human Health The Hastings Center. https://www.thehastingscenter.org/briefingbook/environmental-health/. Accessed 6 Nov 2023
- 792 8. Persson L, Carney Almroth BM, Collins CD, et al (2022) Outside the Safe Operating Space 793 of the Planetary Boundary for Novel Entities. Environ Sci Technol 56:1510–1521. 794 https://doi.org/10.1021/acs.est.1c04158
- Furostat (2023) Chemicals production and consumption statistics Statistics Explained.
 https://ec.europa.eu/eurostat/databrowser/view/sdg_12_10/default/bar?lang=en.
 Accessed 4 Oct 2023
- 798 10. SCHEER (Scientific Committee on Health, Environmental and Emerging Risks) (2018) 799 Statement on emerging health and environmental issues. 800 https://health.ec.europa.eu/system/files/2019-02/scheer s 002 0.pdf
- 801 11. European Commission (2022) Commission Delegated Regulation (EU) 2023/707 of 19
 802 December 2022 amending Regulation (EC) No 1272/2008 as regards hazard classes and
 803 criteria for the classification, labelling and packaging of substances and mixtures.
 804 https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex%3A32023R0707
- Hale SE, Arp HPH, Schliebner I, Neumann M (2020) Persistent, mobile and toxic (PMT) and very persistent and very mobile (vPvM) substances pose an equivalent level of concern to persistent, bioaccumulative and toxic (PBT) and very persistent and very bioaccumulative (vPvB) substances under REACH. Environ Sci Eur 32:155. https://doi.org/10.1186/s12302-020-00440-4
- 810 13. Matthies M, Solomon K, Vighi M, et al (2016) The origin and evolution of assessment criteria for persistent, bioaccumulative and toxic (PBT) chemicals and persistent organic pollutants (POPs). Environ Sci: Processes Impacts 18:1114–1128. https://doi.org/10.1039/C6EM00311G
- Shrestha P, Junker T, Fenner K, et al (2016) Simulation Studies to Explore Biodegradation in Water–Sediment Systems: From OECD 308 to OECD 309. Environ Sci Technol 50:6856–6864. https://doi.org/10.1021/acs.est.6b01095
- 15. Van Dijk J, Figuière R, Dekker SC, et al (2023) Managing PMT/vPvM substances in consumer products through the concepts of essential-use and functional substitution: a

- case-study for cosmetics. Environ Sci: Processes Impacts 25:1067–1081. https://doi.org/10.1039/D3EM00025G
- 821 16. Albergamo V, Escher BI, Schymanski EL, et al (2020) Evaluation of reverse osmosis 822 drinking water treatment of riverbank filtrate using bioanalytical tools and non-target 823 screening. Environ Sci: Water Res Technol 6:103–116. 824 https://doi.org/10.1039/C9EW00741E
- Camacho L, Kelly KP, Beland FA, Gamboa da Costa G (2011) Gene expression of
 biomarkers of nephrotoxicity in F344 rats co-exposed to melamine and cyanuric acid for
 seven days. Toxicol Lett 206:166–171. https://doi.org/10.1016/j.toxlet.2011.07.009
- 828 18. Habotta OA, Abdeen A, Roomi AB, et al (2023) Nootkatone Mitigated Melamine-Evoked 829 Hepatotoxicity by Featuring Oxidative Stress and Inflammation Interconnected 830 Vivo Mechanisms: In and In Silico Approaches. **Toxics** 11:784. 831 https://doi.org/10.3390/toxics11090784
- Hau AK, Kwan TH, Li PK (2009) Melamine Toxicity and the Kidney. Journal of the American Society of Nephrology 20:245–250. https://doi.org/10.1681/ASN.2008101065
- Arp HPH, Hale SE, Schliebner I, Neumann M (2022) S36 | UBAPMT | Prioritised PMT/vPvM substances in the REACH registration database _all versions. https://doi.org/10.5281/ZENODO.2653212
- Aumeier BM, Georgi A, Saeidi N, Sigmund G (2023) Is sorption technology fit for the removal of persistent and mobile organic contaminants from water? Science of The Total Environment 880:163343. https://doi.org/10.1016/j.scitotenv.2023.163343
- Rüdel H, Körner W, Letzel T, et al (2020) Persistent, mobile and toxic substances in the environment: a spotlight on current research and regulatory activities. Environ Sci Eur 32:5. https://doi.org/10.1186/s12302-019-0286-x
- 23. Lütjens LH, Pawlowski S, Silvani M, et al (2023) Melamine in the environment: a critical review of available information. Environmental Sciences Europe 35:2. https://doi.org/10.1186/s12302-022-00707-y
- 846 24. Arp HPH, Hale SE (2022) Assessing the Persistence and Mobility of Organic Substances 847 to Protect Freshwater Resources. ACS Environ Au 2:482–509. 848 https://doi.org/10.1021/acsenvironau.2c00024
- 849 25. Tickner JA, Schifano JN, Blake A, et al (2015) Advancing safer alternatives through 850 functional substitution. Environmental science & technology 49:742–749. 851 https://doi.org/10.1021/es503328m
- 26. Žalmanová T, Hošková K, Nevoral J, et al (2016) Bisphenol S instead of bisphenol A: a story of reproductive disruption by regretable substitution - a review. Czech J Anim Sci 61:433–449. https://doi.org/10.17221/81/2015-CJAS
- 855 27. IUPAC (2019) IUPAC Compendium of Chemical Terminology, "chemical substance." 3rd ed. International Union of Pure and Applied Chemistry; 2006, https://doi.org/10.1351/goldbook.C01039

- 858 28. PubChem (2024) Explore PubChem Documentation. 859 https://pubchem.ncbi.nlm.nih.gov/docs/. Accessed 23 Jan 2024
- 860 29. IUPAC (2019) IUPAC Compendium of Chemical Terminology, "mixture." 3rd ed. 861 International Union of Pure and Applied Chemistry; 2006, https://doi.org/10.1351/goldbook.M03949
- 30. Dimitrov SD, Georgieva DG, Pavlov TS, et al (2015) UVCB substances: Methodology for structural description and application to fate and hazard assessment. Environmental Toxicology and Chemistry 34:2450–2462. https://doi.org/10.1002/etc.3100
- 866 European Commission (2006) EUR-Lex - 02006R1907-20140410 - EN - EUR-Lex 31. Consolidated text: Regulation (EC) No 1907/2006 of the European Parliament and of the 867 868 Council of 18 December 2006 concerning the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH), establishing a European Chemicals Agency, amending 869 870 Directive 1999/45/EC and repealing Council Regulation (EEC) No 793/93 and Commission 871 Regulation (EC) No 1488/94 as well as Council Directive 76/769/EEC and Commission 872 91/155/EEC, 93/67/EEC, 93/105/EC and 2000/21/EC. 873 lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A02006R1907-20140410, Accessed 874 2023-12-20
- 32. Åberg A, MacLeod M, Wiberg K (2008) Physical-Chemical Property Data for Dibenzo-p-dioxin (DD), Dibenzofuran (DF), and Chlorinated DD/Fs: A Critical Review and Recommended Values. Journal of Physical and Chemical Reference Data 37:1997–2008. https://doi.org/10.1063/1.3005673
- 879 European Commission (2017) Commission Regulation (EU) 2017/644 of 5 April 2017 33. laying down methods of sampling and analysis for the control of levels of dioxins, dioxin-880 like PCBs and non-dioxin-like PCBs in certain foodstuffs and repealing Regulation (EU) No. 881 882 589/2014. Official Journal of the European Union L 92/9, https://eur-883 lex.europa.eu/eli/reg/2017/644/oj
- Mukherjee A, Debnath B, Ghosh SK (2016) A Review on Technologies of Removal of Dioxins and Furans from Incinerator Flue Gas. Procedia Environmental Sciences 35:528–540. https://doi.org/10.1016/j.proenv.2016.07.037
- 35. Garmash O, Hermanson MH, Isaksson E, et al (2013) Deposition History of Polychlorinated
 Biphenyls to the Lomonosovfonna Glacier, Svalbard: A 209 Congener Analysis. Environ
 Sci Technol 47:12064–12072. https://doi.org/10.1021/es402430t
- 890 36. Mansouri K, Grulke CM, Judson RS, Williams AJ (2018) OPERA models for predicting physicochemical properties and environmental fate endpoints. J Cheminform 10:10. https://doi.org/10.1186/s13321-018-0263-1
- 893 37. Meylan W, Howard PH, Boethling RS (1992) Molecular topology/fragment contribution 894 method for predicting soil sorption coefficients. Environ Sci Technol 26:1560–1567. 895 https://doi.org/10.1021/es00032a011
- 896 38. Bolton E, Schymanski E, Kondic T, et al (2023) PubChemLite for Exposomics (V. 1.27.0). 897 https://doi.org/10.5281/zenodo.10126889

- 898 39. Peets P, Wang W-C, MacLeod M, et al (2022) MS2Tox Machine Learning Tool for 899 Predicting the Ecotoxicity of Unidentified Chemicals in Water by Nontarget LC-HRMS. 900 Environ Sci Technol 56:15508–15517. https://doi.org/10.1021/acs.est.2c02536
- 901 40. R Core Team (2023) A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. https://cran.r-project.org/bin/windows/base/old/4.3.1/
- 903 41. Wickham H, Averick M, Bryan J, et al (2019) Welcome to the Tidyverse. JOSS 4:1686. 904 https://doi.org/10.21105/joss.01686
- 905 42. Hadley Wickham (2023) tidyverse 2.0.0. https://www.tidyverse.org/blog/2023/03/tidyverse-906 2-0-0/
- 907 43. Wickham H, Chang W, Wickham MH (2016) Package 'ggplot2.' Create elegant data 908 visualisations using the grammar of graphics Version 2:1–189. 909 https://ggplot2.tidyverse.org/reference/ggplot2-package.html
- 910 44. Thomas Lin Pedersen (2024) ggplot2 3.5.0. 911 https://github.com/tidyverse/ggplot2/releases/tag/v3.5.0
- 912 45. Emma L. Schymanski (2020) RChemMass: Various Cheminformatic, Curation and Mass
 913 Spectrometry Functions. https://github.com/schymane/RChemMass
- 914 46. Arp HPH, Hale SE, Schliebner I, Neumann M (2023) Prioritised PMT/vPvM substances in the REACH registration database | Umweltbundesamt. https://www.umweltbundesamt.de/publikationen/prioritised-pmtvpvm-substances-in-the-reach. Accessed 9 Oct 2023
- 918 47. Neumann M, Schliebner I (2019) Protecting the sources of our drinking water: The criteria for identifying persistent, mobile and toxic (PMT) substances and very persistent and very mobile (vPvM) substances under EU Regulation REACH (EC) No 1907/2006. https://www.umweltbundesamt.de/sites/default/files/medien/1410/publikationen/2019-11-29_texte_127-2019_protecting-sources-drinking-water-pmt.pdf
- 48. Holmberg R, Wedebye EB, Nikolov NG, Tyle H (2021) How many potential vPvM/PMT substances have been registered under REACH?-vPvM/PMT-screening by using the Danish (Q) SAR database.
 926 https://backend.orbit.dtu.dk/ws/portalfiles/portal/240040384/111514_909384_DTU_Rapp ort vPvM PMT CB 7kor links.pdf
- 928 49. Kiefer K, Du L, Singer H, Hollender J (2021) S82 | EAWAGPMT | PMT Suspect List from Eawag. https://doi.org/10.5281/ZENODO.5500131
- 930 50. Neuwald I, Muschket M, Zahn D, et al (2021) S84 | UFZHSFPMT | PMT Suspect List from UFZ and HSF. https://doi.org/10.5281/ZENODO.5535287
- 932 51. Schymanski E, Wang Z, Wolf R, Arp HP (2022) S90 | ZEROPMBOX1 | ZeroPM Box 1 Substances. https://doi.org/10.5281/ZENODO.5854251
- 934 52. Schymanski E (2023) S111 | PMTPFAS | Fluorine-containing Compounds in PMT Suspect Lists. https://doi.org/10.5281/ZENODO.8417075

- 936 53. Mohammed Taha H, Aalizadeh R, Alygizakis N, et al (2022) The NORMAN Suspect List Exchange (NORMAN-SLE): facilitating European and worldwide collaboration on suspect screening in high resolution mass spectrometry. Environ Sci Eur 34:104. https://doi.org/10.1186/s12302-022-00680-6
- 940 54. NORMAN Network (2024) NORMAN Suspect List Exchange. https://www.norman-network.com/nds/SLE/. Accessed 30 Jan 2024
- 942 55. Arp HPH, Hale SE (2020) S36 | UBAPMT | Potential Persistent, Mobile and Toxic (PMT) substances_V.0.2. https://doi.org/10.5281/ZENODO.3637611
- 944 56. UBA (2019) REACH: Improvement of guidance and methods for the identification and 945 assessment of PMT/vPvM substances | Umweltbundesamt. 946 https://www.umweltbundesamt.de/en/publikationen/reach-improvement-of-guidance-947 methods-for-the. Accessed 4 Oct 2023
- 948 57. Arp HPH, Hale SE (2020) S36 | UBAPMT | Potential Persistent, Mobile and Toxic (PMT) substances V.0.3. https://doi.org/10.5281/ZENODO.3637611
- 950 58. UBA (2023) Prioritised PMT/vPvM substances in the REACH registration database. 951 https://www.umweltbundesamt.de/sites/default/files/medien/11850/publikationen/21_2023 952 __texte_pmt_vpvm_substances_reach_v23032023.pdf
- 59. Kiefer K, Du L, Singer H, Hollender J (2021) Identification of LC-HRMS nontarget signals
 in groundwater after source related prioritization. Water Research 196:116994.
 https://doi.org/10.1016/j.watres.2021.116994
- 956 60. Fischer S (2023) S17 | KEMIMARKET | KEMI Market List. 957 https://doi.org/10.5281/ZENODO.2628786
- 958 61. Neuwald I, Muschket M, Zahn D, et al (2021) Filling the knowledge gap: A suspect screening study for 1310 potentially persistent and mobile chemicals with SFC- and HILIC-960 HRMS in two German river systems. Water Research 204:117645. 961 https://doi.org/10.1016/j.watres.2021.117645
- 962 62. Neuwald I, Muschket M, Zahn D, et al (2021) A suspect screening list of 1310 persistent and mobile (PM) candidates. https://doi.org/10.5281/zenodo.5503380
- 964 63. Blum A, Behl M, Birnbaum LS, et al (2019) Organophosphate Ester Flame Retardants: Are
 965 They a Regrettable Substitution for Polybrominated Diphenyl Ethers? Environ Sci Technol
 Lett 6:638–649. https://doi.org/10.1021/acs.estlett.9b00582
- 967 64. UNEP (2023) About Montreal Protocol. https://www.unep.org/ozonaction/who-we-968 are/about-montreal-protocol. Accessed 7 Oct 2023
- 969 65. United Nations (1989) Montreal Protocol on Substances that Deplete the Ozone Layer (with 970 Concluded Montreal September 1987. annex). at on 16 971 https://treaties.un.org/doc/publication/unts/volume%201522/volume-1522-i-26369english.pdf Vol. 1522, 1-26369: 972

- 973 66. Stockholm Convention (2023) Overview. 974 https://www.pops.int/TheConvention/Overview/tabid/3351/Default.aspx. Accessed 7 Oct 975 2023
- 976 67. ECHA (2020) Grouping speeds up regulatory action Integrated Regulatory Strategy Annual Report.
- 978 https://echa.europa.eu/documents/10162/5641810/irs_annual_report_2019_en.pdf
- 979 68. European Commission (2022) Consolidated text: Regulation (EC) No 1107/2009 of the
 980 European Parliament and of the Council of 21 October 2009 concerning the placing of plant
 981 protection products on the market and repealing Council Directives 79/117/EEC and
 982 91/414/EEC. https://eur-lex.europa.eu/legal983 content/EN/TXT/?uri=CELEX%3A02009R1107-20221121
- 984 69. European Commission (2022) Consolidated text: Regulation (EU) No 528/2012 of the European Parliament and of the Council of 22 May 2012 concerning the making available on the market and use of biocidal products (Text with EEA relevance)Text with EEA relevance. https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A02012R0528-20220415
- 989 70. European Commission (2021) Commission Regulation (EU) 2021/979 of 17 June 2021 990 amending Annexes VII to XI to Regulation (EC) No 1907/2006 of the European Parliament 991 and of the Council concerning the Registration, Evaluation, Authorisation and Restriction 992 of Chemicals (REACH). https://eur-lex.europa.eu/eli/reg/2021/979/oj. Accessed 16 Oct 993
- 994 71. Gonzalez M, Taddonio KN, Sherman NJ (2015) The Montreal Protocol: how today's successes offer a pathway to the future. J Environ Stud Sci 5:122–129. https://doi.org/10.1007/s13412-014-0208-6
- 997 72. Ravishankara AR, Daniel JS, Portmann RW (2009) Nitrous oxide (N2O): the dominant ozone-depleting substance emitted in the 21st century. Science 326:123–125. https://doi.org/DOI: 10.1126/science.1176985
- 1000 73. Umar SA, Tasduq SA (2022) Ozone Layer Depletion and Emerging Public Health Concerns
 1001 An Update on Epidemiological Perspective of the Ambivalent Effects of Ultraviolet
 1002 Radiation Exposure. Front Oncol 12:866733. https://doi.org/10.3389/fonc.2022.866733
- 74. EEA (2023) EU and global consumption of controlled ozone-depleting substances —
 1004 European Environment Agency. In: European Environmental Agency.
 1005 https://www.eea.europa.eu/data-and-maps/figures/consumption-of-controlled-ozone 1006 depleting-5/. Accessed 7 Oct 2023
- 1007 75. Nunez C (2023) Ozone layer facts and information.
 1008 https://www.nationalgeographic.com/environment/article/ozone-depletion. Accessed 9 Jan
 1009 2024
- 1010 76. ECOLEX (2004) Regulation (EC) No. 850/2004 of the European Parliament and of the Council on persistent organic pollutants and amending Directive 79/117/EEC. https://www.ecolex.org/details/legislation/regulation-ec-no-8502004-of-the-european-

1013 1014	parliament-and-of-the-council-on-persistent-organic-pollutants-and-amending-directive-79117eec-lex-faoc087038/. Accessed 9 Oct 2023
1015 77. 1016	Wheeler AF (2017) Study on the cumulative health and environmental benefits of chemica legislation: final report. https://data.europa.eu/doi/102779/070159
1017 78. 1018 1019	Stockholm Convention (2024) Listing of POPs in the Stockholm Convention https://www.pops.int/TheConvention/ThePOPs/AllPOPs/tabid/2509/Default.aspx. Accessed 7 Oct 2023
1020 79. 1021 1022	United Nations The 12 Initial POPs https://www.pops.int/TheConvention/ThePOPs/The12InitialPOPs/tabid/296/Default.aspx, Accessed 2024-01-30
1023 80. 1024 1025 1026	EEA (2023) National emissions reported to the Convention on Long-range Transboundary Air Pollution (LRTAP Convention). In: European Environmental Agency https://www.eea.europa.eu/en/datahub/datahubitem-view/5be6cebc-ed2b-4496-be59-93736fc4ad78?activeAccordion=1086857. Accessed 7 Oct 2023
1027 81. 1028	Aurich D (2024) Environmental Cheminformatics / chemicalstripes · GitLab https://gitlab.lcsb.uni.lu/eci/chemicalstripes, Accessed 31 Jan 2024
1029 82. 1030 1031	Maertens A, Golden E, Hartung T (2021) Avoiding Regrettable Substitutions: Green Toxicology for Sustainable Chemistry. ACS Sustainable Chem Eng 9:7749–7758 https://doi.org/10.1021/acssuschemeng.0c09435
1032 83. 1033 1034	Young AS, Allen JG, Kim U-J, et al (2018) Phthalate and Organophosphate Plasticizers in Nail Polish: Evaluation of Labels and Ingredients. Environ Sci Technol 52:12841–12850 https://doi.org/10.1021/acs.est.8b04495
1035 84. 1036	Tyl RW (2014) Abbreviated assessment of bisphenol A toxicology literature. Seminars in Fetal and Neonatal Medicine 19:195–202. https://doi.org/10.1016/j.siny.2013.11.010
1037 85. 1038	ECHA (2021) Assessment of regulatory needs https://echa.europa.eu/documents/10162/1bd5525c-432c-495d-9dab-d7806bf34312
1039 86. 1040 1041	McKenzie R, Bernhard G, Liley B, et al (2019) Success of Montreal Protocol Demonstrated by Comparing High-Quality UV Measurements with "World Avoided" Calculations from Two Chemistry-Climate Models. Sci Rep 9:12332. https://doi.org/10.1038/s41598-019-48625-2
1042 87. 1043 1044 1045 1046 1047 1048	UNEP-Ozone secretariat (2023) Consumption of controlled substances-Data in tables Ozone Secretariat. https://ozone.unep.org/countries/data-table?report_type=0&output_type=odp-CO2e-tonnes&party_grouping=individual&eu_member%5Bis_eu_member%5D=is_eu_member ._start=1986._end=2022&ignore_zero=1&baseline=1&group_by=group&op=GENERATE+REPORT&form_id=ozone_country_data_formreport_table_form. Accessed 15 Dec 2023

Tapscott RE, Moore TA, Mather JD, Vitali JA (1998) Halon replacement research—a historical review of technical progress and regulatory decision points. Halons Options

1049

1050

88.

- Technical Working Conference, Albuquerque, NM, https://www.nist.gov/system/files/documents/el/fire_research/R0000266.pdf
- Sukornick B (1989) Potentially acceptable substitutes for the chlorofluorocarbons: properties and performance features of HFC-134a, HCFC-123, and HCFC-141b. Int J Thermophys 10:553–561. https://doi.org/10.1007/BF00507978
- 1056 90. Benhadid-Dib S, Benzaoui A (2012) Refrigerants and their Environmental Impact Substitution of Hydro Chlorofluorocarbon HCFC and HFC Hydro Fluorocarbon. Search for an Adequate Refrigerant. Energy Procedia 18:807–816. https://doi.org/10.1016/j.egypro.2012.05.096
- 1060 91. Heath EA (2017) Amendment to the Montreal Protocol on Substances that Deplete the 1061 Ozone Layer (Kigali Amendment). Int leg mater 56:193–205. https://doi.org/10.1017/ilm.2016.2
- 1063 92. Luecken DJ, Waterland RL, Papasavva S, et al (2010) Ozone and TFA Impacts in North
 1064 America from Degradation of 2,3,3,3-Tetrafluoropropene (HFO-1234yf), A Potential
 1065 Greenhouse Gas Replacement. Environ Sci Technol 44:343–348.
 1066 https://doi.org/10.1021/es902481f
- 1067 93. Rivela CB, Tovar CM, Teruel MA, et al (2019) CFCs replacements: Reactivity and atmospheric lifetimes of a series of Hydrofluoroolefins towards OH radicals and Cl atoms. Chemical Physics Letters 714:190–196. https://doi.org/10.1016/j.cplett.2018.10.078
- 1070 94. Boutonnet JC, Bingham P, Calamari D, et al (1999) Environmental Risk Assessment of 1071 Trifluoroacetic Acid. Human and Ecological Risk Assessment: An International Journal 1072 5:59–124. https://doi.org/10.1080/10807039991289644
- 1073 95. Solomon KR, Carr JA, Du Preez LH, et al (2008) Effects of atrazine on fish, amphibians, and aquatic reptiles: a critical review. Critical reviews in toxicology 38:721–772. https://doi.org/10.1080/10408440802116496
- Dekant W, Dekant R (2023) Mammalian toxicity of trifluoroacetate and assessment of human health risks due to environmental exposures. Arch Toxicol 97:1069–1077. https://doi.org/10.1007/s00204-023-03454-y
- Morgan K, Bonanno F (2022) Sustainability is no longer a "nice to have" goal for the data center industry | S&P Global Market Intelligence.
 https://www.spglobal.com/marketintelligence/en/news-insights/research/sustainability-is-no-longer-a-nice-to-have-goal-for-the-data-center-industry. Accessed 16 Oct 2023
- 1083 98. Wang Z, DeWitt JC, Higgins CP, Cousins IT (2017) A Never-Ending Story of Per- and Polyfluoroalkyl Substances (PFASs)? Environ Sci Technol 51:2508–2518. https://doi.org/10.1021/acs.est.6b04806
- 1086 99. Knutsen H, Mæhlum T, Haarstad K, et al (2019) Leachate emissions of short- and long-1087 chain per- and polyfluoralkyl substances (PFASs) from various Norwegian landfills. Environ 1088 Sci: Processes Impacts 21:1970–1979. https://doi.org/10.1039/C9EM00170K

- 108. Helmer RW, Reeves DM, Cassidy DP (2022) Per- and Polyfluorinated Alkyl Substances (PFAS) cycling within Michigan: Contaminated sites, landfills and wastewater treatment plants. Water Research 210:117983. https://doi.org/10.1016/j.watres.2021.117983
- 101. ECHA (2013) Grouping of substances and read-across approach Part 1: Introductory note.
 1093 https://www.echa.europa.eu/documents/10162/17221/read_across_introductory_note_en.
 1094 pdf/1343b1b8-e5d1-4e72-b9b3-8a99e940ab29
- 1095 102. Lamon L, Asturiol D, Richarz A, et al (2018) Grouping of nanomaterials to read-across hazard endpoints: from data collection to assessment of the grouping hypothesis by application of chemoinformatic techniques. Part Fibre Toxicol 15:37. https://doi.org/10.1186/s12989-018-0273-1
- 1099 103. Schultz TW, Amcoff P, Berggren E, et al (2015) A strategy for structuring and reporting a read-across prediction of toxicity. Regulatory Toxicology and Pharmacology 72:586–601. https://doi.org/10.1016/j.yrtph.2015.05.016
- 1102 104. ECHA (2017) Read-across assessment framework (RAAF). ECHA-17-R-01-EN. https://doi.org/10.2823/619212
- 1104 105. OECD (2017) Guidance on Grouping of Chemicals, Second Edition. https://doi.org/10.1787/9789264274679-en
- 106. Patlewicz G, Helman G, Pradeep P, Shah I (2017) Navigating through the minefield of readacross tools: A review of in silico tools for grouping. Computational Toxicology 3:1–18. https://doi.org/10.1016/j.comtox.2017.05.003
- 1109 107. Tate T, Wambaugh J, Patlewicz G, Shah I (2021) Repeat-dose toxicity prediction with Generalized Read-Across (GenRA) using targeted transcriptomic data: A proof-of-concept case study. Computational Toxicology 19:100171. https://doi.org/10.1016/j.comtox.2021.100171
- 1113 108. ECHA (2023) Substance group and analogy concept ECHA.

 1114 https://echa.europa.eu/de/support/registration/how-to-avoid-unnecessary-testing-on1115 animals/grouping-of-substances-and-read-across. Accessed 20 Dec 2023
- 1116 109. Luo Y-R (2007) Comprehensive Handbook of Chemical Bond Energies. 1117 https://www.taylorfrancis.com/books/9781420007282 9780429128684:
- 1118 110. Hafner J, Fenner K, Scheidegger A (2023) Systematic Handling of Environmental Fate
 1119 Data for Model Development—Illustrated for the Case of Biodegradation Half-Life Data.
 1120 Environ Sci Technol Lett 10:859–864. https://doi.org/10.1021/acs.estlett.3c00526
- 111. Sigmund G, Arp HPH, Aumeier BM, et al (2022) Sorption and Mobility of Charged Organic
 1122 Compounds: How to Confront and Overcome Limitations in Their Assessment. Environ Sci
 1123 Technol 56:4702–4710. https://doi.org/10.1021/acs.est.2c00570
- 112. Zahn D, Neuwald IJ, Knepper TP (2020) Analysis of mobile chemicals in the aquatic 1125 environment—current capabilities, limitations and future perspectives. Anal Bioanal Chem 1126 412:4763–4784. https://doi.org/10.1007/s00216-020-02520-z

- 1127 113. Barratt MD (2000) Prediction of toxicity from chemical structure. Cell Biology and Toxicology 16:1–13. https://doi.org/10.1023/A:1007676602908
- 1129 114. Struyf J (2009) Relating Functional Groups to the Periodic Table. J Chem Educ 86:190. 1130 https://doi.org/10.1021/ed086p190
- 1131 115. Singh PK, Negi A, Gupta PK, et al (2016) Toxicophore exploration as a screening technology for drug design and discovery: techniques, scope and limitations. Arch Toxicol 90:1785–1802. https://doi.org/10.1007/s00204-015-1587-5
- 1134 116. Mansouri K, Abdelaziz A, Rybacka A, et al (2016) CERAPP: Collaborative Estrogen 1135 Receptor Activity Prediction Project. Environ Health Perspect 124:1023–1033. 1136 https://doi.org/10.1289/ehp.1510267
- 1137 117. Liu J, Lei X, Zhang Y, Pan Y (2023) The prediction of molecular toxicity based on BiGRU and GraphSAGE. Computers in Biology and Medicine 153:106524. https://doi.org/10.1016/j.compbiomed.2022.106524
- 1140 118. Freeling F, Scheurer M, Koschorreck J, et al (2022) Levels and Temporal Trends of Trifluoroacetate (TFA) in Archived Plants: Evidence for Increasing Emissions of Gaseous TFA Precursors over the Last Decades. Environ Sci Technol Lett 9:400–405. https://doi.org/10.1021/acs.estlett.2c00164
- 1144 119. Li Q (2020) Application of Fragment-Based Drug Discovery to Versatile Targets. Front Mol Biosci 7:180. https://doi.org/10.3389/fmolb.2020.00180
- 1146 120. US EPA (2012) Estimation programs interface suite[™] for Microsoft® windows, v 4.11.
 1147 United States Environmental Protection Agency, Washington, DC, USA.
 1148 https://www.epa.gov/tsca-screening-tools/epi-suitetm-estimation-program-interface
- 1149 121. Mansouri K, Kleinstreuer N, Abdelaziz AM, et al (2020) CoMPARA: Collaborative Modeling
 1150 Project for Androgen Receptor Activity. Environ Health Perspect 128:027002.
 1151 https://doi.org/10.1289/EHP5580
- 1152 122. Zheng Z, Arp HPH, Peters G, Andersson PL (2021) Combining *In Silico* Tools with
 1153 Multicriteria Analysis for Alternatives Assessment of Hazardous Chemicals: Accounting for
 1154 the Transformation Products of decaBDE and Its Alternatives. Environ Sci Technol
 1155 55:1088–1098. https://doi.org/10.1021/acs.est.0c02593
- 1156 123. Schymanski EL, Bolton EE (2022) FAIRifying the exposome journal: Templates for chemical structures and transformations. Exposome 2:osab006. https://doi.org/10.1093/exposome/osab006
- 1159 124. Schymanski E, Bolton E, Cheng T, et al (2023) Transformations in PubChem Full Dataset. 1160 https://doi.org/105281/zenodo8117741
- 1161 125. Schymanski EL, Zhang J, Thiessen PA, et al (2023) Per- and Polyfluoroalkyl Substances (PFAS) in PubChem: 7 Million and Growing. Environ Sci Technol 57, 44, 16918–1632 16928:acs.est.3c04855. https://doi.org/10.1021/acs.est.3c04855

- 1164 126. Shi Z-Q, Liu Y-S, Xiong Q, et al (2019) Occurrence, toxicity and transformation of six typical benzotriazoles in the environment: A review. Science of The Total Environment 661:407–1166 421. https://doi.org/10.1016/j.scitotenv.2019.01.138
- 1167 127. Cheng T, Zhao Y, Li X, et al (2007) Computation of Octanol-Water Partition Coefficients 1168 by Guiding an Additive Model with Knowledge. J Chem Inf Model 47:2140–2148. 1169 https://doi.org/10.1021/ci700257y
- 1170 128. Wang R, Fu Y, Lai L (1997) A New Atom-Additive Method for Calculating Partition Coefficients. J Chem Inf Comput Sci 37:615–621. https://doi.org/10.1021/ci960169p
- 1172 129. Meylan WM, Howard PH (1995) Atom/Fragment Contribution Method for Estimating Octanol–Water Partition Coefficients. Journal of Pharmaceutical Sciences 84:83–92. https://doi.org/10.1002/jps.2600840120
- 130. Card ML, Gomez-Alvarez V, Lee W-H, et al (2017) History of EPI Suite[™] and future perspectives on chemical property estimation in US Toxic Substances Control Act new chemical risk assessments. Environmental Science: Processes & Impacts 19:203–212. https://doi.org/10.1039/C7EM00064B
- 131. Cassano A, Manganaro A, Martin T, et al (2010) CAESAR models for developmental toxicity. Chemistry Central Journal 4:S4. https://doi.org/10.1186/1752-153X-4-S1-S4
- 132. Honma M (2020) An assessment of mutagenicity of chemical substances by (quantitative) structure—activity relationship. Genes and Environ 42:23. https://doi.org/10.1186/s41021-020-00163-1
- 1184 133. Schymanski EL, Kondić T, Neumann S, et al (2021) Empowering large chemical knowledge 1185 bases for exposomics: PubChemLite meets MetFrag. J Cheminform 13:19. 1186 https://doi.org/10.1186/s13321-021-00489-0
- 134. Wang Y, Ning X, Li G, Sang N (2022) New insights into potential estrogen agonistic activity
 of triazole fungicides and coupled metabolic disturbance. Journal of Hazardous Materials
 424:127479. https://doi.org/10.1016/j.jhazmat.2021.127479
- 1190 135. Goss K-U, Arp HPH, Bronner G, Niederer C (2009) Nonadditive effects in the partitioning behavior of various aliphatic and aromatic molecules. Environmental Toxicology and Chemistry 28:52–60. https://doi.org/10.1897/08-189.1
- 1193 136. Villeneuve DL, Coady K, Escher BI, et al (2019) High-throughput screening and environmental risk assessment: State of the science and emerging applications.

 1195 Environmental Toxicology and Chemistry 38:12–26. https://doi.org/10.1002/etc.4315
- 1196 137. Hollender J, Schymanski EL, Ahrens L, et al (2023) NORMAN guidance on suspect and non-target screening in environmental monitoring. Environmental Sciences Europe 35:75. https://doi.org/10.1186/s12302-023-00779-4
- 1199 138. Cramer GM, Ford RA, Hall RL (1976) Estimation of toxic hazard—A decision tree approach.
 1200 Food and Cosmetics Toxicology 16:255–276. https://doi.org/10.1016/S00151201 6264(76)80522-6

- 1202 139. EFSA Scientific Committee, More SJ, Bampidis V, et al (2019) Guidance on the use of the Threshold of Toxicological Concern approach in food safety assessment. EFS2 17:. https://doi.org/10.2903/j.efsa.2019.5708
- 1205 140. Wohlleben W, Mehling A, Landsiedel R (2023) Lessons Learned from the Grouping of Chemicals to Assess Risks to Human Health. Angew Chem Int Ed 62:e202210651. https://doi.org/10.1002/anie.202210651
- 1208 141. Zhu H, Zhang J, Kim MT, et al (2014) Big Data in Chemical Toxicity Research: The Use of High-Throughput Screening Assays To Identify Potential Toxicants. Chem Res Toxicol 27:1643–1651. https://doi.org/10.1021/tx500145h
- 1211 142. Enoch SJ, Roberts DW (2011) Approaches for Grouping Chemicals into Categories. https://doi.org/10.1039/9781849731744-00030
- 1213 143. Ankley GT, Bennett RS, Erickson RJ, et al (2010) Adverse outcome pathways: A 1214 conceptual framework to support ecotoxicology research and risk assessment. Enviro 1215 Toxic and Chemistry 29:730–741. https://doi.org/10.1002/etc.34
- 1216 144. Vinken M (2013) The adverse outcome pathway concept: A pragmatic tool in toxicology. 1217 Toxicology 312:158–165. https://doi.org/10.1016/j.tox.2013.08.011
- 1218 145. Rodríguez-Carrillo A, Rosenmai AK, Mustieles V, et al (2021) Assessment of chemical mixtures using biomarkers of combined biological activity: A screening study in human placentas. Reproductive Toxicology 100:143–154. https://doi.org/10.1016/j.reprotox.2021.01.002
- 1222 146. Zare Jeddi M, Hopf NB, Viegas S, et al (2021) Towards a systematic use of effect 1223 biomarkers in population and occupational biomonitoring. Environment International 1224 146:106257. https://doi.org/10.1016/j.envint.2020.106257
- 147. Hansson SO (2013) Chapter 9 ALARA: What is Reasonably Achievable? In: Oughton D,
 1226 Hansson SO (eds) Radioactivity in the Environment. Elsevier, pp 143–155