

Working Group XI - High-redshift Galaxies

Daniel Ceverino

Working Group XI – 28 Participants:

Oscar Agertz, Peter Behroozi, Daniel Ceverino, Jun-Hwan Choi, Romeel Dave, Maria Emilia De Rossi, Fabrice Durier, Jan Engels, Michele Fumagalli, Nick Gnedin, Oleg Gnedin, Javiera Guedes, Jason Jaacks, Dusan Keres, Ji-hoon Kim, Andrey Kravtsov, Mike Kuhlen, Sam Leitner, Piero Madau, Ken Nagamine, Brian O'Shea, Joel Primack, Brant Robertson, Emilio Romano-Diaz, Sijing Shen, Robert Thompson, Matteo Tomassetti, John Wise.

Code representatives

- All codes are well represented, with the exception of GADGET-SPHS


Data needed for high- z

- $M_h=10^{11}, M^{12}, M^{13} \text{ Msun}$
- Minimum number of snapshots:
200 timesteps equally spaced in expansion parameter ($da=0.005$) + redshift snapshots at **$z = 6, 3, 2, 1, 0.5, 0.2, 0.0$** at the very least).
- Do we want to address questions about reionization?

Working Group XI - High-redshift Galaxies: Cold flows, clumpiness, kinematics, ...


- Broad field
- Highly biased set of scientific questions

Cold streams




Nelson et al. (2013)

Clumpiness



Joel's slice

kinematics



Förster Schreiber et al (2009)

ART-I

runs without Cooling



Standard refinement



AGORA refinement

thanks