

Taylor polynomials and Maple

Ex 1. Find first and second degree Taylor polynomials
a) expanded at $x_0=0$.

$$P_n(x) = \sum_{i=0}^n \frac{f^{(i)}(x_0)}{i!} (x-x_0)^i$$

1. $f(x) = e^x$, $x \in \mathbb{R}$ $f'(x) = e^x$, $e^0 = 1$

$$P_1(x) = 1 + x$$

$$P_2(x) = 1 + x + \frac{x^2}{2}$$

2. $f(x) = \cos(x)$, $x \in \mathbb{R}$ $f'(x) = -\sin(x)$, $-\sin(0) = 0$

$$f''(x) = -\cos(x), \quad -\cos(0) = -1$$

$$P_1(x) = 1$$

$$P_2(x) = 1 - \frac{x^2}{2}$$

3. $f(x) = e^{\sin(x)}$, $x \in \mathbb{R}$ $f'(x) = \cos(x) \cdot e^{\sin(x)}$, $f'(0) = 1$

$$f''(x) = -\sin(x) \cdot e^{\sin(x)} + \cos^2(x) \cdot e^{\sin(x)}, \quad f''(0) = 1$$

$$P_1(x) = 1 + x$$

$$P_2(x) = 1 + x + \frac{x^2}{2}$$

b) Let $x_0 = 1$ and $f(x) = \frac{1}{x}$, $x > 0$.

$$P_1(x) = 1 - (x-1) = 2 - x$$

$$f'(x) = -\frac{1}{x^2}, \quad f'(1) = -1$$

$$P_2(x) = 1 - (x-1) + \frac{2}{2}(x-1)^2$$

$$f''(x) = \frac{2}{x^3}, \quad f''(1) = 2$$

$$= 2 - x + x^2 - 2x + 1 = x^2 - 3x + 3$$

Ex.2 Try different prompts with Maple.

```
> 2+2;
```

4


Simple calculations

```
> diff(sin(x), x);
```


$\cos(x)$

Function taking 2 inputs


```
> plot(sin(x), x=0..5);
```



```
> plot(sin(x), x=0..5, scaling=constrained);
```




```
> plot([sin(x), x^2], x=0..5, y=-2..2, color=[red, blue], scaling=constrained);
```




Plotting has rather many options. For now it's plenty to plot several graphs or using display to show several graphs together.

Ex3. Plot the functions from Ex1.


```
> plot([exp(x), ntaylor(exp(x), x=0, 2), ntaylor(exp(x), x=0, 3)]);
```




```
> plot([cos(x), ntaylor(cos(x), x=0, 2), ntaylor(cos(x), x=0, 3)], x=-2*Pi..2*Pi, y=-2..2, scaling=constrained);
```



```
> plot([exp(sin(x)), ntaylor(exp(sin(x)), x=0, 2), ntaylor(exp(sin(x)), x=0, 3)], x=-2*Pi..2*Pi, y=-5..5, scaling=constrained);
```




```
> plot([1/x, ntaylor(1/x, x=1, 2), ntaylor(1/x, x=1, 3)]);
```



Ex4. Approximate $f(x) = \sin(x)$ and see how well P_9 does expanded at $x_0 = 0$.

```
> f:=x->sin(x);
P9:=unapply(taylor(f(x),x=0,10),x);
P17:=unapply(taylor(f(x),x=0,18),x);
```

$$P9 = x + x^3 - \frac{1}{6}x^5 + \frac{1}{120}x^7 - \frac{1}{5040}x^9 + \frac{1}{362880}x^{11} - \frac{1}{39916800}x^{13} + \frac{1}{625038080}x^{15} - \frac{1}{130767436800}x^{17} + \frac{1}{35504742099200}x^{19}$$

$$P17 = x + x^3 - \frac{1}{6}x^5 + \frac{1}{120}x^7 - \frac{1}{5040}x^9 + \frac{1}{362880}x^{11} - \frac{1}{39916800}x^{13} + \frac{1}{625038080}x^{15} - \frac{1}{130767436800}x^{17} + \frac{1}{35504742099200}x^{19}$$


We just use `mtaylor`.

P_9 is quite good around 0, but falls off around $\pm \frac{3\pi}{2}$. We see P_{17} approximates well on the entirety of $[-2\pi, 2\pi]$.

Ex5. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be given by $f(x) = \sqrt{2x-1}$.

a) Since $\text{Dom}(\sqrt{x}) = [0, \infty[$ we have

$$\begin{aligned} \sqrt{2x-1} = 0 &\Leftrightarrow 2x-1=0 \\ &\Leftrightarrow x = \frac{1}{2}, \end{aligned}$$

and so $\text{Dom}(f) = [\frac{1}{2}, \infty[$.

b) Determine P_3 of f at $x_0 = 1$.

$$f'(x) = \frac{1}{2\sqrt{2x-1}} \cdot 2 = \frac{1}{\sqrt{2x-1}}, \quad f'(1) = 1$$

$$\begin{aligned} f''(x) &= ((2x-1)^{-\frac{1}{2}})' = -\frac{1}{2} (2x-1)^{-\frac{3}{2}} \cdot 2 \\ &= -\frac{1}{(2x-1)^{\frac{3}{2}}}, \quad f''(1) = -1 \end{aligned}$$

$$f'''(x) = \left(-(2x-1)^{-\frac{3}{2}} \right)' = \frac{3}{2} (2x-1)^{-5/2} \cdot 2$$

$$= \frac{3}{(2x-1)^{5/2}}, \quad f'''(1) = 3$$

$$P_3(x) = 1 + (x-1) - \frac{1}{2} (x-1)^2 + \frac{3}{6} (x-1)^3$$

$$= x - \frac{1}{2} (x^2 - 2x + 1) + \frac{1}{2} (x^3 - 3x^2 + 3x - 1)$$

$$= x - \frac{1}{2} x^2 + x - \frac{1}{2} + \frac{1}{2} x^3 - \frac{3}{2} x^2 + \frac{3}{2} x - \frac{1}{2}$$

$$= -1 + \frac{7}{2} x - 2x^2 + \frac{1}{2} x^3$$

c) Determine R_3 and that the error at $x = \frac{3}{2}$ is at most $\frac{5}{2^7}$.

Using lemma 4.5 we get

$$R_3(x) = \frac{f^{(4)}(\xi)}{4!} (x-1)^4, \quad \xi \in]1, x[.$$

$$f^{(4)}(x) = -\frac{15}{(2x-1)^{7/2}} \quad (\text{just follow the pattern})$$

Thus we get

$$R_3(x) = -\frac{15}{4!(2\xi-1)^{7/2}} (x-1)^4 = -\frac{5}{8} \frac{1}{(2\xi-1)^{7/2}} (x-1)^4.$$

The error is at most

$$\left| R_3\left(\frac{3}{2}\right) \right| = \left| -\frac{5}{8} \frac{1}{(2\xi-1)^{7/2}} \left(\frac{3}{2}-1\right)^4 \right| \leq \frac{5}{8} \frac{1}{(2 \cdot 1 - 1)^{7/2}} \left(\frac{1}{2}\right)^4$$

$$= \frac{5}{8} \cdot 1 \cdot \frac{1}{2^4} = \frac{5}{2^7}.$$

Ex 6. Past exercises to complete with Maple.

```

a) > I^2;
      I^3;
      (-I)^4;
      (-I)^5;

b) > (-2+3*I)/I

c) > w:=1-I;
      abs(w);
      argument(w);
      abs(exp(w));
      argument(exp(w));

d) > exp(I*pi/2);
      3*exp(1+pi*I);

e) > solve(z^2+(2+2*I)*z-2*I=0,z);

f) > diff(t^2+I*exp(t),t);
      diff(1-I*t^5,t);
      diff(t^5-I,t);
      diff(3*exp(1+t),t);
      diff(1*exp(2+I*t+3*I*t),t);

```

$$\begin{aligned}
 & -1 \\
 & -I \\
 & 1 \\
 & -I \\
 & 3 + 2I \\
 & \sqrt{2} \\
 & -\frac{\pi}{4} \\
 & e \\
 & -1 \\
 & 1 \\
 & -3e \\
 & -1 - 1 + \sqrt{2} + 1\sqrt{2}, -1 - 1 - \sqrt{2} - 1\sqrt{2} \\
 & 2t + 1\cos(t) \\
 & 5t^4 \\
 & 5t^4 \\
 & 31e^{2t} \\
 & (-3 + 2I)e^{1+3It}
 \end{aligned}$$

Ex 7. Let $f(x) = 2 \cos(x) + i \sin(2x)$, $x \in \mathbb{R}$.

a) Determine P_3 for f expanded at $x_0 = 0$.

Let's use Maple, since we know differentiation works the same for complex functions.

```

> f:=x-> 2*cos(x)+I*sin(2*x);
> P3:=unapply(mtaylor(f(x),x=0,4),x);

```

$$P3 := x \mapsto 2 + 2 \cdot 1 x - x^2 - \frac{4 \cdot 1 x^3}{3}$$

b) Expand Q_3 at $x_1 = \frac{\pi}{2}$.

```


> Q3:=unapply(mtaylor(f(x),x=pi/2,4),x);
expand(Q3(x));

```

$$\begin{aligned}
 Q3 &= x \mapsto -(2 + 2i) \left(x - \frac{\pi}{2} \right) + \left(\frac{1}{3} + \frac{4i}{3} \right) \left(x - \frac{\pi}{2} \right)^2 \\
 & - 2ix - 2ix + \pi + 1i + \frac{x^3}{3} + \frac{4ix^3}{3} - \frac{\pi x^3}{2} - 2i\pi x^2 + \frac{x^2}{4} + 1ix^2 - \frac{x^2}{24} - \frac{1ix^2}{6}
 \end{aligned}$$

c) Why is P_3 smarter to use than Q_3 for approximating $f(1)$?

It has a significantly simpler form. Though we can see that Q_3 performs better at $x=1$, it is hard labour unless we delegate to Maple:



Ex 8. Let $f(x) = \ln(1+x)$.

- a) State the limit formula (4.8) for f at $x_0 = 0$ of degrees 1, 2 and 3.

$$f'(x) = \frac{1}{1+x}, \quad f''(x) = -\frac{1}{(1+x)^2}, \quad f'''(x) = \frac{2}{(1+x)^3}$$

deg 1: $f(x) = x + x \cdot \varepsilon(x)$

deg 2: $f(x) = x - \frac{1}{2}x^2 + x^2 \cdot \varepsilon(x)$

deg 3: $f(x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 + x^3 \cdot \varepsilon(x)$

- b) Which result from a) can't be used to determine

$$\lim_{x \rightarrow 0} \frac{\ln(1+x) - x}{x^2} ?$$

What is the limit?

First degree is insufficient: $\lim_{x \rightarrow 0} \frac{x + x \cdot \varepsilon(x) - x}{x^2} = \lim_{x \rightarrow 0} \frac{\varepsilon(x)}{x}$

Second degree and above is fine:

$$\lim_{x \rightarrow 0} \frac{x - \frac{1}{2}x^2 + x^2 \varepsilon(x) - x}{x^2} = \lim_{x \rightarrow 0} \left(-\frac{1}{2} + \varepsilon(x) \right) = -\frac{1}{2}.$$

c) Compute $\lim_{x \rightarrow 0} \frac{x(e^x + 1) - 2(e^x - 1)}{x^3}$

We need to have a degree 3 expansion to deal with the denominator.

Numerator:	$x e^x + x - 2e^x + 2$	$x = 0$
		0
1	$e^x + x e^x + 1 - 2e^x$	0
2	$e^x + e^x + x e^x - 2e^x = x e^x$	0
3	$e^x + x e^x$	1

Thus $x(e^x + 1) - 2(e^x - 1) = \frac{1}{6} x^3 + x^3 \cdot \varepsilon(x)$,
as all lower degree terms vanish.

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{x(e^x + 1) - 2(e^x - 1)}{x^3} &= \lim_{x \rightarrow 0} \frac{\frac{1}{6} x^3 + x^3 \cdot \varepsilon(x)}{x^3} \\ &= \lim_{x \rightarrow 0} \left(\frac{1}{6} + \varepsilon(x) \right) = \frac{1}{6}. \end{aligned}$$

d) Use $f(x) = P_1(x) + R_1(x)$ to compute b) more gracefully.

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\ln(1+x) - x}{x^2} &= \lim_{x \rightarrow 0} \frac{P_1(x) + R_1(x) - x}{x^2} \\ &= \lim_{x \rightarrow 0} \frac{x - \frac{1}{2(1+\xi)^2} \cdot x^2 - x}{x^2} = \lim_{x \rightarrow 0} -\frac{1}{2(1+\xi)^2} = -\frac{1}{2}. \end{aligned}$$

Since $\xi \in]0, x[$ it follows that $\xi \rightarrow 0$ as $x \rightarrow 0$.