

Test case: 2D Lennard-Jones fluid versus lammps output

Morten Ledum

 github.com/mortele

October 26, 2018




FIG. 1. Positions read in by the subroutine `read_state_lammps` in the `file_writer` module, after being dumped by `dump (...) x y z (...)` in `lammps.in` input script. Difference between LAMMPS output and the imported Fortran positions shown—should be zero to machine precision, which it is.




FIG. 2. Positions after a single time step at $\Delta t = 0.001$, as compared to the LAMMPS trajectory positions.

Positions

Loading initial positions from a LAMMPS output file gives equal positions to within machine precision, c.f. Fig. 1. After a single time step at $\Delta t = 0.001$, positions deviate from the LAMMPS trajectories by between 10^{-15} and 10^{-16} , as seen in Fig. 2.

Note that the first 100 points represent the x coordinate of all $N = 100$ particles, the subsequent 100 points represents the y coordinate, while the last 100 are the z component of the positions. Since we are working in 2D, the positions and forces in the z -direction should always vanish.

Forces

The initial forces—calculated at the exact positions of the LAMMPS particles—deviate from the LAMMPS values by up to 10^{-12} , c.f. Fig. 3.




FIG. 3




FIG. 4




FIG. 5

Energies

Total energies calculated deviate from




FIG. 6