



خدمتكم عبادة نتقرب بها الى الله

2021

# أسئلة سنوات ورقي فاينال تفاضل وتكامل 2



سافر البولي تكنك 2021



**Question 1** Choose the correct answer for the questions (1-18) and complete the table

|       |       |       |       |       |       |       |       |      |       |
|-------|-------|-------|-------|-------|-------|-------|-------|------|-------|
| Q(1)  | Q(2)  | Q(3)  | Q(4)  | Q(5)  | Q(6)  | Q(7)  | Q(8)  | Q(9) | Q(10) |
|       |       |       |       |       |       |       |       |      |       |
| Q(11) | Q(12) | Q(13) | Q(14) | Q(15) | Q(16) | Q(17) | Q(18) |      |       |
|       |       |       |       |       |       |       |       |      |       |

**Q(1)** By converting from cartesian to polar, the cartesian point  $(x, y) = (-3, 0)$  becomes

- a)  $(13, \pi)$       b)  $(3, \pi \pm 2n\pi)$       c)  $(-3, \pi \pm 2n\pi)$       d)  $(16, \pi \pm 2n\pi)$

**Q(2)** Converting from polar to cartesian, the polar point  $(r, \theta) = \left(6, \frac{2\pi}{3}\right)$  becomes

- a)  $(3, 3)$       b)  $(-3, 3\sqrt{3})$       c)  $(-3, 3)$       d)  $((-\sqrt{3}, 3\sqrt{3}))$

**Q(3)** In Polar system,  $4 \cos \theta = 3$  defines a

- a) Horizontal line      b) Vertical Line      c) Circle      d) Square

**Q(4)** The sequence  $\left\{\frac{n^2}{2n^2+3}\right\}$  converges to

- a)  $\frac{3}{2}$       b)  $-\frac{1}{2}$       c)  $\frac{5}{2}$       d)  $\frac{1}{2}$

**Q(5)** By telescoping method we get  $\sum_{k=1}^{\infty} \frac{2}{k(k+1)} =$

- a)  $\frac{1}{4}$       b)  $\frac{1}{3}$       c)  $\frac{1}{2}$       d) 2

**Q(6)** The series  $\sum_{k=1}^{\infty} \frac{1}{\sqrt{2k}}$

- a) converges conditionally      b) converges absolutely      c) diverges      d) None

**Q(7)** The series  $\sum_{k=1}^{\infty} \frac{k^3 + 2k + 22}{2k^8 + 1}$

- a) converges      b) diverges      c) absolutely diverges      d) None

**Q(8)** The Maclaurin series for  $\sin x^3$  is

$$Q(9) \sum_{k=1}^{\infty} (\ln(2k+1))^k$$

- a) converges   b) converges conditionally   c) diverges   d) diverges absolutely

$$Q(10) \text{ The value of the improper integral } \int_1^2 \frac{3}{1-x} dx \text{ is}$$

- a) 1   b) 2   c) 6   d) diverges

$$Q(11) \text{ The value of the derivative } \frac{d}{dx} x^x \text{ at 2 is}$$

- a) 4   b)  $1 + \ln 4$    c)  $4 + \ln 4$    d)  $4 + 4 \ln 4$

$$Q(12) \lim_{x \rightarrow 0} (1 + 2x)^{\frac{1}{x}} =$$

- a)  $e^6$    b) 6   c) -6   d)  $e^{-6}$

$$Q(13) \lim_{x \rightarrow \infty} \frac{x^{11}}{e^{2x}} =$$

- a) 1   b) 0   c) 4   d) 3

$$Q(14) \lim_{x \rightarrow \infty} \left( 2 (\ln x)^{\frac{1}{2}} \right) =$$

- a) 1   b) 2   c)  $\ln 2$    d)  $\ln 3$

$$Q(15) \lim_{x \rightarrow 0} (e^{2x} + x)^{\frac{1}{x}} =$$

- a) 3   b) 4   c)  $e$    d)  $e^2$

$$Q(16) \int_1^{\sqrt{3}} \frac{4dx}{x^2 \sqrt{4-x^2}} dx$$

- a)  $\sqrt{3} - 2$    b)  $2\sqrt{3} - 1$    c)  $\sqrt{3} - 1$    d)  $\sqrt{3} - 3$

$$Q(17) \int_0^{\pi} 40 \sin 7x \cos 3x dx =$$

- a) -17   b) -7   c) 0   d) -71

$$Q(18) \int_0^{\ln 3} x e^x dx =$$

- a)  $3 \ln 3 - 2$    b)  $3 \ln 3$    c)  $\ln 3 - 2$    d)  $5 \ln 3 - 2$

$$\text{Question 2 Find the value of the integral } \int_1^{\infty} \frac{dx}{x^{12}} =$$

$$\text{Question 3 Find the value of the integral } \int_{-1}^0 \frac{dx}{x^2 + x - 2} = \frac{1}{2}$$

Q<sub>1</sub>)  $r^2 = x^2 + y^2$


$$r^2 = (-3)^2 + 0^2$$

$$r^2 = 9$$

$$r = 3$$

$$\theta = \pi \pm 2n\pi$$

$$(3, \pi \pm 2n\pi). \quad \textcircled{b}$$



1

Q<sub>2</sub>)  $x = r \cos \theta = 6 \cos \frac{2\pi}{3} = 6 \cdot \left(-\frac{1}{2}\right) = -3$

$$y = r \sin \theta = 6 \sin \frac{2\pi}{3} = 6 \cdot \frac{1}{2} = 3$$

$$(-3, 3) \quad \textcircled{c}$$

Q4)  $\lim_{n \rightarrow \infty} \frac{n^2}{2n^2+3} = \lim_{n \rightarrow \infty} \frac{n^2}{2n^2}$   
 $= \frac{1}{2} \quad (d)$

(2)

Q5)  $\frac{A}{k} + \frac{B}{(k+1)} = \frac{2}{k(k+1)}$

$$2 = A(k+1) + Bk$$

If  $k=0 \rightarrow A=2$

$k=-1 \rightarrow 2 = -B$   
 $B = -2$

$$S_n = \sum_{i=1}^n \left( \frac{2}{i} - \frac{2}{i+1} \right) = 2 \sum_{i=1}^n \left( \frac{1}{i} - \frac{1}{i+1} \right)$$

$$S_n = 2 \left[ \left( 1 - \frac{1}{2} \right) + \left( \frac{1}{2} - \frac{1}{3} \right) + \left( \frac{1}{3} - \frac{1}{4} \right) + \dots + \left( \frac{1}{n-1} - \frac{1}{n} \right) + \left( \frac{1}{n} - \frac{1}{n+1} \right) \right]$$

$$S_n = 2 \left( 1 - \frac{1}{n+1} \right)$$

$$\lim_{n \rightarrow \infty} S_n = 2 \left( 1 - \frac{1}{\infty} \right)$$

$$= 2 \cdot 1$$

$$= 2 \quad (d)$$



$$6) \sum_1^{\infty} \frac{1}{\sqrt{2} \sqrt{k}} = \sum \frac{1}{\sqrt{2} k^{1/2}} \quad p = \frac{1}{2} < 1$$

Div.  
(c)

(3)

$$7) b_k = \frac{k^3}{k^8} = \frac{1}{k^5}$$

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{a_k}{b_k} &= \lim_{k \rightarrow \infty} \frac{k^3 + 2k + 22}{2k^8 + 1} \cdot \frac{k^5}{1} \\ &= \lim_{k \rightarrow \infty} \frac{k^8 + 2k^6 + 22k^5}{2k^8} \\ &= \lim_{k \rightarrow \infty} \frac{\cancel{k^8} + 2\cancel{k^6} + 22\cancel{k^5}}{2\cancel{k^8}} \\ &= \frac{1}{2} < 1 \end{aligned}$$

$$\Rightarrow \sum b_k = \sum \frac{1}{k^5} \quad p = 5 > 1$$

Conu.  
(a)

$$\begin{aligned} Q8) \sin x^3 &= \sum_{k=0}^{\infty} \frac{(-1)^k (x^3)^{2k+1}}{(2k+1)!} \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k x^{6k+3}}{(2k+1)!} \end{aligned}$$

Q9)  $\sum_{k=1}^{\infty} \frac{1}{(\ln(2k+1))^k}$  Root.

(4)

$$\lim_{k \rightarrow \infty} \left( \frac{1}{\ln(2k+1)} \right)^{\frac{1}{k}}$$

$$= \lim_{k \rightarrow \infty} \frac{1}{\ln(2k+1)} = \frac{1}{\infty} = 0 < 1$$

Conv. (a)

Q10)  $\lim_{x \rightarrow 1^+} \int_m^2 \frac{3}{1-x} dx = \lim_{m \rightarrow 1^+} 3 \ln|1-x| \Big|_m^2$

$$= \lim_{m \rightarrow 1^+} [3 \ln|1-2| - 3 \ln|1-m|]$$

$$= 0 - 3(-\infty)$$

$$= \infty$$

Div. (d)

Q11)  $y = x^x$

$$\ln y = \ln x^x$$

$$\ln y = x \ln x$$

$$\frac{y'}{y} = x \cdot \frac{1}{x} + \ln x \cdot 1$$

$$y' = y(1 + \ln x)$$

$$y' = x^x (1 + \ln x)$$

$$y'(2) = 2^2 (1 + \ln 2)$$

$$= 4(1 + \ln 2) = 4 + 4 \ln 2$$

موجوده

(5)

$$Q_{12}) \lim_{x \rightarrow 0} (1 + 2x)^{\frac{-3}{x}}$$

$$\text{let } x = \frac{1}{y}$$

$$\begin{matrix} x \rightarrow 0 \\ y \rightarrow \infty \end{matrix}$$

$$\lim_{y \rightarrow \infty} \left(1 + \frac{2}{y}\right)^{-3y} = e^{2 \cdot (-3)} = e^{-6} \quad (d)$$

$$Q_{13}) \lim_{x \rightarrow \infty} \frac{x^{11}}{e^{2x}} = \text{لویٹال}$$

$$\text{Finally } \lim_{x \rightarrow \infty} \frac{11!}{2^{11} e^{2x}} = \frac{11!}{\infty} = 0 \quad (b)$$

$$Q_{14}) \lim_{x \rightarrow \infty} (2(\ln x)^{\frac{1}{x}})$$

$$2 \lim_{x \rightarrow \infty} (\ln x)^{\frac{1}{x}}$$

$$= 2 \cdot 1$$

$$= 2 \quad (b)$$

$$y = (\ln x)^{\frac{1}{x}}$$

$$\ln y = \ln (\ln x)^{\frac{1}{x}}$$

$$\ln y = \frac{\ln(\ln x)}{x}$$

$$\ln y = \lim_{x \rightarrow \infty} \frac{\ln(\ln x)}{x} \quad \frac{\infty}{\infty}$$

$$\ln y = \text{L'H} \lim_{x \rightarrow \infty} \frac{\frac{1}{x}}{\frac{1}{\ln x}}$$

$$\ln y = \lim_{x \rightarrow \infty} \frac{1}{x \ln x}$$

$$\ln y = 0$$

$$y = e^0$$

$$y = 1$$



$$\text{Q15) } \lim_{x \rightarrow 0} (e^{2x} + x)^{\frac{1}{x}} =$$

$$= e^2$$

$$y = (e^{2x} + x)^{\frac{1}{x}} \quad (6)$$

$$\ln y = \ln (e^{2x} + x)^{\frac{1}{x}}$$

$$\ln y = \frac{\ln(e^{2x} + x)}{x} \quad \frac{\infty}{\infty}$$

$$\ln y = \lim_{x \rightarrow \infty} \frac{\ln(e^{2x} + x)}{x}$$

$$\ln y \stackrel{L'H}{=} \lim_{x \rightarrow \infty} \frac{\frac{2e^{2x} + 1}{e^{2x} + x}}{1} \quad \frac{\infty}{\infty}$$

$$\ln y \stackrel{L'H}{=} \lim_{x \rightarrow \infty} \frac{4e^{2x}}{2e^{2x} + 1} \quad \frac{\infty}{\infty}$$

$$\ln y \stackrel{L'H}{=} \lim_{x \rightarrow \infty} \frac{8e^{2x}}{4e^{2x}}$$

$$\ln y = 2$$

$$y = e^2$$

Q16)  $x = 2\sin\theta$

$$x^2 = 4\sin^2\theta$$

$$\sqrt{4-x^2} = \sqrt{4-4\sin^2\theta} = 2\sqrt{1-\sin^2\theta} = 2\cos\theta$$

$$dx = 2\cos\theta d\theta$$

$$\text{If } x=1 \rightarrow 1 = 2\sin\theta \rightarrow \frac{1}{2} = \sin\theta \rightarrow \theta = \frac{\pi}{6}$$

$$x = \sqrt{2} \rightarrow \sqrt{2} = 2\sin\theta \rightarrow \frac{1}{\sqrt{2}} = \sin\theta \rightarrow \theta = \frac{\pi}{4}$$

$$\int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \frac{4 \cdot 2\cos\theta d\theta}{4\sin^2\theta \cdot 2\cos\theta} = \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \csc^2\theta d\theta$$

$$= -\cot\theta \Big|_{\frac{\pi}{6}}^{\frac{\pi}{4}}$$

$$= -1 + \sqrt{3}$$

$$= \sqrt{3} - 1 \quad (C)$$

Q17)  $\int_0^{\pi} 40 \sin 7x \cos 3x dx$

$$40 \times \frac{1}{2} \int_0^{\pi} \sin(4x) + \sin(10x) dx$$

$$20 \left( -\frac{\cos 4x}{4} + \frac{\cos 10x}{10} \right) \Big|_0^{\pi}$$

$$= 20 \left[ \left( -\frac{1}{4} - \frac{1}{10} \right) - \left( -\frac{1}{4} - \frac{1}{10} \right) \right]$$

$$= 20 \left( \cancel{-\frac{1}{4}} - \cancel{\frac{1}{10}} + \cancel{\frac{1}{4}} + \cancel{\frac{1}{10}} \right) = 0 \quad (C)$$

Q18)  $\int_0^{\ln 3} x e^x dx$

8

$$\begin{array}{rcl} x & & e^x \\ & \searrow + & \\ & & e^x \\ & \swarrow - & \\ 1 & & e^x \\ & \searrow - & \\ & & e^x \\ & \swarrow + & \\ 0 & & 0 \end{array}$$

$$= x e^x - e^x \Big|_0^{\ln 3}$$

$$= (\ln 3 \cdot e^{\ln 3} - e^{\ln 3}) - (0 - e^0)$$

$$= 3 \ln 3 - 3 + 1$$

$$= 3 \ln 3 - 2 \quad (a)$$

Q2)  $\lim_{m \rightarrow \infty} \int_1^m x^{-12} dx = \lim_{m \rightarrow \infty} \frac{x^{-11}}{-11} \Big|_1^m$

$$= \lim_{m \rightarrow \infty} \frac{-1}{11 x^{11}} \Big|_1^m$$

$$= \lim_{m \rightarrow \infty} \frac{-1}{11 m^{11}} + \frac{1}{11}$$

$$= \frac{-1}{\infty} + \frac{1}{11}$$

$$= \frac{1}{11}$$

$$Q_3) \int_{-1}^0 \frac{dx}{x^2 + x - 2}$$

9

$$\frac{1}{(x+2)(x-1)} = \frac{A}{(x+2)} + \frac{B}{(x-1)}$$

$$1 = A(x-1) + B(x+2)$$

$$\text{If } x=1 \rightarrow 1 = 3B \rightarrow B = \frac{1}{3}$$

$$x=-2 \rightarrow 1 = -3A \rightarrow A = -\frac{1}{3}$$

$$\int_{-1}^0 \frac{-\frac{1}{3}}{x+2} + \frac{\frac{1}{3}}{x-1} dx$$

$$-\frac{1}{3} \int_{-1}^0 \frac{1}{x+2} - \frac{1}{x-1} dx$$

$$-\frac{1}{3} \left( \ln(x+2) - \ln(x-1) \right) \Big|_{-1}^0$$

$$-\frac{1}{3} \ln \left| \frac{x+2}{x-1} \right| \Big|_{-1}^0$$

$$= -\frac{1}{3} \ln|2| + \frac{1}{3} \ln \left| \frac{1}{2} \right|$$