
Approximate Hyperparameter Marginalisation for Gaussian Processes

Anonymous Author(s)

Affiliation

Address

email

Abstract

Enter Abstract here

1 Introduction

2 Gaussian Processes

Gaussian processes (GPs) constitute a powerful method for performing Bayesian inference about functions using a limited set of observations [1]. A GP is defined as a distribution over the functions $f : \mathcal{X} \rightarrow \mathbb{R}$ such that the distribution over the possible function values on any finite set of \mathcal{X} is multivariate Gaussian. Given some arbitrary size n dataset, the observations $\mathbf{y} = \{y_1, \dots, y_n\}$ could be viewed as a single point sampled from a n -variate Gaussian distribution and can be partnered with a GP.

A GP is completely defined by its first and second moments: a mean function $\mu : \mathcal{X} \rightarrow \mathbb{R}$ which describes the overall trend of the function, and a symmetric positive semidefinite covariance function $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ which describes how function values are correlated as a function of their locations in the domain. Given a function $f : \mathcal{X} \rightarrow \mathbb{R}$ and a set of input points $\mathbf{x} \subset \mathcal{X}$, the Gaussian process prior distribution over the function values $\mathbf{f} = f(\mathbf{x})$ is given by:

$$p(\mathbf{f}|\mathbf{x}, \boldsymbol{\theta}) = \mathcal{N}(\mathbf{f}; \mu(\mathbf{x}, \boldsymbol{\theta}), K(\mathbf{x}, \mathbf{x}, \boldsymbol{\theta})) \quad (1)$$

$$= \frac{1}{\sqrt{\det 2\pi K_{\mathbf{f}}}} \exp\left(-\frac{1}{2}(\mathbf{f} - \mu_{\mathbf{f}})^{\top} K_{\mathbf{f}}^{-1}(\mathbf{f} - \mu_{\mathbf{f}})\right) \quad (2)$$

where $\boldsymbol{\theta}$ is a vector containing any parameters required by μ and K , and which form the *hyperparameters* of the model. There exists a wide variety of mean and covariance functions which can be chosen in order to reflect any prior knowledge available about the function of interest.

3 Approximate Hyperparameter Marginalisation

3.1 Proof of Positive Semi-Definiteness

A sum of kernels is itself a kernel, which by definition fulfils the necessary condition of positive semi-definiteness. Therefore:

$$K = \int k(\beta)p(\beta)d\beta \quad (3)$$

054
055
056
057
058
059
060
061
062
063
064
065
066
067
068
069
070
071
072
073
074
075
076
077
078
079
080
081
082
083
084
085
086
087
088
089
090
091
092
093
094
095
096
097
098
099
100
101
102
103
104
105
106
107

4 Experiments

5 Related Work

6 Conclusion

Acknowledgments

Do we have any? Aladdin / Orchid?

References

- [1] C. E. Rasmussen and C. K. I. Williams. *Gaussian Processes for Machine Learning*. MIT Press Cambridge, MA, USA, 2006.