

---

# Approximate Hyperparameter Marginalisation for Gaussian Processes

---

Anonymous Author(s)

Affiliation

Address

email

## Abstract

Enter Abstract here

## 1 Introduction

## 2 Gaussian Processes

Gaussian processes (GPs) constitute a powerful method for performing Bayesian inference about functions using a limited set of observations [1]. A GP is defined as a distribution over the functions  $f : \mathcal{X} \rightarrow \mathbb{R}$  such that the distribution over the possible function values on any finite set of  $\mathcal{X}$  is multivariate Gaussian. Given some arbitrary size  $n$  dataset, the observations  $\mathbf{y} = \{y_1, \dots, y_n\}$  could be viewed as a single point sampled from a  $n$ -variate Gaussian distribution and can be partnered with a GP.

A GP is completely defined by its first and second moments: a mean function  $\mu : \mathcal{X} \rightarrow \mathbb{R}$  which describes the overall trend of the function, and a symmetric positive semidefinite covariance function  $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  which describes how function values are correlated as a function of their locations in the domain. Given a function  $f : \mathcal{X} \rightarrow \mathbb{R}$  and a set of input points  $\mathbf{x} \subset \mathcal{X}$ , the Gaussian process prior distribution over the function values  $\mathbf{f} = f(\mathbf{x})$  is given by:

$$p(\mathbf{f}|\mathbf{x}, \boldsymbol{\theta}) = \mathcal{N}(\mathbf{f}; \mu(\mathbf{x}, \boldsymbol{\theta}), K(\mathbf{x}, \mathbf{x}, \boldsymbol{\theta})) \quad (1)$$

$$= \frac{1}{\sqrt{\det 2\pi K_{\mathbf{f}}}} \exp\left(-\frac{1}{2}(\mathbf{f} - \mu_{\mathbf{f}})^{\top} K_{\mathbf{f}}^{-1}(\mathbf{f} - \mu_{\mathbf{f}})\right) \quad (2)$$

where  $\boldsymbol{\theta}$  is a vector containing any parameters required by  $\mu$  and  $K$ , and which form the *hyperparameters* of the model. There exists a wide variety of mean and covariance functions which can be chosen in order to reflect any prior knowledge available about the function of interest.

## 3 Approximate Hyperparameter Marginalisation

### 3.1 Proof of Positive Semi-Definiteness

A sum of kernels is itself a kernel, which by definition fulfils the necessary condition of positive semi-definiteness. Therefore:

$$K = \int k(\beta)p(\beta)d\beta \quad (3)$$

054  
055  
056  
057  
058  
059  
060  
061  
062  
063  
064  
065  
066  
067  
068  
069  
070  
071  
072  
073  
074  
075  
076  
077  
078  
079  
080  
081  
082  
083  
084  
085  
086  
087  
088  
089  
090  
091  
092  
093  
094  
095  
096  
097  
098  
099  
100  
101  
102  
103  
104  
105  
106  
107

## **4 Experiments**

## **5 Related Work**

## **6 Conclusion**

### **Acknowledgments**

Do we have any? Aladdin / Orchid?

### **References**

- [1] C. E. Rasmussen and C. K. I. Williams. *Gaussian Processes for Machine Learning*. MIT Press Cambridge, MA, USA, 2006.