

LinesSlope of a line:

Let  $P = (x_1, y_1)$  and  $Q = (x_2, y_2)$  be two distinct points.

If  $x_1 \neq x_2$ , the slope  $m$  of the nonvertical line  $L$  containing  $P$  and  $Q$  is defined by the formula


$$m = \frac{y_2 - y_1}{x_2 - x_1} \quad x_1 \neq x_2$$

If  $x_1 = x_2$ ,  $L$  is a vertical line and the slope  $m$  is undefined.

In other words, slope is the angle a line makes with  $x$ -axis.


$$m = \tan \theta$$

If a line is vertical, then the angle the line will make with  $x$ -axis will be  $\pi/2$  and  $\tan \frac{\pi}{2}$  is undefined. So the slope is undefined.



Slope of  $L$  is

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$



Slope is undefined,  
 $L$  is vertical

$$\text{Slope, } m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{\text{change in } y}{\text{change in } x} = \frac{\Delta y}{\Delta x}$$

The expression  $\frac{\Delta y}{\Delta x}$  is called the average rate of change of  $y$  with respect to  $x$ .

Example:

Find the slope of the line containing points  $(1, 2)$  and  $(5, -3)$

Soln.: Here  $P = (1, 2)$   $Q = (5, -3)$

$$x_1 = 1, y_1 = 2, x_2 = 5, y_2 = -3$$

$$\therefore m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 - 2}{5 - 1} = \frac{-5}{4}$$

If  $x$  increases by 4 units the  $y$  will decrease by 5 units. The average rate of change of  $y$  with respect to  $x$  is  $-\frac{5}{4}$ .

Example:

Compute slopes of the lines

$$L_1 : P = (2, 3) \quad Q_1 = (-1, -2)$$

$$L_2 : P = (2, 3) \quad Q_2 = (3, -1)$$

$$L_3 : P = (2, 3) \quad Q_3 = (5, 3)$$

$$L_4 : P = (2, 3) \quad Q_4 = (2, 5)$$


### Example: 3

Draw a graph of the line that contains the point  $(3, 2)$  and has a slope of (a)  $\frac{3}{4}$  (b)  $-\frac{4}{5}$

Soln:

$$(a) \text{ Slope} = \frac{\text{Rise}}{\text{Run}} = \frac{3}{4} = \frac{\Delta y}{\Delta x}$$


The point given is  $(3, 2)$



### Equation of a vertical line:

A vertical line is given by the equation of the form  $x=a$ .

where  $a$  is the  $x$ -intercept.




### Example:

Graph the equation  $x=3$

Solution:


The graph of a equation  $x=3$  is a vertical line with  $x$ -intercept 3 and undefined slope.



## # Point-slope form of a line:

An equation of a nonvertical line with slope  $m$  that contains the point  $(x_1, y_1)$  is

$$y - y_1 = m(x - x_1)$$



### Example:

An equation of the line with slope 4 and containing the point  $(1, 2)$  can be found by using the point-slope form with  $m=4$ ,  $x_1=1$ , and  $y_1=2$ .

$$\begin{aligned} y - y_1 &= m(x - x_1) \\ \Rightarrow y - 2 &= 4(x - 1) \\ \Rightarrow y - 2 &= 4x - 4 \\ \Rightarrow y &= 4x - 4 + 2 \\ \Rightarrow y &= 4x - 2 \end{aligned}$$


### Exercise:

1. Find an equation of the line with slope 3 and containing point  $(-2, 3)$ .
2. Find an equation of the line with slope 2 and containing point  $(4, -3)$ .

## Equation of a Horizontal Line:

A horizontal line is given by an equation of the form  $y = b$

where  $b$  is the  $y$ -intercept.



### Example:

Find the equation of the horizontal line containing point  $(3, 2)$

### Soln:


All the  $y$ -values are equal on a horizontal line, the slope of a horizontal line is 0.

Here we have  $m=0$ ,  $x_1=3$ ,  $y_1=2$ .

$$\text{So } y - y_1 = m(x - x_1)$$

$$\Rightarrow y - 2 = 0(x - 3)$$

$$\Rightarrow y - 2 = 0 \Rightarrow y = 2$$



### Exercises

1. Find the equation of the horizontal line containing Point  $(-3, 2)$
2. Find the equation of the vertical line containing Point  $(4, -5)$

# Find the equation of a line given two points:

Find an equation of line containing the points  $(2, 3)$  and  $(-4, 5)$ . Graph the line.

Solution:

First compute the slope of the line;

$$m = \frac{y_2 - y_1}{x_2 - x_1} \quad \text{Hence, } x_1 = 2, y_1 = 3 \\ x_2 = -4, y_2 = 5$$

$$\Rightarrow m = \frac{5 - 3}{-4 - 2} = \frac{2}{-6} = -\frac{1}{3}$$


Use the point  $(2, 3)$  and the slope  $m = -\frac{1}{3}$  to get the point-slope form of the equation of the line,

$$y - 3 = -\frac{1}{3}(x - 2)$$

$$\Rightarrow y - 3 = -\frac{1}{3}x + \frac{2}{3}$$

$$\Rightarrow y = -\frac{1}{3}x + \frac{2}{3} + 3$$

$$\Rightarrow y = -\frac{1}{3}x + \frac{11}{3}$$




# Slope-intercept form of an eqn of line:

An equation of a line with slope  $m$  and

$y$ -intercept  $b$  is

$$y = mx + b$$

Hence  $m$  is the slope and  $b$  is the  $y$ -intercept.



For example:

$y = -2x + 7$ , if we compare this equation to  $y = mx + b$   
we get  $m = -2$  and  $b = 7$ .

so the slope of the line is  $-2$  and its y intercept is  $7$ .

Example:

Find the slope and y intercept  $b$  of the equation

$2x + 4y = 8$ . Graph the equation.

Solution:

To obtain the slope and y intercept, write the equation in slope-intercept form by solving for  $y$ :

$$2x + 4y = 8$$


$$\Rightarrow 4y = -2x + 8$$

$$\Rightarrow y = -\frac{1}{2}x + 2$$

Compare this equation with  $y = mx + b$ , we got the  
slope  $m = -\frac{1}{2}$  and y-intercept  $b = 2$

Exercise:

- Find the slope and y intercept of the  
equation  $x + 2y = 4$ ,  $2x - 3y = 6$ ,  $y = -3x + 4$



# The equation of a line is in general form when it is written as

$$Ax + By = C$$

where  $A$ ,  $B$  and  $C$  are real numbers and  $A$  and  $B$  are not both zero.

If  $B=0$  and  $A \neq 0$  then  $Ax = C$  or  $x = \frac{C}{A}$  is a equation of vertical line. Ex:  $x = a \Rightarrow 1 \cdot x + 0 \cdot y = a$

If  $A=0$  and  $B \neq 0$  then  $By = C$  or  $y = \frac{C}{B}$  is a equation of horizontal line. Ex:  $y = b \Rightarrow 0 \cdot x + 1 \cdot y = b$

If  $B \neq 0$  and  $A \neq 0$  then we can solve the equation for  $y$  and write the equation in-slope-intercept form.

Example:

Graph the equation  $2x + 4y = 8$  by finding its intercepts.

Solution:

To obtain  $x$ -intercept, let  $y=0$  in the eqn and solve for  $x$ .

$$2x + 4y = 8$$

$$\Rightarrow 2x + 4 \cdot 0 = 8$$

$$\Rightarrow 2x = 8 \Rightarrow x = 4$$

To obtain  $y$ -intercept, let  $x=0$  in the eqn and solve for  $y$ ,

$$2x + 4y = 8$$

$$\Rightarrow 2 \cdot 0 + 4y = 8$$


$$\Rightarrow 4y = 8 \Rightarrow y = 2$$

$\therefore$  The  $x$ -intercept is 4 and the point  $(4,0)$  is on the graph of the equation.

The  $y$ -intercept is 2 and the point  $(0,2)$  is on the graph of the equation.


Plot the points  $(4,0)$  and  $(0,2)$

and draw the line through the points.



# Find equations of Parallel lines:

Two non-vertical lines are parallel if and only if their slopes are equal and they have different  $y$ -intercepts. i.e  $m_1 = m_2$



Example:

Show that the lines given by the following equations are parallel.

$$L_1: 2x + 3y = 6 \quad ; \quad L_2: 4x + 6y = 0$$

Soln: To determine whether these lines have equal slopes and different  $y$ -intercepts, write each equation in slope-intercept form:

$$L_1: 2x + 3y = 6$$

$$\Rightarrow 3y = -2x + 6$$

$$\Rightarrow y = -\frac{2}{3}x + 2$$

$$\text{Slope} = -\frac{2}{3}, y\text{-intercept} = 2$$

$$L_2: 4x + 6y = 0$$


$$\Rightarrow 6y = -4x$$

$$\Rightarrow y = -\frac{4}{6}x$$

$$\Rightarrow y = -\frac{2}{3}x$$

$$\text{Slope} = -\frac{2}{3}, y\text{-intercept} = 0$$

Since these lines have the same slope  $-\frac{2}{3}$  but different y-intercepts, the lines are parallel.



Example:

Find an equation for the line that contains the point  $(2, -3)$  and is parallel to the line  $2x+y=6$ .

Solution:

Since the two lines to be parallel, the slope of the line that we seek equals the slope of the line  $2x+y=6$ . Begin by writing the eqn of the line  $2x+y=6$  in the slope-intercept form,

$$2x+y=6 \Rightarrow y=-2x+6$$

The slope is  $-2$ . Since the line that we seek also has slope  $-2$  and contains the point  $(2, -3)$ , use the point-slope form to obtain its equation,

$$y-y_1 = m(x-x_1)$$

$$\Rightarrow y-(-3) = -2(x-2)$$

$$\Rightarrow y+3 = -2x+4$$

$$\Rightarrow y = -2x+1$$

$$\Rightarrow 2x+y = 1$$


This is the line parallel to the line  $2x+y=6$  and contains the point  $(2, -3)$

## # Find Equations of perpendicular lines:

Two non-vertical lines are perpendicular if and only if the product of their slope is  $-1$ .

$$\text{i.e } m_1 \cdot m_2 = -1$$

$$\text{or } m_1 = -\frac{1}{m_2} \text{ or } m_2 = -\frac{1}{m_1}$$



If a line has slope  $\frac{3}{2}$ , any line having slope  $-\frac{2}{3}$  is perpendicular to it.

### Example:

Find an equation of the line that contains the point  $(1, -2)$  and is perpendicular to the line  $x+3y=6$ .

Graph the two lines.

### Solution:

First write the equation of the given line in slope intercept form to find its slope.

$$x+3y=6 \Rightarrow 3y = -x+6 \\ \Rightarrow y = -\frac{1}{3}x + 2$$

Compare this equation with  $y = mx+b$  we get  $m = -\frac{1}{3}$ .

Any line perpendicular to this line will have

$$\text{slope } m_1 \cdot m_2 = -1$$

$$\Rightarrow m_2 = -\frac{1}{m_1}$$

$$\Rightarrow m_2 = -\frac{1}{(-\frac{1}{3})} = (-1) \cdot (-3) = 3$$


We require the point  $(1, -2)$  to be on the line with slope 3, we use the point-slope form of the equation of a line.

$$y - y_1 = m(x - x_1)$$

$$\Rightarrow y - (-2) = 3(x - 1)$$

$$\Rightarrow y + 2 = 3x - 3$$

$$\Rightarrow 3x - y = 5$$



Exercise:

Find an equation of line that is perpendicular to the line  $y = \frac{1}{2}x + 4$  and contain point  $(1, -2)$ .

## Summary of the Chapter 2

1. Slope,  $m = \frac{y_2 - y_1}{x_2 - x_1} = -\frac{y_1 - y_2}{x_1 - x_2}$ ;  $x_1 \neq x_2$

2. Equation of a vertical line:  $x=a$

where  $a$  is the  $x$ -intercept.  
slope undefined.

3. Equation of a horizontal line:  $y=b$

where  $b$  is the  $y$ -intercept, slope is zero.

4. Point-slope form of an equation of a line:

$$y - y_1 = m(x - x_1)$$

where  $m$  is the slope of the line and  $(x_1, y_1)$  point contains lies on the line.

5. Slope-intercept form of an equation of a line:

$$y = mx + b$$

where  $m$  is the slope and  $b$  is the  $y$ -intercept.

6. General form of equation:

$$Ax + By = C$$

If  $B=0$  then  $Ax=C$  vertical line. If  $A=0$  then  $By=C$  horizontal line.

7. Two lines are parallel if their slopes are equal.

$$\text{i.e. } m_1 = m_2$$

8. Two lines are perpendicular if their slopes are  
perpendicular i.e  $m_1 m_2 = -1$

$$\text{or, } m_1 = -\frac{1}{m_2} \text{ or } m_2 = -\frac{1}{m_1}$$

### Exercise:

1. Graph plot each pair of points and determine the slope containing them. Graph the line

i.  $(-2, 3); (2, 1)$  ii)  $(-1, 2); (-1, -2)$  iii)  $(4, 2); (-5, 2)$

2. Graph the line containing the point  $P$  and having slope  $m$

i.  $P = (1, 2); m = 3$  ii)  $P = (0, 3)$ ; slope undefined  
iii)  $P = (2, -4); m = 0$  iv)  $P = (1, 3); m = -\frac{2}{5}$

3. i) Find an equation of the line containing the points  $(-1, 2)$  and  $(-1, -2)$ . Graph the line.

ii) Find an equation of the line containing the points  $(2, 0)$  and  $(2, 2)$ . Graph the line.

4. Find the slope and  $y$ -intercept of each line. Graph the line.

i)  $x + y = 1$  ii)  $x = 2$  iii)  $y = \frac{1}{2}x + 2$  iv)  $y = 5$

5. Find the intercepts of the graph of each equation.

i)  $x - \frac{2}{3}y = 4$  ii)  $7x + 2y = 21$  iii)  $\frac{1}{2}x + \frac{1}{3}y = 1$

6. The equations of two lines are given, determine if the lines are parallel, perpendicular or neither.

i)  $y = 2x - 3$  ii)  $y = 4x + 5$  iii)  $y = -2x + 3$   
 $y = 2x + 4$   $y = -4x + 2$   $y = -\frac{1}{2}x + 2$

7. Find an eqn perpendicular to the line  $y = 2x - 3$  containing the point  $(1, -2)$ .


## Circles

A circle is a set of points in the  $xy$ -plane that are a fixed distance  $r$  from a fixed point  $(h, k)$ . The fixed distance  $r$  is called the radius and the fixed point  $(h, k)$  is called the center of the circle.

Let  $(x, y)$  represents the coordinates of any points on a circle with radius  $r$  and center  $(h, k)$ . Then the distance between the points  $(x, y)$  and  $(h, k)$  must always equal to  $r$ . Thus by distance formula:

$$\sqrt{(x-h)^2 + (y-k)^2} = r$$

$$\Rightarrow (x-h)^2 + (y-k)^2 = r^2$$




# The standard form of an equation of circle with radius  $r$  and center  $(h, k)$  is

$$(x-h)^2 + (y-k)^2 = r^2$$


# The standard form of an equation of a circle with radius  $r$  and center at the origin  $(0, 0)$  is

$$x^2 + y^2 = r^2$$



# If the radius  $r=1$ , then the circle whose center is at origin is called unit circle and has the equation

$$x^2 + y^2 = 1$$



Example:

Write the standard form of an equation of the circle with radius 5 and center  $(-3, 6)$ .

Solution: Hence radius  $r=5$ ,  $h=-3$ ,  $k=6$ .

we have,  $(x-h)^2 + (y-k)^2 = r^2$

$$\Rightarrow [x - (-3)]^2 + (y - 6)^2 = 5^2$$

$$\Rightarrow (x+3)^2 + (y-6)^2 = 25$$

Example:

Graph the equation of circle.

$$(x+3)^2 + (y-2)^2 = 16$$

Solution:

Given

$$(x+3)^2 + (y-2)^2 = 16$$


$$\Rightarrow [x - (-3)]^2 + (y-2)^2 = 4^2$$

Compare with the standard form of the eqn of circle

$$(x-h)^2 + (y-k)^2 = r^2$$

Get  $h = -3$ ,  $k = 2$  and  $r = 4$

So the circle has center  $(-3, 2)$  and radius 4.



Exercise:

1. Write the standard form of an equation of the circle with radius 8 and center (4, 2).
2. Graph the equation of the circle  
$$(x+2)^2 + (y+1)^2 = 4$$

Example:

For the circle  $(x+3)^2 + (y-2)^2 = 16$ , Find the intercepts.

Solution:

To find x-intercepts let  $y=0$ , then

$$(x+3)^2 + (0-2)^2 = 16$$

$$\Rightarrow (x+3)^2 + (0-2)^2 = 16$$

$$\Rightarrow (x+3)^2 + 4 = 16$$

$$\Rightarrow (x+3)^2 = 12$$

$$\Rightarrow x+3 = \pm\sqrt{12}$$

$$\Rightarrow x = -3 \pm \sqrt{12}$$

To find y-intercepts, let  $x=0$ , then

$$(0+3)^2 + (y-2)^2 = 16$$

$$\Rightarrow 9 + (y-2)^2 = 16$$

$$\Rightarrow (y-2)^2 = 7$$

$$\Rightarrow y-2 = \pm\sqrt{7}$$

$$\Rightarrow y = 2 \pm \sqrt{7}$$

The x-intercepts are  $-3-\sqrt{12}$  and  $-3+\sqrt{12}$

The y-intercepts are  $2-\sqrt{7}$  and  $2+\sqrt{7}$

# General form of the equation of a circle:

When its graph is a circle, the equation

$$x^2 + y^2 + ax + by + c = 0$$

is referred to as the general form of the equation of a circle.

\* \* This eqn has a graph of a circle or a point or has no graph at all.

For example  $x^2 + y^2 = 0$  is the single point  $(0,0)$ .

The equation  $x^2 + y^2 + 5 = 0$  has no graph, because sum of squares of real numbers are never negative.

Problem:

Graph the equation

$$x^2 + y^2 + 4x - 6y + 12 = 0$$

Soln:

Group the terms involving  $x$ , group the term involving  $y$  and put the constant on the right hand side of the equation. The result is

$$(x^2 + 4x) + (y^2 - 6y) = -12$$

$$\Rightarrow (x^2 + 4x + 4) + (y^2 - 6y + 9) = -12 + 4 + 9$$

$$\Rightarrow (x+2)^2 + (y-3)^2 = 1$$

$$\Rightarrow (x-(-2))^2 + (y-3)^2 = 1$$

This eqn is the standard form of the equation of a circle with radius 1 and center  $(-2, 3)$ .

1. Find the standard form of the equation and graph the circle

i)  $x^2 + y^2 + 4x - 4y - 1 = 0$       ii)  $2x^2 + 2y^2 - 12x + 8y - 24 = 0$

2. Find the intercepts, center, radius

i.  $2(x-3)^2 + 2y^2 = 8$       ii)  $x^2 + (y-1)^2 = 1$

### Exercise / Homework:

1. Write the standard form and the general form of the circle of radius  $r$  and origin  $(h, k)$ . Graph each circle.

i)  $r=2; (h, k)=(0, 2)$       ii)  $r=\frac{1}{2}; (h, k)=\left(\frac{1}{2}, 0\right)$

iii)  $r=4; (h, k)=(-2, 1)$

2. Find the center and radius of each circle, also find the intercepts.

i)  $2x^2 + 2y^2 + 8x + 7 = 0$       ii)  $x^2 + y^2 + 4x + 2y - 20 = 0$

3. Find the standard form of the equation of each circle;

i) center at the origin and containing the point  $(-2, 3)$

ii) center  $(2, 3)$  and tangent to the  $x$ -axis

iii) with endpoints of a diameter at  $(1, 4)$  and  $(-3, 2)$