

MAT361.4  
Final Assignment

S.M. Sajid Hasan Shanta  
ID: 1831238 642

(1)

Here, Probability of getting any side of number,  $P = \frac{1}{5}$

and, Since the spinner spun 5 times.

$$\therefore n = 5$$

It follows binomial distribution.

$\therefore$  Probability of getting at most two 5's,

$$\begin{aligned} P(X \leq 2) &= P(X=0) + P(X=1) + P(X=2) \\ &= \binom{5}{0} \left(\frac{1}{5}\right)^0 \left(1-\frac{1}{5}\right)^{5-0} + \binom{5}{1} \left(\frac{1}{5}\right)^1 \left(1-\frac{1}{5}\right)^{5-1} \\ &\quad + \binom{5}{2} \left(\frac{1}{5}\right)^2 \left(1-\frac{1}{5}\right)^{5-2} \\ &= 0.94 \end{aligned}$$

(2)

Solar power Plants failures occur with an average of 5 failures every year.

$$\therefore \lambda_y = 5$$

for a week,

$$\lambda_w = \frac{5}{\frac{365}{7}}$$

$$[\because 1 \text{ year} = 365 \text{ days} = \frac{365}{7} \text{ weeks}]$$

$$= 0.0959$$

$\therefore$  Probability that there will be more than one failure during a particular week,

$$\begin{aligned} P(X \geq 1) &= 1 - P(X=0) - P(X=1) \\ &= 1 - \frac{e^{-0.0959} (0.0959)^0}{0!} - \frac{e^{-0.0959} (0.0959)^1}{1!} \\ &= 1 - 0.909 - 0.087 \\ &= 0.004 \end{aligned}$$


(3)

Probability that a adult people height  
is greater than 184 cm,

we have,  
normally distributed,  
mean,  $E(x) = \mu = 185 \text{ cm}$   
variance,  $V(x) = \sigma^2 = 2 \text{ cm}$   
 $\therefore \sigma = \sqrt{2}$

$$\begin{aligned}
 &P(X > 184) \\
 &= P(184 < X < \infty) \\
 &= P\left(\frac{184 - \mu}{\sigma} < \frac{x - \mu}{\sigma} < \frac{\infty - \mu}{\sigma}\right) \\
 &= P\left(\frac{184 - 185}{\sqrt{2}} < Z < \infty\right) \\
 &= F(\infty) - F\left(\frac{184 - 185}{\sqrt{2}}\right) \\
 &= 1 - F(-0.707) \\
 &= 1 - F(-0.71) \\
 &= 1 - 0.2389 = 0.7611
 \end{aligned}$$

Figure:





(4)

$$\bar{x} = \frac{60 + 75 + 72 + 65 + 68}{5} = 68$$

$$\mu_0 = 70, n = 5$$

$$s^2 = \frac{\sum (x_i - \bar{x})^2}{n-1}$$

$$= \frac{(60-68)^2 + (75-68)^2 + (72-68)^2 + (65-68)^2 + (68-68)^2}{5-1}$$

$$= 34.5$$

$$\therefore \text{Test statistic is } \frac{68-70}{\sqrt{\frac{34.5}{5}}} = -0.76$$

$$\begin{aligned} \therefore \text{The rejection region is } & ]-\infty, -t_{\alpha}] \\ & = ]-\infty, -t_{0.05}] \\ & = ]-\infty, -2.132] \end{aligned} \quad \left| \begin{array}{l} \alpha = 0.05 \end{array} \right.$$

P.T.O

Comment:

$\therefore$  Test statistic value,  $-0.76$ , does not fall in the rejection region.

$\therefore$  We can not reject null hypothesis ( $H_0$ ).

So that, the researcher's assumption, the mean weight of the adult men in Bangladesh is less than 70kg is incorrect.

(5)

Blood samples of 5 people were sent to each of two laboratories (Lab 1 and 2) for cholesterol determinations.  
∴ it is matched paired t test.

$$H_0: \mu_D = 0$$

$$H_1: \mu_D < 0$$

where,

$$\mu_D = \mu_Y - \mu_X$$

here,  
 $\mu_X$  = The mean cholesterol <sup>reported</sup> by Lab 1  
 $\mu_Y$  = The mean cholesterol reported by Lab 2

$$\text{Test Statistic} = \frac{\bar{D}}{\sqrt{\frac{S_D^2}{n}}} \sim t_{n-1}$$

from the data we get,

Person (i)	$D_i = Y_i - X_i$
1	42
2	17
3	20
4	-38
5	16

$$\begin{aligned} \therefore \text{Sample mean difference, } \bar{D} &= \frac{42 + 17 + 20 - 38 + 16}{5} \\ &= 11.4 \end{aligned}$$



$$\therefore S_D^2 = \frac{\sum_{i=1}^5 (D_i - \bar{D})^2}{n-1}$$

$$= \frac{(42-11.4)^2 + (17-11.4)^2 + (20-11.4)^2 + (-38-11.4)^2 + (16-11.4)^2}{5-1}$$

$$= 875.8$$

and,  $n = 5$

$$\therefore \text{Test static} = \frac{11.4}{\sqrt{\frac{875.8}{5}}}$$

$$= 0.86$$

$\therefore$  Rejection region is:  $]-\infty, -t_{\alpha, n-1}]$

$$= ]-\infty, -t_{0.1, 4}]$$

$$= ]-\infty, -1.533]$$

$$\left\{ \begin{array}{l} \text{here} \\ \alpha = 0.1 \\ n-1 = 5-1 = 4 \end{array} \right.$$

Comment:

$\therefore$  Test statistics value (0.86) does not fall in the rejection region.

$\therefore$  We can not reject null hypothesis ( $H_0$ )

So that, the assumption, the (population) mean cholesterol levels by Lab 1 is greater than the (population) mean cholesterol mean levels reported by Lab 2 is incorrect.