

MS2STEP

Do you mind if I ask you a few questions, so I can get a better idea of what is best from my experience?

Can I ask who is it for? (Is this medication for you or someone else?)		
Has the pregnancy been confirmed by ultrasound?		(MUST be YES and MUST be IN UTERUS, not ectopic)
How long has your pregnancy been?		(MUST be UNDER 63 DAYS)
Do you currently have an IDU?		(MUST be NO)
Do you have?		
	Have a bleeding problem	(MUST be NO)
	Have a problem with your adrenal gland	(MUST be NO)
	Have taken corticosteroids for a long time, or take blood thinners	(MUST be NO)
	Are UNABLE to get emergency medical care in the next 14 days	(MUST be NO)

How do I take MS-2-Step?

To take MS-2 Step:

Step 1	Take one mifepristone tablet by swallowing it with a glass of water . It should be taken on an empty stomach 2 hours before or 2 hours after a meal.
Step 2	Take four misoprostol tablets at the same time, 36 to 48 hours after the mifepristone. Hold the 4 tablets in your mouth between the cheek and gum for 30 minutes before <u>swallowing</u> any tablet fragments with water. They should be taken on an empty stomach – 2 hours before or 2 hours after a meal.

If you forget to take the step-2 tablets (misoprostol) and it is **more than 48 hours** after you have taken step-1 tablet (mifepristone), **contact your doctor immediately**.

What can I expect after taking MS-2 Step?

Vaginal bleeding

Step-1	After taking mifepristone, it is likely that you will be able to continue with your normal activities. VAGINAL BLEEDING usually starts <u>1 to 2 days after</u> taking the mifepristone tablet.
Step-2	After taking the misoprostol tablets, you should <u>stay at home and rest for 3 hours.</u> VAGINAL BLEEDING will occur usually <u>within 4 hours</u> and can range from light to heavy – it is usually more than your usual period. VAGINAL BLEEDING lasts on average for <u>10-16 days</u> , although it may last <u>up to 6 weeks</u> . If bleeding does NOT occur, contact your doctor straight away.

Cramping and pain may also occur and can be mild to severe.

You should stay **within a 2-hour drive of a hospital** for 14 days after starting MS-2 Step in case you have heavy bleeding or complications.

Before taking this medicine, I would like to inform you of the risks and the expected side effects if that is, ok?

Like all medicine, MS-2 Step can have side effects. These are usually minor and temporary and can include:	Diarrhea Headache Nausea and vomiting Dizziness Flushing Shivering or chills
Complications and risks associated with MS-2-step are rare but you need to <u>see your doctor and get checked out</u> if you have:	Have heavy bleeding filling more than 2 pads an hour, for 2 hours in a row Have large clots the size of your fist Have severe pain Feel weak or have nausea, vomiting or diarrhoea more than 24 hours after taking the misoprostol Have fevers or chills that are ongoing or more than 38 degrees Have smelly or abnormal vaginal discharge

You should arrange a follow-up appointment with your healthcare practitioner 14 to 21 days after your first dose of MS-2-Step to ensure that the termination was complete.