

Emergency contraception

Questions	Please answer here	Triggers for Pharmacist
General Information & Triage		
Who is this requested for? Are you filling this form out for yourself or someone else?	Myself / For	[Trigger: If third party cannot provide adequate history → Contact patient directly before supply.
Age: What is your age / date of birth?	 years old	(Triggers: If <18 years, assess Gillick competency/maturity) (if the patient does not have a mature understanding of what is proposed → Refer)
Current symptoms: Are you experiencing severe abdominal pain or unexpected heavy bleeding?	Yes / No	[Trigger: Red flag for acute pathology / existing ectopic pregnancy → Medical referral / investigation.]
Physical Characteristics		
Weight: What is your weight in kg?	 kg	BMI = kg/m²
Height: What is your height in cm? (Used to calculate BMI)	 cm	[Triggers: Weight >70 kg or BMI >26 kg/m ² directs product selection → (Ulipristal or Levonorgestrel double dose 3 mg preferred) ; Weight >85 kg or BMI >30 kg/m ² → requires Supply & Refer.]
Timing of Unprotected Sexual Intercourse (UPSI)		
Timing: How many hours or days ago did the unprotected sex occur?	 hours ago.	[Triggers: ≤96 hours → (Ulipristal or Levonorgestrel) ; 96–120 hours → (Ulipristal only); >120 hours → (Urgent Referral).]
Frequency: How many times have you had unprotected sex in this current menstrual cycle ?	 times	[Triggers: Multiple episodes increase the risk of pre-existing pregnancy from an earlier event.]
Menstrual History & Pregnancy Risk		
Average cycle length: How many days long is your normal menstrual cycle?	 days	
Last Menstrual Period (LMP): When was the <u>first day</u> of your last period?	On .../.../...	
Expected period: When is your next period expected?	On .../.../...	
Period characteristics: Was your last period normal in duration and flow, or was it lighter/shorter than usual?		[Triggers: Lighter/unusual flow indicates risk of pre-existing pregnancy.]
Overdue period: Is your current period late?	Late / Not late	[Triggers: Late period indicates risk of pre-existing pregnancy.]
Medication & Health History		
Current contraception: Are you currently using any hormonal contraception (e.g., pill, ring, patch, implant, injection, IUD)? If yes, which one?	Yes / No	[Triggers: Missed/delayed doses guide whether EC is required and influence product selection.]
Recent EC use: Have you taken emergency contraception in the past 7 days? If yes, which one?	Yes / No	[Triggers: Ulipristal in past 5 days requires Ulipristal; Levonorgestrel in past 7 days requires Levonorgestrel.] [ulipristal EC is required more than once in the same menstrual cycle (possible risk of hepatotoxicity → Supply & Refer.)]
Recent progestogens: Have you used any progestogen-containing medicines or contraceptives in the past 7 days?	Yes / No	[Triggers: Requires Levonorgestrel (must avoid Ulipristal).]
Other medications: Have you taken any prescription or herbal medicines in the past 4 weeks (e.g., epilepsy meds, St John's Wort, antibiotics like griseofulvin)?	Yes / No	[Triggers: CYP3A4 enzyme inducers require Levonorgestrel double dose (3 mg) or Copper IUD, plus Supply & Refer.]
Gastrointestinal conditions: Do you have Crohn's disease, severe diarrhea, or recent vomiting?	Yes / No	[Triggers: Malabsorption lowers efficacy (Supply & Refer.)]
Asthma: Do you have severe asthma that requires oral steroid tablets ?	Yes / No	[Triggers: If UPSI >96 hours ago, requires Urgent Referral (as ulipristal EC is may worsen their asthma); if ≤96 hours ago, use Levonorgestrel.]
Breastfeeding: Are you currently breastfeeding?	Yes / No	[Triggers: Levonorgestrel is 1st choice; Ulipristal requires express & discard milk counselling.]
Safety & Support		
Consent & Safety: Did this sexual intercourse occur with your full consent?	Yes / No	[Triggers: Suspected sexual assault requires offering support, 1800RESPECT contact, and Supply & Refer.]

Decision-Making & Product Selection Questions

Field / Question Area	Clinical Reason / Decision Implication
Menstrual History	Late period or lighter than usual: Indicates risk of pre-existing pregnancy.
Exact Time Since UPSI	0–96 hours: Ulipristal (1st choice), Levonorgestrel (2nd choice). / 96–120 hours: Ulipristal only. / >120 hours: Urgent referral.
Weight / BMI	>70 kg / BMI >26 kg/m ² : Copper IUD preferred; Ulipristal is 1st oral choice; Levonorgestrel double dose (3 mg) is 2nd choice.
Previous EC Use	Ulipristal in past 5 days: Use Ulipristal. / Levonorgestrel in past 7 days: Use Levonorgestrel.
Current Contraception	Progestogens in past 7 days / Resuming immediately: Use Levonorgestrel. (Ulipristal requires a 5-day delay before starting progestogens).
Enzyme-Inducing Meds	CYP3A4 inducers / Griseofulvin in past 4 weeks: Double dose Levonorgestrel (3 mg) only (or Copper IUD).
Severe Asthma	Severe asthma on oral steroids (UPSI ≤96h): Use Levonorgestrel.
Breastfeeding	Breastfeeding mothers: Levonorgestrel is 1st choice. (If Ulipristal is used, express & discard milk for 24h/1 week per guidelines).